

A ONE DIMENSIONAL MANIFOLD IS OF COHOMOLOGICAL DIMENSION 2

SATYA DEO¹

ABSTRACT. G. Bredon defines the cohomological Dimension of a topological space X to be the supremum of all cohomological ϕ -dimensions of X , where ϕ varies over the entire families of supports on X . He has proved that if X is a topological n -manifold then the cohomological Dimension of X is n or $n + 1$. He was not able to decide which one it is, even for a space as simple as the real line. The objective of this paper is to solve his problem for $n = 1$. In particular, we have shown that the cohomological Dimension of the real line is 2.

Let ϕ be a family of supports on a topological space X . Godement [2] defines the ϕ -dimension ($\dim_{\phi}(X)$) of X to be the largest integer n (or ∞) if there is a sheaf \mathcal{Q} of abelian groups on X for which the Grothendieck cohomology [3] group $H_{\phi}^n(X, \mathcal{Q}) \neq 0$. If ϕ runs over all those paracompactifying families of supports on X whose extents equal X , then $\dim_{\phi}(X)$ is independent of ϕ and is called the *cohomological dimension* ($\dim(X)$) of X . The class of all such spaces for which $\dim(X)$ has a meaning is large enough to include all locally paracompact Hausdorff spaces. The $\dim(X)$ of a n -manifold X is n and is dominated by the covering dimension [4] for paracompact Hausdorff X . Quite naturally, Bredon [1] defines the *cohomological Dimension* ($\text{Dim}(X)$) of X to be the $\text{Sup}_{\phi} \{ \dim_{\phi}(X) \}$ for all families of supports ϕ on X . Among other things, he proves that for a topological n -manifold X , $\text{Dim}(X) = n$ or $n + 1$ leaving it open which one it is. The present objective is to indicate its pathological or nonpathological nature by proving the following

Theorem. *Let X be a one-dimensional topological manifold. Then $\text{Dim}(X)$ is two.*

Recall that if \mathcal{Q} is a sheaf of abelian groups on a space X and $\mathcal{Q}^s(U)$ denotes the group of all serrations of \mathcal{Q} on U , then the sheaf generated by the presheaf $U \rightarrow \mathcal{Q}^s(U)/\mathcal{Q}(U)$ on X is denoted by $\mathcal{Z}^1(X, \mathcal{Q})$ and $\text{Dim}(X) \leq 1$ if and only if $\mathcal{Z}^1(X, \mathcal{Q})$ is flabby for every sheaf \mathcal{Q} on X [1, p. 110]. Consider \mathbf{R} as imbedded in the one-dimensional manifold X . Then the proof of the theorem follows from

Received by the editors July 16, 1974.

AMS (MOS) subject classifications (1970). Primary 55B30; Secondary 55C10.

Key words and phrases. Sheaves, sections and serrations of sheaves, extent of a family of supports, paracompactifying family of supports, manifolds and dimension of a topological space.

¹ This work was done while the author was a Visiting Member at the Tata Institute of Fundamental Research, Bombay.

Lemma. Let $A = \{0\} \cup \{1/m \mid m \in \mathbf{N}\}$ be the subset of X and $U = X - A$. Then the sheaf $\mathcal{Q} = \mathbf{Z}_U$ on X is such that $\mathcal{Z}^1(X, \mathcal{Q})$ is not flabby.

Proof. Notice that any section of \mathcal{Q} on any subset P of X such that $P \cap A \neq \emptyset$, is a zero section. Consider the set

$$\{f_m : (1/(m+2), 1/m) \rightarrow \mathcal{Q} \mid m \in \mathbf{N}\}$$

of serrations of \mathcal{Q} defined by

$$f_m(x) = \begin{cases} m, & x \neq 1/(m+1), \\ 0, & x = 1/(m+1). \end{cases}$$

These serrations give rise to a well-defined section S of $\mathcal{Z}^1(X, \mathcal{Q})$ on $(0, 1)$ because of the fact that $f_{m+1} - f_m$ is a section of \mathcal{Q} on their common domain $(1/(m+2), 1/(m+1))$. We claim that this section cannot be extended to any neighbourhood of the set $[0, 1)$ in X . For, suppose S^1 is an extension of S in a neighbourhood $(-\delta, 1)$ of $[0, 1)$ where $\delta > 0$, and let S^1 be represented by a serration g of \mathcal{Q} defined on some neighbourhood $(-\epsilon, \epsilon)$ of $0 \in X$ where $\delta > \epsilon > 0$. Since S and S^1 agree on $(0, \epsilon)$, $g - f_m$ must be a section of \mathcal{Q} for each m sufficiently large. But that means, by our remark at the outset, that there is a positive integer N such that $\forall m \geq N$, $g - f_m = 0$ on $(1/(m+2), 1/m)$. This means f_{N+k} and f_{N+k+1} must agree for each integer $k \geq 0$ on their common domains, i.e., $N = N+k$, $\forall k \geq 0$, a contradiction.

Remark. It seems to me that for a topological n -manifold X , $\text{Dim}(X) = n + 1$ is true for every n . However, the difficulty in extending the proof lies in our inability to recognise precisely the stalks and the topology of the sheaf $\mathcal{Z}^n(X, \mathcal{Q})$, $n \geq 1$, and for any given sheaf \mathcal{Q} on X .

REFERENCES

1. G. E. Bredon, *Sheaf theory*, McGraw-Hill, New York, 1967. MR 36 #4552.
2. R. Godement, *Topologie algébrique et théorie des faisceaux*, Hermann, Paris, 1958. MR 21 #1583.
3. A. Grothendieck, *Sur quelques points d'algèbre homologique*, Tôhoku Math J. (2) 9 (1957), 119–221. MR 21 #1328.
4. W. Hurewicz and H. Wallmann, *Dimension theory*, Princeton Math. Ser., vol. 4, Princeton Univ. Press, Princeton, N. J., 1948. MR 3, 312.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALLAHABAD, ALLAHABAD-211002, INDIA (Current address)

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, COLABA, BOMBAY 400 005, INDIA