

SEQUENTIAL WEAK CONVERGENCE IN THE BIDUAL OF A BANACH LATTICE

JOHN L. B. GAMLEN¹

ABSTRACT. Let X be a Banach lattice with separable dual. Then the bidual X'' is the closure of X under sequential monotone weak convergence.

This paper proves a special case of a conjecture relating two types of weak sequential closure in the bidual X'' of a Banach lattice X . Specifically, if Y is a sublattice of X'' , write Y^σ for the smallest subset of X'' which contains Y and is closed under sequential monotone $\sigma(X'', X')$ convergence. It turns out that Y^σ is a sublattice. We will study its relation with the space \tilde{Y} generated by sequential $\sigma(X'', X')$ convergence.

Conjecture. The spaces X^σ, \tilde{X} coincide.

We prove a special case below (Theorem 1). Incidentally, note that, for a more general sublattice Y , the spaces Y^σ, \tilde{Y} may not coincide. An example is with $X = L^1[0, 1]$. Let K be the maximal ideal space of $L^\infty[0, 1]$, and identify X'' with the regular measures on K . Take Y to be the atomic measures, noting that $Y^\sigma = Y$, but $\tilde{Y} = X''$.

Theorem 1. *If X is a Banach lattice with separable dual, and Y is a sublattice of X'' , then $Y^\sigma = \tilde{Y}$. In particular $X^\sigma = \tilde{X} = X''$.*

Another special case of the conjecture is known already, and is a deep result of T. Ogasawara [6] stating (essentially) that if X is a Banach lattice in which every monotone bounded sequence converges weakly, then X is weakly complete. This may be rephrased: "if $X^\sigma = X$, then $\tilde{X} = X''$ ".

The space $C[0, 1]$ is a nice nontrivial example of a Banach lattice in which the conjecture holds. In this case both X^σ, \tilde{X} may be identified with the space of bounded Borel-measurable functions on $[0, 1]$. It is easy to see that a sequence of such functions converges for $\sigma(C'', C')$ if and only if it is uniformly bounded and pointwise convergent.

If Y is a linear subspace of X'' , write Y^2 for the smallest subspace of X'' containing Y and closed under weak unconditional series convergence. It is clear that, in general,

Received by the editors March 26, 1973.

AMS (MOS) subject classifications (1970). Primary 46A40, 46B99.

Key words and phrases. Weak sequential convergence.

¹ The author is partially supported by N.R.C. Grant A7552 and NSF GP-30671.

$$(*) \quad X^\sigma \subseteq X^\Sigma \subseteq \tilde{X} \subseteq X''.$$

Note here that monotone weak limits may be expressed as limits of positive term series which converge unconditionally (weakly) because members of X' are differences of positive members. Thus Theorem 1 implies

Corollary 2. *For a Banach lattice with separable dual, equality holds in (*).*

This result can be used to give another proof of the following result of T. Ogasawara [6]:

Corollary 3. *If X is a Banach lattice with separable bidual then X is reflexive.*

Proof of Corollary 3. Since X'' is separable, it is clear that X cannot have a subspace isomorphic to c_0 , and thus by a theorem of Bessaga and Pelczynski [1], X^Σ must coincide with X . Corollary 2 now shows $X'' = X$. \square

One can also argue above using Ogasawara's first-mentioned result, instead of Corollary 2. (Since $X^\Sigma = X$, bounded monotone sequences converge weakly so X is weakly complete, but X' is separable, proving reflexivity.)

It is well known that Corollary 3 does not extend to general Banach spaces, as the famous counterexample of R. C. James [3] shows.

For expositions of the elementary Banach lattice theory we will use, the reader is referred to [2] or [9]. For further results in the spirit of Ogasawara's theorems, but in a more general setting, the reader is referred to the recent paper of L. Tzafriri [8].

The proof of Theorem 1. For simplicity we deal only with the case $Y = X$. Once we know that $X^\sigma = X''$, treatment of the general sublattice Y is just an imitation.

Lemma 4. *If X is a Banach lattice, then X^σ is a sublattice of X'' .*

Proof. The idea is the same as in the analogous result for Boolean algebras (the "monotone class" theorem).

Let $X_1 = \{x \text{ in } X'' : x \vee 0 \in X^\sigma\}$. A routine argument shows that X_1 is closed under monotone sequential $\sigma(X'', X')$ convergence. This means X_1 contains X^σ , which shows X^σ is a sublattice. \square

We introduce a new topology \mathcal{J} on X'' , defined by the seminorms p_ϕ , $\phi \in X'$ where $p_\phi(x) = \langle |x|, |\phi| \rangle$ for $x \in X''$. Incidentally, this is the "locally solid" topology generated by $\sigma(X'', X')$ in the sense of I. Namioka [5].

Lemma 5. (a) X is dense in (X^σ, \mathcal{J}) .

(b) The weak topology of (X^σ, \mathcal{J}) is $\sigma(X^\sigma, X')$.

Proof. From the definition, monotone convergence for \mathcal{J} and $\sigma(X'', X')$ coincide. This proves (a).

License or copyright restrictions may apply to redistribution; see <https://www.ams.org/journal-terms-of-use>

Clearly the members of X' are \mathcal{J} -continuous functionals. For the con-

verse note that each \mathcal{J} -continuous linear functional ψ on X has a restriction to X which is a member ϕ of X' . By part (a), ϕ and ψ coincide on X^σ . \square

The proof of Theorem 1 goes as follows. Let $\{\phi_n\}$ be a sequence of positive norm-one members of X' with dense span. Construct an increasing norm ρ on X^σ as follows.

Define

$$\rho(x) = \sum_1^\infty 2^{-n} \langle |x|, \phi_n \rangle \quad \text{for } x \in X''.$$

It is clear that ρ gives the same topology as \mathcal{J} on bounded subsets of X'' . Write L for the completion of X under the norm ρ . Not only is L a Banach lattice, but by Kakutani's theorem [4] L is isomorphic to some $L^1(\mu)$, and is thus weakly complete. We make two claims:

(1) the weak topology of L coincides on bounded subsets of X^σ with $\sigma(X^\sigma, X')$;

(2) the unit ball B in X^σ is ρ -complete, and hence by (1) sequentially complete for $\sigma(X^\sigma, X')$.

Obviously (1), (2) finish the proof of Theorem 1.

Proof of Claim (1). Since the topology of ρ is weaker than \mathcal{J} (on X''), the weak topology of L is weaker than the weak topology of (X^σ, \mathcal{J}) , which by Lemma 5 is $\sigma(X^\sigma, X')$ on X^σ . For the converse, note that the functionals ϕ_n are ρ -continuous, and on bounded sets they determine the topology $\sigma(X^\sigma, X')$. \square

Proof of Claim (2). Let $\{x_n\}$ be a ρ -Cauchy sequence in B , say $\{x_n\}$ converges in L to x_- . Since x_n^+ converges in L to x^+ , we may assume that x_n, x , are positive, and each x_n is in B . Write $y_k = \bigvee_{n=k}^\infty x_n$, which exists as a monotone $\sigma(X'', X')$ limit, and thus lies in X^σ . Let us assume that $\rho(x_n - x) < 2^{-n}$, all n . We claim that $\rho(y_k - x_k) < 2^{2-k}$, all k .

To prove this, write $y_k - x_k$ as $\bigwedge_{n=k}^\infty (x_n - x_k)$, to obtain:

$$\rho(y_k - x_k) \leq \rho\left(\sum_{n=k}^\infty |x_n - x_k|\right) < 2^{2-k}, \quad \text{as required.}$$

(The series on the right converges for ρ and for $\sigma(X'', X')$ to an element of X^σ . The fact that ρ is an increasing norm then yields the above inequality.)

Claim (2) now follows from the observation that $\{y_k\}$ is a monotone increasing sequence in B , so must converge for ρ and for $\sigma(X'', X')$ to a member of B which the above reasoning shows is x . \square

Remarks on the proof of Theorem 1. It might be thought that the Theorem could be proved directly by some variant of the above argument involving $\{y_k\}$. The trouble is that if x_k converges to x for $\sigma(X'', X')$, then x_k^+ need not converge to x^+ . (For example, in $L^2[0, \pi]$, $x_k(t) = \sin kt$.) The point of our proof is that ρ -convergence does not suffer from this defect.

License or copyright restrictions may apply to redistribution; see <https://www.ams.org/journal-terms-of-use>

It might also occur to the reader that a theorem of H. Schaefer (cf. A.

Peressini [7, p. 146]) concerning metrizable locally convex lattices could be modified to deal with our situation. This fails because the weak topology $\sigma(X'', X')$ is not metrizable, even though the bounded sets are metrizable.

Our results raise the question of which sequentially monotone complete topological vector lattices are already sequentially complete? The facts we have used about X^σ , $\sigma(X^\sigma, X')$ in proving its sequential completeness are:

- (a) $\sigma(X^\sigma, X')$ is the weak topology of a locally solid space;
- (b) the bounded sets are metrizable (for the weak topology, in this case $\sigma(X, X')$).

Unfortunately (a) may fail even for the topology $\sigma(X', X)$ of the dual X' of a Banach lattice X . (Take $X = C[0, 1]$. The locally solid topology generated by $\sigma(X', X)$ is the norm topology, whose weak topology is not $\sigma(X', X)$.)

REFERENCES

1. C. Bessaga and A. Pelczynski, *On bases and unconditional convergence of a series in Banach spaces*, *Studia. Math.* 17 (1958), 151–164. MR 22 #5872.
2. Garrett Birkhoff, *Lattice theory*, 3rd ed., Amer. Math. Soc. Colloq. Publ., vol. 25, Amer. Math. Soc., Providence, R. I., 1967. MR 37 #2638.
3. R. C. James, *A non-reflexive Banach space isometric with its second conjugate space*, *Proc. Nat. Acad. Sci. U.S.A.* 37 (1951), 174–177. MR 13, 356.
4. S. Kakutani, *Concrete representation of abstract (L)-spaces and the mean ergodic theorem*, *Ann. Math. (2)* 42 (1941), 523–537. MR 2, 318.
5. I. Namioka, *Partially ordered linear topological spaces*, *Mem. Amer. Math. Soc. No. 24* (1957). MR 20 #1193.
6. T. Ogasawara, *Theory of vector lattices. I*, *J. Sci. Hiroshima Univ. Ser. A* 12 (1942), 37–100. MR 10, 545.
7. A. Peressini, *Ordered topological vector spaces*, Harper and Row, New York, 1967. MR 37 #3315.
8. L. Tzafriri, *Reflexivity in Banach lattices and their subspaces*, *J. Functional Analysis* 10 (1972), 1–18.
9. A. C. Zaanen, and W. A. J. Luxemburg, *Riesz spaces*, North-Holland, London, 1971.

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CONNECTICUT 06520

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA, CANADA