

ARCS DEFINED BY ONE-PARAMETER SEMIGROUPS
OF OPERATORS IN BANACH SPACES WITH
THE RADON-NIKODYM PROPERTY

CHARLES L. BYRNE

ABSTRACT. It is shown that a recent theorem of Junghenn and Taam concerning the domain of the infinitesimal generator of a strongly continuous one-parameter semigroup of operators on a reflexive, locally convex topological vector space remains valid if the domain of the operators is a Banach space with the Radon-Nikodym property. A partial result is obtained for general Banach spaces.

The following theorem is proved in a recent paper by Junghenn and Taam:

Theorem. *Let X be a reflexive, locally convex topological vector space. Let $T(t)$, for $t \geq 0$, be a strongly continuous semigroup of operators on X , such that for all $c > 0$, $T(c)$ is an isomorphism (into), and let A be the infinitesimal generator of the semigroup. The following are equivalent:*

- (1) x is in the domain of A ;
- (2) $T(\cdot)x$ is absolutely continuous on the interval $[0, c]$ for any $c > 0$;
- (3) $T(\cdot)x$ is of bounded variation on the interval $[0, c]$ for any $c > 0$.

In the absence of reflexivity, each condition implies the next. It is the purpose of this note to show that if X is a Banach space, then (2) and (3) are equivalent, and if, in addition, X has the Radon-Nikodym property, then all three are equivalent. Among spaces with the Radon-Nikodym property are reflexive Banach spaces and separable dual Banach spaces. For a discussion of the Radon-Nikodym property, see the paper of Rieffel [2]. Recent results characterizing Banach spaces with this property are to be found in [4], [5], and [6].

To prove the above assertions we need two lemmas. Our notation is that of [1]. Terms not defined in this note are as used in that paper.

Lemma 1. *Let X be a Banach space, and $x \in X$, for which (3) holds. If $c > 0$ and L is the total variation of $T(\cdot)x$ on $[0, c]$, then $\|T(c+h)x - T(c)x\|/h \leq KL/(c-h)$ for all $0 < h < c$, where K is a constant not dependent upon the choice of h (but dependent upon c).*

Received by the editors October 4, 1974.

AMS (MOS) subject classifications (1970). Primary 46A05, 47D05; Secondary 46A25.

Proof. Choose n so that $nb \leq c < (n + 1)b$, and let $1 \leq j \leq n$. Pick K so that $\|T(t)x\| \leq K\|x\|$ for t in $[0, c]$. Then

$$\begin{aligned} \|T(c + b)x - T(c)x\| &= \|T(c + b - jb)\{T(jb)x - T(jb - b)x\}\| \\ &\leq K\|T(jb)x - T(jb - b)x\|. \end{aligned}$$

Therefore

$$n\|T(c + b)x - T(c)x\| \leq \sum_1^n K\|T(jb)x - T(jb - b)x\| \leq KL.$$

The assertion follows since we have $c - b < nb$.

Lemma 2. *Under the same assumptions as Lemma 1, the set $\{\|T(h)x - x\|/b \mid 0 < b \leq c\}$ is bounded.*

Proof. We have

$$\|T(h)x - x\|/b = \|T(c)^{-1}\{T(c + b)x - T(c)x\}\|/b \leq M\|T(c + b)x - T(c)x\|/b,$$

for $M = \|T(c)^{-1}\|$. Then $\|T(h)x - x\|/b \leq MKL/(c - b)$, and if $0 < b < c/2$ we have $\|T(h)x - x\|/b \leq 2MKL/c = Q$. Clearly, for b in $[c/2, c]$, the numbers are bounded.

We prove now that (3) implies (2) in any Banach space. Let $\epsilon > 0$ be given and let (a_i, b_i) , $i = 1, 2, \dots, n$, be any collection of nonoverlapping intervals in $[0, c]$. If we first assume that $b_i - a_i < c/2$, then

$$\|T(b_i)x - T(a_i)x\| = \|T(a_i)\{T(b_i - a_i)x - x\}\| \leq K\|T(b_i - a_i)x - x\|.$$

Therefore the choice of $d < \epsilon/QK$ gives $\sum_1^n \|T(b_i)x - T(a_i)x\| < \epsilon$ if $\sum_1^n (b_i - a_i) < d$.

Banach spaces X with the Radon-Nikodym property are precisely those spaces with the property that every function of bounded variation from the real line into X is differentiable almost everywhere. This is a classical result due to Bochner and Taylor [3]. If we assume that X has this property, then there is a $\bar{T} > 0$ at which point the function $T(\cdot)x$ is differentiable, if (3) holds. Then, as h approaches 0, $[T(\bar{T} + h)x - T(\bar{T})x]/h$ approaches a limit. It follows that $[T(h)x - x]/h$ does also, since it is equal to $[T(\bar{T})^{-1}\{T(\bar{T} + h)x - T(\bar{T})x\}]/h$. Therefore, if X is assumed to have the Radon-Nikodym property, the three conditions given in the Theorem are equivalent.

REFERENCES

1. H. Junghenn and C. T. Taam, *Arcs defined by one-parameter semigroups of operators*, Proc. Amer. Math. Soc. 44 (1974), 113–120.
2. M. A. Rieffel, *The Radon-Nikodym theorem for the Bochner integral*, Trans. Amer. Math. Soc. 131 (1968), 466–487. MR 36 #5297.

3. S. Bochner and A. E. Taylor, *Linear functionals on certain spaces of abstractly-valued functions*, *Ann. of Math. (2)* 39 (1938), 913–944.
4. W. J. Davis and R. R. Phelps, *The Radon-Nikodym property and dentable subsets in Banach spaces*, *Proc. Amer. Math. Soc.* 45 (1974), 119–122.
5. R. E. Huff, *Dentability and the Radon-Nikodym property*, *Duke Math. J.* 41 (1974), 111–114.
6. R. R. Phelps, *Dentability and extreme points in Banach spaces*, *J. Functional Analysis* 17 (1974), 78–90.

DEPARTMENT OF MATHEMATICS, THE CATHOLIC UNIVERSITY OF AMERICA,
WASHINGTON, D.C. 20064