

A NOTE ON CONTINUITY OF SEMIGROUPS OF MAPS

PAUL R. CHERNOFF¹

ABSTRACT. An example is given of a separately continuous semigroup of transformations on Hilbert space which fails to be jointly continuous at $t = 0$.

1. Let X be a topological space and $\{F_t: t \geq 0\}$ a one-parameter semigroup of continuous maps of X into itself; that is, $F_{s+t} = F_s \circ F_t$ and $F_0 = \text{identity}$. Suppose also that for each x in X the mapping $t \mapsto F_t(x)$ is continuous. Thus $(t, x) \mapsto F_t(x)$ is continuous in each variable separately. Under additional hypotheses we can conclude that this map is jointly continuous. Specifically, in [1] it is shown that if X is metrizable, then every point (t, x) with $t > 0$ is a point of joint continuity. Moreover, Dorroh [2] has shown that if X is locally compact and σ -compact then every point (t, x) , including points with $t = 0$, is a point of joint continuity.

On the other hand, it was stated in [1] that there is an example for which joint continuity fails at $t = 0$, with X a certain subset of R^2 . (There is a misprint in [1] on p. 1046; it is erroneously stated that such an example exists with $X = R^2$. This is, of course, ruled out by Dorroh's result.)

The aim of this note is to present an example illustrating failure of joint continuity at $t = 0$ with X a Hilbert space. Thus Dorroh's result does not generalize from finite-dimensional to infinite-dimensional manifolds.

2. Before giving the construction, it seems worthwhile to present a proof of joint continuity at $t = 0$ which is substantially more elementary than Dorroh's argument. We shall assume that the space X is locally compact and metrizable (rather than locally compact and σ -compact as in [2]).

Suppose that $\{F_t: t \geq 0\}$ is a separately continuous semigroup of maps on X . By [1] we have joint continuity for $t > 0$. We must establish joint continuity at $t = 0$. That is, given x in X and sequences $x_n \rightarrow x$, $t_n \rightarrow 0$, we have to show that $F_{t_n}(x_n) \rightarrow x$. If this is not the case, then there is a compact neighborhood K of x such that $F_{t_n}(x_n) \notin K$ for arbitrarily large n . Since $x_n \rightarrow x$, we may as well assume that $x_n \in K$, but $F_{t_n}(x_n) \notin K$, for all n . But then, because $F_t(x_n)$ is continuous in t , a connectedness argument

Received by the editors December 14, 1974.

AMS (MOS) subject classifications (1970). Primary 20M20, 54H15; Secondary 57A20.

¹ Research partially supported by NSF grant GP-30798X.

Copyright © 1975, American Mathematical Society

shows that $F_{s_n}(x_n)$ lies on the boundary B of K for some s_n between 0 and t_n . Since B is compact, we may assume that $F_{s_n}(x_n)$ converges to a point y in B . Now observe that if $t > 0$,

$$(1) \quad F_t(y) = \lim F_t F_{s_n}(x_n) = \lim F_{t+s_n}(x_n) = F_t(x),$$

where joint continuity at (t, x) is used at the last step. If we finally let t converge to 0 in (1), we conclude that $y = x$. But this is a contradiction, since x is in the interior of K .

3. To prepare the ground for the construction of the Hilbert space example, we will exhibit the example on the subset of R^2 which we mentioned above.

We define the subset X as follows. Let p be the point $(1, 0)$. Introducing polar coordinates in the usual way, let $A = \{(r, \theta) : r > 1, 0 < \theta < 2\pi\}$. Then let $X = A \cup \{p\}$. (Thus X is obtained from R^2 by deleting the positive x -axis and the closed unit disk, then restoring the point p . Note that X is locally compact except at p .)

Define $h(\theta) = \theta/(2\pi - \theta)$. The function h maps the interval $(0, 2\pi)$ homeomorphically onto $(0, \infty)$.

Now, for $t \geq 0$, define the map $F_t: X \rightarrow X$ in the following way. Set $F_t(p) = p$; and if $r > 1$, put $F_t(r, \theta) = (r, h^{-1}[h(\theta) + t/(r-1)])$. It is straightforward to check that $\{F_t: t \geq 0\}$ is a semigroup. (Roughly speaking, the flow F_t makes the circular arc with radius $r > 1$ collapse in a counterclockwise sense with increasing velocity as $r \rightarrow 1$.) To see that F_t is continuous for fixed $t > 0$, it is only necessary to worry about what happens at the point p . So let $x_n = (r_n, \theta_n)$ with $r_n \rightarrow 1$, $\theta_n \rightarrow 0$ or 2π . Then $h(\theta_n) + t/(r_n - 1) \rightarrow \infty$, and so $h^{-1}[h(\theta_n) + t/(r_n - 1)] \rightarrow 2\pi$. Hence, $F_t(r_n, \theta_n) \rightarrow (1, 2\pi) = p = F_t(p)$. If $t = 0$, F_t is identity. Also, it is clear that $F_t(x)$ is continuous in t for fixed x in X .

Finally, we verify the failure of joint continuity at p with $t = 0$. Take $x_n = (r_n, \theta_n)$ with $r_n = 1 + 1/n$ and $\theta_n = 1/n$. Take $t_n = 1/n$. Then $x_n \rightarrow p$ and $t_n \rightarrow 0$, but

$$(2) \quad F_{t_n}(x_n) = (r_n, h^{-1}[h(1/n) + 1]) \rightarrow (1, h^{-1}(1)) = (1, \pi) \neq p.$$

We can now obtain an example on an open subset of Hilbert space quite cheaply (modulo infinite-dimensional topology!). Indeed, let H be a separable, infinite-dimensional Hilbert space. The space X considered above is easily seen to be a locally finite-dimensional simplicial complex; that is, the two-dimensional space X can be triangulated and homeomorphically embedded as a piecewise linear subset of H so that the vertices of the triangulation

correspond to mutually orthogonal unit vectors. Hence, by [5, Theorem 3], the product space $X \times H$ is a manifold modelled on H . By the results in [3], $X \times H$ is homeomorphic to an open subset of H . We then simply take as our semigroup on $X \times H$ the maps $G_t = F_t \times I$.

The referee has observed that a simple modification of this construction yields a semigroup acting on the whole space H . Consider the metric cone C of X : if X is embedded as a piecewise linear subset of H , then C is the subset of $H \times \mathbf{R}$ consisting of the points $(\lambda x, 1 - \lambda)$ with $x \in X$ and $0 \leq \lambda \leq 1$. The semigroup F_t extends in the obvious way to C : we define $F'_t(\lambda x, 1 - \lambda) = (\lambda F_t(x), 1 - \lambda)$. Then $F'_t \times I$ is the desired semigroup on $C \times H$. The point is that C is a *contractible* locally finite-dimensional simplicial complex, and so, by [3, Corollary 3], $C \times H$ is homeomorphic to H .

Acknowledgement. I wish to thank Professor Victor Klee for some interesting correspondence concerning the proof that $X \times H$ is homeomorphic to an open subset of H . In particular, Klee has shown that this can be obtained via the machinery developed in [4].

REFERENCES

1. P. R. Chernoff and J. Marsden, *On continuity and smoothness of group actions*, Bull. Amer. Math. Soc. 76 (1970), 1044–1049. MR 42 #419.
2. J. R. Dorroh, *Semi-groups of maps in a locally compact space*, Canad. J. Math. 19 (1967), 688–696. MR 37 #4777.
3. D. W. Henderson, *Infinite-dimensional manifolds are open subsets of Hilbert space*, Topology 9 (1970), 25–33. MR 40 #3581; erratum, 40, p. 1706.
4. V. L. Klee, Jr., *Convex bodies and periodic homeomorphisms in Hilbert space*, Trans. Amer. Math. Soc. 74 (1953), 10–43. MR 14, 989.
5. J. E. West, *Products of complexes and Fréchet spaces which are manifolds*, Trans. Amer. Math. Soc. 166 (1972), 317–337. MR 45 #2756.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY,
CALIFORNIA 94720