

EXTENSIONS OF MONOTONE OPERATOR FUNCTIONS

J. D. CHANDLER, JR.¹

ABSTRACT. It is shown that a monotone operator function f defined on an open subset Δ of the real numbers may be extended to a monotone operator function on the convex hull of Δ .

Let f be a bounded real-valued Borel-measurable function defined on a Borel subset Δ of the real numbers. Let A be a bounded selfadjoint operator on a separable complex Hilbert space H such that the spectrum $\sigma(A)$ of A is contained in Δ . Then $f(A)$ is the selfadjoint operator on H defined by

$$\langle f(A)\phi, \psi \rangle = \int_{\sigma(A)} f(\lambda) \langle E(d\lambda)\phi, \psi \rangle, \quad \phi, \psi \in H,$$

where E is the resolution of the identity corresponding to A .

The function f is said to be a *monotone operator function* on Δ if $f(A) \leq f(B)$ whenever A and B are bounded selfadjoint operators on a Hilbert space H such that $\sigma(A) \subseteq \Delta$, $\sigma(B) \subseteq \Delta$, and $A \leq B$. If f satisfies this monotonicity condition for the totality of finite-dimensional complex Hilbert spaces, then f is said to be a *monotone matrix function* on Δ .

The following characterization of monotone operator functions is essentially due to Loewner [5]. Loewner proved the theorem for monotone matrix functions; Bendat and Sherman [1] proved that a function f is monotone matrix on an open interval (a, b) if and only if f is monotone operator on (a, b) .

THEOREM 1. *A real-valued function f defined on an open interval (a, b) is a monotone operator function on (a, b) if and only if f is analytic on (a, b) , can be analytically continued onto the upper half-plane, and represents there a holomorphic function with nonnegative imaginary part.*

It will be shown that there is a similar characterization of monotone operator functions defined on an arbitrary open subset Δ of the real numbers.

Let f be a monotone operator function on an open subset Δ of the real numbers. By Theorem 1, f is continuously differentiable on Δ ; see also [4, p. 73]. Associated with f is the kernel K , defined on $\Delta \times \Delta$ by

$$(1) \quad K(\lambda, \mu) = \frac{f(\lambda) - f(\mu)}{\lambda - \mu} \quad (\lambda \neq \mu), \quad K(\lambda, \lambda) = f'(\lambda).$$

Loewner made extensive use of this kernel in his study of monotone matrix

Received by the editors May 21, 1975.

AMS (MOS) subject classifications (1970). Primary 47A60, 47B15; Secondary 30A14, 30A76.

Key words and phrases. Monotone operator functions, functions with positive imaginary part.

¹ Research supported by NSF Grant GP-31483XZ.

functions. The following theorem is essentially due to Loewner [5]; see also [1] and [4].

THEOREM 2. *Let f be a monotone operator function on an open subset Δ of the real numbers. Then K is a positive matrix; that is,*

$$\sum_{j=1}^n \sum_{k=1}^n K(\lambda_j, \lambda_k) c_j \bar{c}_k \geq 0$$

whenever $\lambda_1, \dots, \lambda_n$ belong to Δ and c_1, \dots, c_n are complex numbers, for $n = 1, 2, 3, \dots$

PROOF. Let A and B be self adjoint operators on an n -dimensional Hilbert space H , and let A and B have spectral representations

$$A = \sum_{j=1}^n \lambda_j E_j, \quad B = \sum_{k=1}^n \mu_k F_k,$$

where $\lambda_1, \dots, \lambda_n$ and μ_1, \dots, μ_n belong to Δ . An easy computation shows that

$$(2) \quad f(B) - f(A) = \sum_{j=1}^n \sum_{k=1}^n K(\lambda_j, \mu_k) E_j (B - A) F_k.$$

Let $B = A + \epsilon P$ be a one-dimensional perturbation of A , say $P = \langle \cdot, \phi \rangle \phi$ for some ϕ in H , such that $\epsilon > 0$ and $\sigma(A + \epsilon P) \subseteq \Delta$. Then B has a spectral representation

$$B = \sum_{k=1}^n \lambda_k(\epsilon) E_k(\epsilon),$$

where $\lambda_k(\epsilon) \rightarrow \lambda_k$ as $\epsilon \downarrow 0$ ($k = 1, \dots, n$). Equation (2) shows that, for any ψ in H ,

$$\begin{aligned} 0 &\leq \langle (1/\epsilon)[f(B) - f(A)]\psi, \psi \rangle \\ &= \sum_{j=1}^n \sum_{k=1}^n K(\lambda_j(\epsilon), \lambda_k) \langle E_j P E_k(\epsilon) \psi, \psi \rangle; \end{aligned}$$

take $\epsilon \downarrow 0$ to show that

$$\sum_{j=1}^n \sum_{k=1}^n K(\lambda_j, \lambda_k) \langle E_k \psi, \phi \rangle \langle \phi, E_j \psi \rangle \geq 0.$$

Now choose $\phi = \sum_{j=1}^n c_j e_j$ and $\psi = \sum_{k=1}^n e_k$, where $E_j e_j = e_j$ and $\|e_j\| = 1$ ($j = 1, \dots, n$), to show that

$$\sum_{j=1}^n \sum_{k=1}^n K(\lambda_j, \lambda_k) c_j \bar{c}_k \geq 0.$$

Loewner [5] proved that the converse of Theorem 2 is true if Δ is an open interval (a, b) . This result, together with Theorem 1, shows that a real-valued continuously differentiable function f , defined on an open interval (a, b) , admits an analytic continuation F onto the upper half-plane Π^+ with

$\text{Im}(F) \geq 0$ on Π^+ , if and only if the kernel K associated with f is a positive matrix.

This aspect of Loewner's work was generalized by Rosenblum and Rovnyak [6, Theorem 8]; the following theorem is an immediate consequence of their generalization.

THEOREM 3. *Let f be a real-valued continuously differentiable function on an open subset Δ of the real numbers, and let K be the kernel associated with f , as in (1). If K is a positive matrix, then there exists a function F such that*

- (i) F is separately holomorphic on the upper and lower half-planes,
- (ii) F has nonnegative imaginary part on the upper half-plane, and
- (iii) F may be analytically continued by reflection across Δ so that

$$f(x) = F(x + i0) = F(x - i0), \quad x \in \Delta.$$

The next theorem is due to Šmul'jan [7]; it depends heavily on the work of Davis [2], [3] on convex operator functions.

THEOREM 4. *Let f be a monotone operator function on $\{a\} \cup (b, c)$, where $-\infty < a < b < c < \infty$. Then there exists a unique monotone operator function g defined on (a, c) such that $g = f$ on (b, c) and $g(a + 0) \geq f(a)$.*

Šmul'jan proved this theorem for the special case $a = 0$, $b = 1$, and $c = 2$. As noted in [7], the general case follows by considering $f \circ h$, where h is the unique linear fractional transformation such that $h(0) = a$, $h(1) = b$, and $h(2) = c$.

Theorems 3 and 4 are now combined to yield a characterization of monotone operator functions defined on arbitrary open subsets of the real numbers.

THEOREM 5. *Let f be a monotone operator function defined on an open subset Δ of the real numbers. Then there exists a monotone operator function h defined on the convex hull D of Δ such that $f = h$ on Δ .*

PROOF. If f is a monotone operator function on Δ , then f is continuously differentiable on Δ . By Theorem 2, the kernel K associated with f is a positive matrix. By Theorem 3, f admits an analytic continuation F onto the upper half-plane that satisfies (i)—(iii) of that theorem.

Let a and c be arbitrary points in Δ with $a < c$. Since Δ is open, there is a point b in Δ such that $a < b < c$ and $(b, c) \subseteq \Delta$. By Theorem 4, there exists a monotone operator function g on (a, c) such that $f = g$ on (b, c) . By Theorem 3, g admits an analytic continuation G onto the upper half-plane that satisfies (i)—(iii) of that theorem, with Δ replaced by (a, c) .

Let F and G be analytically continued by reflection across Δ and (a, c) , respectively, as in Theorem 3(iii). But $F = f$ on Δ , $G = g$ on (a, c) , and $f = g$ on (b, c) ; since the holomorphic functions F and G coincide on (b, c) , they coincide on the intersection of their domains. Thus $F = G$ on Π^+ , and so F may be analytically continued across $\Delta \cup (a, c)$ by reflection. Since a and c are arbitrary points in Δ , it follows that F may be analytically continued across the convex hull D of Δ by reflection.

Let H be this extension of F to the complex plane slit along $(-\infty, \infty) - D$, and let h be the restriction of H to D . Since H has nonnegative imaginary part in the upper half-plane and since h is real-valued and analytic on D , h is a monotone operator function on D , by Theorem 1. Since $h = f$ on Δ , h is the desired extension of f to D .

The converse of Theorem 5 follows directly from the definition of "monotone operator function": if g is a monotone operator function on an interval D and if Δ is an open subset of D , then $g|_{\Delta}$ is a monotone operator function on Δ .

Note that Theorem 3 was essential to the proof of Theorem 5; Loewner's original theorem (Theorem 1 above) does not provide the necessary single analytic continuation of f onto the upper half-plane.

The author is grateful to Professors Marvin Rosenblum and James Rovnyak for suggesting this problem.

REFERENCES

1. J. Benda and S. Sherman, *Monotone and convex operator functions*, Trans. Amer. Math. Soc. **79** (1955), 58—71. MR **18**, 588.
2. C. Davis, *A Schwarz inequality for convex operator functions*, Proc. Amer. Math. Soc. **8** (1957), 42—44. MR **18**, 812.
3. ———, *Notions generalizing convexity for functions defined on spaces of matrices*, Proc. Sympos. Pure Math., vol. 7, Amer. Math. Soc., Providence, R.I., 1963, pp. 187—201. MR **27** #5771.
4. W. Donoghue, *Monotone matrix functions and analytic continuation*, Springer-Verlag, New York, Heidelberg, Berlin, 1974.
5. K. Löwner, *Über monotone Matrixfunktionen*, Math. Z. **38** (1934), 177—216.
6. M. Rosenblum and J. Rovnyak, *Restrictions of analytic functions. I*, Proc. Amer. Math. Soc. **48** (1975), 113—119.
7. Ju. L. Šmul'jan, *Monotone operator functions on a set consisting of an interval and a point*, Ukrain. Mat. Ž. **17** (1965), no. 1, 130—136; English transl., Amer. Math. Soc. Transl. (2) **67** (1968), 25—32. MR **32** #6239.

GRADUATE SCHOOL OF ARTS AND SCIENCES, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VIRGINIA 22901