

ON A FIXED POINT THEOREM OF CONTRACTIVE TYPE¹

CHI SONG WONG

ABSTRACT. A refinement of the proof of a recent fixed point theorem of Caristi is obtained.

The main purpose of this paper is to refine the proof of the following recent result of James Caristi [3].

THEOREM. *Let f be a self-map on a nonempty complete metric space (X, d) . Suppose that there exists a lower semicontinuous function V of X into $[0, \infty)$ such that*

$$d(x, f(x)) \leq V(x) - V(f(x)), \quad x \in X.$$

Then f has a fixed point.

PROOF. Let $x_0 \in X$ and ν be a nonzero ordinal. Suppose that there exists a net $\{x_\beta\}_{\beta < \nu}$ in X such that:

- (i) if β is an isolated nonzero ordinal less than ν , then $x_\beta = f(x_{\beta-1})$;
- (ii) if β is a limiting ordinal less than ν , then the net $\{x_\alpha\}_{\alpha < \beta}$ converges to x_β ;
- (iii) for all $0 \leq \alpha < \beta < \nu$, $d(x_\alpha, x_\beta) \leq V(x_\alpha) - V(x_\beta)$. If ν is an isolated ordinal, define $x_\nu = f(x_{\nu-1})$ and note that for $\alpha < \nu$,

$$\begin{aligned} d(x_\alpha, x_\nu) &\leq d(x_\alpha, x_{\nu-1}) + d(x_{\nu-1}, x_\nu) \\ &\leq (V(x_\alpha) - V(x_{\nu-1})) + (V(x_{\nu-1}) - V(x_\nu)) = V(x_\alpha) - V(x_\nu). \end{aligned}$$

Suppose now that ν is a limiting ordinal. We claim that $\{x_\beta\}_{\beta < \nu}$ is a Cauchy net. Suppose not. Then there exists $\epsilon > 0$ and a strictly increasing sequence $\{\alpha_n\}$ in $(0, \nu)$ such that $d(x_{\alpha_{2n}}, x_{\alpha_{2n-1}}) \geq \epsilon$ for all n . Thus

$$\begin{aligned} \infty &= \sum_{n=1}^{\infty} d(x_{\alpha_{2n}}, x_{\alpha_{2n-1}}) \\ &\leq \sum_{n=1}^{\infty} (V(x_{\alpha_{2n}}) - V(x_{\alpha_{2n-1}})) \leq V(x_0), \end{aligned}$$

a contradiction. So $\{x_\beta\}_{\beta < \nu}$ is Cauchy and therefore, by completeness of

Received by the editors July 23, 1975.

AMS (MOS) subject classifications (1970). Primary 47H10; Secondary 54H25.

Key words and phrases. Completeness, lower semicontinuity, transfinite induction.

¹ This research was partially supported by National Research Council of Canada, Grant A8518 and a grant from Canadian Mathematical Congress. It was prepared while the author was at the Summer Research Institute at Dalhousie University.

© American Mathematical Society 1976

(X, d) , $\{x_\alpha\}_{\alpha < \nu}$ converges to some point which we shall define to be x_ν . Let $\alpha < \nu$. Since V is lower semicontinuous,

$$\begin{aligned} d(x_\alpha, x_\nu) &= \lim_{\beta < \nu} d(x_\alpha, x_\beta) \leq \lim_{\beta < \nu} \inf_{\beta < \nu} (V(x_\alpha) - V(x_\beta)) \\ &= V(x_\alpha) - \limsup_{\beta < \nu} V(x_\beta) \leq V(x_\alpha) - V(x_\nu). \end{aligned}$$

This shows that for any ordinal ν , there exists a net $\{x_\beta\}_{\beta < \nu}$ in X with properties (i), (ii), and (iii). If f has no fixed point, then $\{V(x_\alpha)\}_{\alpha < \nu}$ is strictly decreasing and so a contradiction is obtained if we take the cardinality of ν larger than that of X (or $\{1, 2, \dots\}$). Q.E.D.

The next result follows from the above theorem.

PROPOSITION. *Let (X, d) be a nonempty complete metric space. Let f be a self-map on X . Suppose that there exists a lower semicontinuous function of X into $[0, \infty)$ such that for any x in X with $x \neq f(x)$, there exists y in $X \setminus \{x\}$ such that $d(x, y) \leq V(x) - V(y)$. Then f has a fixed point.*

We are informed by the referee and W. A. Kirk that a theorem which is equivalent to Caristi's theorem was announced by I. Ekeland in 1972 [5], a proof [8] of Ekeland's theorem using Zorn's lemma can be found in a recent paper of A. Brønsted [2], and a proof of Caristi's theorem without using Zorn's lemma was obtained by F. E. Browder [1]. Applications of the above theorem of Caristi can be found in [3], [4], [6], [7], [8].

REFERENCES

1. F. E. Browder, *On a theorem of Caristi and Kirk*, Proc. Seminar on Fixed Point Theory and its Applications, Dalhousie University, June 1975 (to appear).
2. A. Brønsted, *On a lemma of Bishop and Phelps*, Pacific J. Math. **55** (1974), 335-341.
3. J. Caristi, *Fixed point theorems for mappings satisfying inwardness conditions*, Trans. Amer. Math. Soc. **215** (1976), 241-251.
4. J. Caristi and W. A. Kirk, *Geometric fixed point theory and inwardness conditions*, Proc. Conf. on Geometry of Metric and Linear Spaces, Michigan State University, 1974 (to appear).
5. I. Ekeland, *Sur les problèmes variationnels*, C. R. Acad. Sci. Paris Sér. A-B **275** (1972), A1057-A1059. MR **46** #9768.
6. W. A. Kirk, *Caristi's fixed point theorem and the theory of normal solvability*, Proc. Seminar on Fixed Point Theory and its Applications, Dalhousie University, 1975 (to appear).
7. W. A. Kirk and J. Caristi, *Mapping theorems in metric and Banach spaces*, Bull. Acad. Polon. Sci. **23** (1975).
8. W. A. Kirk, *Caristi's fixed point theorem and metric convexity* (to appear).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WINDSOR, WINDSOR, ONTARIO, CANADA N9B 3P4