

EIGENVALUES OF HOPF MANIFOLDS

ERIC BEDFORD¹ AND TATSUO SUWA²

ABSTRACT. The eigenvalues of the Laplacians Δ and \square on the Hopf manifolds are described. Some isospectral results are also given.

On a complex manifold M there are the de Rham and Dolbeault complexes, with operators d and $\bar{\partial}$, respectively. If we fix a hermitian metric on the manifold M , then d and $\bar{\partial}$ will have formal adjoints δ and $\bar{\delta}$ with respect to the hermitian volume element. With these, we define the real Laplacian $\Delta = d\delta + \delta d$ and the complex Laplacian $\square = \bar{\partial}\bar{\delta} + \bar{\delta}\bar{\partial}$. We have $\Delta = 2\square = 2\bar{\square}$ if M is Kähler.

In this note, we study the eigenvalues of the Laplacians Δ and \square on the Hopf manifolds M_α , $0 < |\alpha| < 1$, which are not Kähler. The notation $\text{Sp}(M, A)$ will be used to denote the eigenvalues with multiplicity (spectrum) of the operator A on the manifold M . In Theorem 1, the eigenvalues of Δ and \square are described. The computation of the eigenfunctions shows that the eigenspaces of \square refine those of Δ . Some isospectral results are also given. For α in a certain domain, $\text{Sp}(M_\alpha, \Delta)$ determines M_α up to isometry (Theorem 2). Moreover, it is shown that for "most" values of α , M_α may be determined up to isometry from either $\text{Sp}(M_\alpha, \Delta)$ or $\text{Sp}(M_\alpha, \square)$ (Theorem 3).

Let W denote the n -dimensional complex vector space $\mathbf{C}^n = \{z \mid z = (z_1, \dots, z_n)\}$ minus the origin; $W = \mathbf{C}^n - \{0\}$, and let α be a complex number with $0 < |\alpha| < 1$. Consider the analytic automorphism g_α of W defined by $g_\alpha(z_1, \dots, z_n) = (\alpha z_1, \dots, \alpha z_n)$. The group G_α generated by g_α is an infinite cyclic group acting on W freely and properly discontinuously. Thus the quotient $M_\alpha = W/G_\alpha$ is an n -dimensional complex manifold, which is called a (homogeneous) Hopf manifold. It is easy to see that M_α is diffeomorphic to $S^1 \times S^{2n-1}$, where S^r denotes the standard r -sphere (cf. the proof of Lemma 3). If $n = 1$, M_α is the complex torus whose lattice is generated by 1 and $(2\pi i)^{-1} \log \alpha$, and if $n > 1$, M_α is a non-Kähler manifold. The hermitian metric

$$(1) \quad g = \|z\|^{-2} \sum_{i=1}^n dz_i d\bar{z}_i, \quad \|z\|^2 = \sum_{i=1}^n z_i \bar{z}_i,$$

Received by the editors June 9, 1975.

AMS (MOS) subject classifications (1970). Primary 53C55, 32C10; Secondary 35P99, 58G99.

Key words and phrases. Hopf manifold, hermitian metric, Laplacian, eigenvalues, isospectral problem.

¹ Research supported in part by a Sloan Foundation Grant to CIMS and Army Research Office No. DAHC 04-74-G-0159.

² Research supported in part by NSF Grant GP 38878.

Copyright © 1977, American Mathematical Society

on W is G_α -invariant. Hence, it induces a hermitian metric on M_α . From now on we think of M_α as a hermitian manifold with this metric. It is easy to check that if $n = 1$, M_α is a flat torus. If α is real, M_α is isometric to $S^1 \times S^{2n-1}$ as a riemannian manifold. We denote by $\mathcal{C}^\infty(M_\alpha)$ the space of complex valued smooth functions on M_α . Consider the real and complex Laplacians Δ and \square induced by the hermitian metric on M_α . Since the operator δ maps a (p, q) -form to a $(p, q - 1)$ -form, we have

$$(2) \quad \Delta = \square + \bar{\square} \quad \text{on } \mathcal{C}^\infty(M_\alpha).$$

A straightforward computation [3, p. 97] shows that the complex Laplacian on $\mathcal{C}^\infty(M_\alpha)$ is

$$\square = -\|z\|^2 \sum_{i=1}^n \frac{\partial^2}{\partial z_i \partial \bar{z}_i} + (n - 1) \sum_{i=1}^n \bar{z}_i \frac{\partial}{\partial \bar{z}_i}.$$

Let

$$\Delta_0 = - \sum_{i=1}^n \frac{\partial^2}{\partial z_i \partial \bar{z}_i}$$

be the standard Laplacian on \mathbf{C}^n . Moreover, let $\mathcal{H}_{p,q}$ be the space of harmonic polynomials of type (p, q) , i.e., the polynomials f on \mathbf{C}^n such that $\Delta_0 f = 0$ and $f(z) = \sum_{|\mu|=p; |\nu|=q} c_{\mu\nu} z^\mu \bar{z}^\nu$, where $\mu = (\mu_1, \dots, \mu_n)$ and $\nu = (\nu_1, \dots, \nu_n)$ are n -tuples of nonnegative integers, $|\mu| = \sum_{i=1}^n \mu_i$, $|\nu| = \sum_{i=1}^n \nu_i$, and $z^\mu = z_1^{\mu_1} \cdots z_n^{\mu_n}$, $\bar{z}^\nu = \bar{z}_1^{\nu_1} \cdots \bar{z}_n^{\nu_n}$.

LEMMA 1. For $f \in \mathcal{H}_{p,q}$ and for a complex number γ ,

$$(3) \quad \square (\|z\|^\gamma f) = -(\gamma/2)^2 - (p + q)\gamma/2 + q(n - 1) \|z\|^\gamma f.$$

PROOF. Since $\Delta_0 f = 0$, we have

$$\begin{aligned} \square (\|z\|^\gamma f) &= -\left(\frac{\gamma}{2}\right)^2 \|z\|^\gamma f - \frac{\gamma}{2} \|z\|^\gamma \sum_{i=1}^n z_i \frac{\partial f}{\partial z_i} \\ &\quad + \left(n - 1 - \frac{\gamma}{2}\right) \|z\|^\gamma \sum_{i=1}^n \bar{z}_i \frac{\partial f}{\partial \bar{z}_i}. \end{aligned}$$

Substituting

$$\sum_{i=1}^n z_i \frac{\partial f}{\partial z_i} = pf \quad \text{and} \quad \sum_{i=1}^n \bar{z}_i \frac{\partial f}{\partial \bar{z}_i} = qf$$

in the above equation, we get (3). Q.E.D.

Let ω be a complex number such that $e^{2\pi i \omega} = \alpha$. Since $|\alpha| < 1$, we have $\text{Im } \omega > 0$.

DEFINITION. For nonnegative integers p and q , let $\Gamma_{p,q}$ be the set of complex numbers γ which can be expressed as $\gamma = -\gamma_1 + \bar{\gamma}_2$ with $\text{Re } \gamma_1 = p$, $\text{Re } \gamma_2 = -q$, and $\text{Re}((\gamma_1 + \gamma_2)\bar{\omega}) \in \mathbf{Z}$.

REMARKS. 1°. $\gamma_1 + \gamma_2$ is in the lattice dual to the one generated by 1 and ω (cf. [1, p. 146]).

2°. Let $a = \text{Re } \omega$ and $b = \text{Im } \omega$. Then

$$(4) \quad \Gamma_{p,q} = \{-(p + q) + ((a(p - q) - k)/b)i \mid k \in \mathbf{Z}\}.$$

LEMMA 2. For $f \in \mathcal{K}_{p,q}$, the function $\|z\|^\gamma f$ is G_α -invariant if and only if $\gamma \in \Gamma_{p,q}$.

PROOF. $\|z\|^\gamma f$ is G_α -invariant if and only if $|\alpha|^\gamma \alpha^p \bar{\alpha}^q = 1$, or equivalently,

$$\begin{cases} \text{Re } \gamma = -(p + q) \text{ and} \\ -\text{Im } \omega \cdot \text{Im } \gamma + (p - q) \text{Re } \omega = k, \text{ for some } k \in \mathbf{Z}. \end{cases}$$

In view of (4), this is equivalent to $\gamma \in \Gamma_{p,q}$. Q.E.D.

LEMMA 3. The vector subspace of $\mathcal{C}^\infty(M_\alpha)$ generated by $\cup_{p,q \geq 0} \{\|z\|^\gamma f \mid f \in \mathcal{K}_{p,q}, \gamma \in \Gamma_{p,q}\}$ is dense in $\mathcal{C}^\infty(M_\alpha)$.

PROOF. Consider the unit $(2n - 1)$ -sphere $S^{2n-1} = \{z \in \mathbf{C}^n \mid \|z\| = 1\}$. The map $\varphi: \mathbf{R} \times S^{2n-1} \rightarrow W$ defined by $\varphi(t, z_1, \dots, z_n) = (e^{2\pi i \omega t} z_1, \dots, e^{2\pi i \omega t} z_n)$ is clearly a diffeomorphism. Moreover, φ induces a diffeomorphism, which we denote also by φ , from $S^1 \times S^{2n-1}$ onto M_α , where $S^1 = \mathbf{R}/\mathbf{Z}$. Take a function $\|z\|^\gamma f$ with $f \in \mathcal{K}_{p,q}$ and $\gamma \in \Gamma_{p,q}$. The pullback of $\|z\|^\gamma f$ by φ is given by

$$((\|z\|^\gamma f) \circ \varphi)(t, z) = \exp(2\pi i t(i\gamma \text{Im } \omega + p\omega - q\bar{\omega})) f(z) = e^{2\pi i k t} f(z),$$

where $k = \text{Re}((\gamma_1 + \gamma_2)\bar{\omega}) \in \mathbf{Z}$. On the other hand, the vector subspace of $\mathcal{C}^\infty(S^1)$ generated by the functions $\{e^{2\pi i k t}\}_{k \in \mathbf{Z}}$ is dense in $\mathcal{C}^\infty(S^1)$ and the harmonic polynomials are dense in $\mathcal{C}^\infty(S^{2n-1})$ [1, p. 160]. Q.E.D.

THEOREM 1. (i) The eigenvalues of \square on $\mathcal{C}^\infty(M_\alpha)$ are $|\gamma|^2/4 + q(n - 1)$, $p, q \in \mathbf{Z}^+$, $\gamma \in \Gamma_{p,q}$.

(ii) The eigenvalues of Δ on $\mathcal{C}^\infty(M_\alpha)$ are $|\gamma|^2/2 + (p + q)(n - 1)$, $p, q \in \mathbf{Z}^+$, $\gamma \in \Gamma_{p,q}$, where \mathbf{Z}^+ denotes the nonnegative integers.

PROOF. If $\gamma \in \Gamma_{p,q}$, we have $-(\gamma/2)^2 - (p + q)\gamma/2 + q(n - 1) = |\gamma|^2/4 + q(n - 1)$. The theorem follows from (2), Lemmas 1-3, and Lemma A.II.1 in [1, p. 143]. Q.E.D.

Now let us discuss the isospectral problem. Let $a + bi = \omega = (2\pi i)^{-1} \log \alpha$ as before. Since $a = (2\pi)^{-1} \arg \alpha$, we may assume that $|a| \leq \frac{1}{2}$. Note that if $|a| = \frac{1}{2}$, then α is real and M_α is isometric to $S^1 \times S^{2n-1}$.

THEOREM 2. Suppose (i) $b > 1/2\sqrt{2n + 1}$ or (ii) $|a| < b\sqrt{2n + 1}$. Then $\text{Sp}(M_\alpha, \Delta)$ determines M_α up to isometry, namely, $\text{Sp}(M_\alpha, \Delta) = \text{Sp}(M_{\alpha'}, \Delta)$ implies M_α is isometric to $M_{\alpha'}$.

PROOF. Let

$$a' + b'i = \omega' = (2\pi i)^{-1} \log \alpha', \quad |\alpha'| \leq \frac{1}{2}.$$

If $\text{Sp}(M_\alpha, \Delta) = \text{Sp}(M_{\alpha'}, \Delta)$, then the volumes of M_α and $M_{\alpha'}$ are the same [1, Corollaire E.IV.2, p. 216], see also [4]. This implies $|\alpha| = |\alpha'|$ or equivalently $b = b'$. Consider the eigenvalues

$$\lambda(p, q, k, a^{(\nu)}) = (p + q)^2/2 + (a^{(\nu)}(p - q) - k)^2/2b^2 + (p + q)(n - 1),$$

$p, q \in \mathbf{Z}^+, k \in \mathbf{Z}$, of Δ on $\mathcal{C}^\infty(M_{\alpha^{(\nu)}})$, $\nu = 0, 1$, $a^{(0)} = a, a^{(1)} = a'$. Note that if $p + q \geq 2$, then $\lambda(p, q, k, a^{(\nu)}) \geq 2n$ and that $\lambda(p, q, k, a^{(\nu)}) = 0$ if and only if $p = q = k = 0$.

Case (i). $b > 1/2\sqrt{2n + 1}$. Since $|a| \leq \frac{1}{2}$, we have

$$\begin{aligned} \lambda(1, 0, 0, a) &= a^2/2b^2 + n - \frac{1}{2} < 2a^2(2n + 1) + n - \frac{1}{2} \leq \frac{1}{2}(2n + 1) + n - \frac{1}{2} \\ &= 2n. \end{aligned}$$

Similarly, we have $\lambda(1, 0, 0, a') < 2n$. Hence, we have

$$\lambda(1, 0, 0, a) = \lambda(p_1, q_1, k_1, a') \quad \text{and} \quad \lambda(1, 0, 0, a') = \lambda(p_2, q_2, k_2, a)$$

for some $p_j, q_j \in \mathbf{Z}^+$ and $k_j \in \mathbf{Z}$ with $0 \leq p_j + q_j \leq 1, j = 1, 2$. If $p_1 + q_1 = 1$, we have $a^2 = (a' \pm k_1)^2$. Since $|a^{(\nu)}| \leq \frac{1}{2}, \nu = 0, 1$, we get $a' = \pm a$. If $p_2 + q_2 = 1$, we also get $a' = \pm a$. If $p_j + q_j = 0, j = 1, 2$, we have $a^2 - a'^2 = k_1^2 - k_2^2$. Since $|a^2 - a'^2| \leq \frac{1}{2}$, we get $a' = \pm a$.

Case (ii). $0 < b \leq 1/2\sqrt{2n + 1}, |a| < b\sqrt{2n + 1}$. We have

$$\lambda(1, 0, 0, a) = \frac{a^2}{2b^2} + n - \frac{1}{2} < n + \frac{1}{2} + n - \frac{1}{2} = 2n.$$

On the other hand

$$\lambda(0, 0, k, a') = \frac{k^2}{2b^2} \geq 2k^2(2n + 1) > 2n, \quad \text{for } |k| \geq 1.$$

Hence, we have $\lambda(1, 0, 0, a) = \lambda(p_1, q_1, k_1, a')$, with $p_1 + q_1 = 1$. Thus, we get $a' = \pm a$.

If $a' = -a$, then $\alpha' = \bar{\alpha}$ and the diffeomorphism $W \rightarrow W$ defined by $(z_1, \dots, z_n) \mapsto (\bar{z}_1, \dots, \bar{z}_n)$ induces an isometry between M_α and $M_{\alpha'}$. Q.E.D.

THEOREM 3. *Suppose a and b are algebraically independent over \mathbf{Q} . Then M_α may be determined up to isometry from either $\text{Sp}(M_\alpha, \Delta)$ or $\text{Sp}(M_\alpha, \square)$.*

PROOF. We show that $\text{Sp}(M_\alpha, \square)$ determines $|a|$ and b . The same argument with only small modifications will apply to $\text{Sp}(M_\alpha, \Delta)$. Let α' be a complex number with $a' + ib' = (2\pi i)^{-1} \log \alpha', -\frac{1}{2} \leq a' \leq \frac{1}{2}$. We shall assume that $\text{Sp}(M_\alpha, \square) = \text{Sp}(M_{\alpha'}, \square)$ and show that $|a| = |a'|$ and $b = b'$. Since the volume of M_α determines b , and the volume of M_α is determined by the asymptotic behavior of the spectrum (see, for instance, Gilkey [4]), it follows that $b = b'$. Let

$$\lambda(p, q, k, a, b) = (p + q)^2/4 + (a(p - q) - k)^2/4b^2 + (n - 1)q.$$

It is easily seen that the linear span of $\{b^2\lambda \mid \lambda \in \text{Sp}(M_\alpha, \square)\}$ over \mathbf{Q} is the linear span over \mathbf{Q} of $\{1, a, a^2, b^2\}$. Thus if $\text{Sp}(M_\alpha, \square) = \text{Sp}(M_{\alpha'}, \square)$, then $\{1, a, a^2, b^2\}$ and $\{1, a', (a')^2, b^2\}$ have the same linear span over \mathbf{Q} . Thus there exist rationals $r_j, r'_j, 1 \leq j \leq 4$, such that

$$a' = r_1 + ar_2 + a^2r_3 + b^2r_4, \quad (a')^2 = r'_1 + ar'_2 + a^2r'_3 + b^2r'_4.$$

By the algebraic independence of a and b , it follows by squaring the left-hand equation that $r_3 = r_4 = 0$. Thus $a' = r_1 + ar_2$.

Now we show that $r_1 = 0, r_2 = \pm 1$. Observe that if

$$\text{Sp}(M_\alpha, \square) = \text{Sp}(M_{\alpha'}, \square)$$

then for integers (p, q, k) there exists (p', q', k') such that

$$(5) \quad \lambda(p, q, k, a, b) = \lambda(p', q', k', r_1 + ar_2, b).$$

Conversely, for (p', q', k') there exists (p, q, k) . Multiplying (5) by b^2 and comparing the coefficients of 1 and a^2 , we obtain

$$(6) \quad k^2 = ((p' - q')r_1 - k')^2,$$

$$(7) \quad (p - q)^2 = ((p' - q')r_2)^2.$$

Since (7) must always have integer solutions, $r_2 = \pm 1$. From equation (6) it follows that r_1 must be an integer, and so $r_1 = 0$ since $-\frac{1}{2} \leq a' \leq \frac{1}{2}$. Thus $|a|$ and b are determined. Q.E.D.

REMARKS. 1. If $n = 1, M_\alpha$ and $M_{\alpha'}$ are biholomorphic, as is well known, if and only if $\begin{pmatrix} \omega \\ 1 \end{pmatrix} = u \begin{pmatrix} \omega' \\ 1 \end{pmatrix}$ for some $u \in \text{SL}(2, \mathbf{Z})$, where

$$\omega = (2\pi i)^{-1} \log \alpha \quad \text{and} \quad \omega' = (2\pi i)^{-1} \log \alpha'.$$

If $n \geq 2, M_\alpha$ and $M_{\alpha'}$ are biholomorphic, by Hartogs' theorem, if and only if $\alpha' = \alpha$ [2, Theorem 15.1].

2. It does not seem easy to find the dimension (multiplicity) of each eigenspace. The dimension of $\mathcal{H}_{p,q}$ can be computed as follows. Let $\mathcal{P}_{p,q}$ denote the space of polynomials on \mathbf{C}^n of type (p, q) and set $r = \|z\|$. Observe that \mathcal{H}_k , the harmonic polynomials homogeneous of degree k , is subdivided into the spaces $\mathcal{H}_{p,q}$, i.e., $\mathcal{H}_k = \bigoplus_{p+q=k} \mathcal{H}_{p,q}$. This may be done since Δ_0 maps $\mathcal{P}_{p,q}$ into $\mathcal{P}_{p-1,q-1}$. By repeating the argument in Berger [1, p. 161], we may conclude that for $p \geq q$,

$$\mathcal{P}_{p,q} = \mathcal{H}_{p,q} \oplus r^2 \mathcal{H}_{p-1,q-1} \oplus \dots \oplus r^{2q} \mathcal{H}_{p-q,0},$$

and the summands are pairwise orthogonal in $L^2(S^{2n-1})$. Thus, $\mathcal{P}_{p,q} = \mathcal{H}_{p,q} \oplus r^2 \mathcal{P}_{p-1,q-1}$ from which it follows that

$$\dim \mathcal{H}_{p,q} = \dim \mathcal{P}_{p,q} - \dim \mathcal{P}_{p-1,q-1}.$$

Since

$$\dim \mathcal{P}_{p,q} = \binom{n-1+p}{p} \binom{n-1+q}{q},$$

we get

$$\dim \mathcal{H}_{p,q} = \binom{n+p-1}{p} \binom{n+q-1}{q} - \binom{n+p-2}{p-1} \binom{n+q-2}{q-1}.$$

REFERENCES

1. M. Berger, P. Gauduchon and E. Mazet, *Le spectre d'une variété riemannienne*, Lecture Notes in Math., vol. 194, Springer-Verlag, Berlin and New York, 1971. MR 43 #8025.
2. K. Kodaira and D. C. Spencer, *On deformations of complex analytic structures. II*, Ann. of Math. (2) 67 (1958), 403–466. MR 22 #3009.
3. J. Morrow and K. Kodaira, *Complex manifolds*, Holt, Rinehart and Winston, New York, 1971. MR 46 #2080.
4. P. Gilkey, *The spectral geometry of real and complex manifolds*, Proc. Sympos. Pure Math., vol. 27, Amer. Math. Soc., Providence, R. I., 1975, pp. 265–280.

COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, NEW YORK, NEW YORK 10012