

NONIMMERSIONS OF FLAG MANIFOLDS

S. A. ILORI

ABSTRACT. Certain useful quadratic identities in the cohomology of classifying spaces induce quadratic equations in the cohomology of a manifold M under the classifying map for the normal bundle of M . In low dimensional flag manifolds, one can show that the quadratic equation has no root, thus establishing a nonimmersion.

1. Introduction. In integral cohomology, one has that

$$(1.1) \quad p_r = \chi^2 \in H^{4r}(\text{BSO}(2r); \mathbb{Z})$$

where $\text{BSO}(2r)$ is the classifying space for oriented real r -plane bundles, p_r is the r th Pontryagin class and χ is the Euler class. An oriented manifold M immerses in codimension k if and only if the stable normal bundle has dimension k . (Cf. [3].) We obtain nonimmersion by showing that the quadratic equation in $H^*(M; \mathbb{Z})$, induced by (1.1), has no integral solution.

If $n = n_1 + \cdots + n_q$, let

$$F(n_1, \dots, n_q) = U(n)/U(\tilde{n}_1) \times \cdots \times U(n_q)$$

denote a complex incomplete flag manifold. We shall show that:

(a) If $M = F(2,4)$, then $\dim M = 16$, $M \subseteq \mathbb{R}^{31}$, $\not\subseteq \mathbb{R}^{24}$.

(b) If $M = F(2,5)$, then $\dim M = 20$, $M \subseteq \mathbb{R}^{39}$, $\not\subseteq \mathbb{R}^{30}$.

The following flag manifolds are bundles with fibre or base equal to $F(2,4)$ or $F(2,5)$ and so results (a) and (b) together with [8] imply the following results.

If $M = F(2,2,2)$, then $\dim M = 24$, $M \subseteq \mathbb{R}^{33}$, $\not\subseteq \mathbb{R}^{32}$.

If $M = F(2,1,3)$, then $\dim M = 22$, $M \subseteq \mathbb{R}^{33}$, $\not\subseteq \mathbb{R}^{30}$.

If $M = F(2,2,3)$, then $\dim M = 32$, $M \subseteq \mathbb{R}^{46}$, $\not\subseteq \mathbb{R}^{42}$.

If $M = F(2,3,2)$, then $\dim M = 32$, $M \subseteq \mathbb{R}^{46}$, $\not\subseteq \mathbb{R}^{42}$.

If $M = F(2,1,4)$, then $\dim M = 28$, $M \subseteq \mathbb{R}^{46}$, $\not\subseteq \mathbb{R}^{38}$.

If $M = F(1,2,4)$, then $\dim M = 28$, $M \subseteq \mathbb{R}^{46}$, $\not\subseteq \mathbb{R}^{36}$.

If $M = F(1,2,5)$, then $\dim M = 34$, $M \subseteq \mathbb{R}^{61}$, $\not\subseteq \mathbb{R}^{44}$.

An interesting aspect of the above results is that the questions of immersion and nonimmersion are now completely solved for the flag manifold $F(2,2,2)$.

The method of this paper was used by Connell [2] to obtain nonimmersions of $F(2,2)$, $F(2,3)$, $F(1,2,2)$ and $F(2,2,2)$. This paper gives an improvement on the result of Connell [2] in the case of the flag manifold $F(2,2,2)$.

It seems that the following conjecture is reasonable.

Received by the editors April 4, 1975 and, in revised form, April 30, 1976.

AMS (MOS) subject classifications (1970). Primary 57A35, 57F15; Secondary 55B15, 55F40.

Key words and phrases. Normal bundle, Chern class, Pontryagin class, Euler class, classifying space.

Copyright © 1977, American Mathematical Society

CONJECTURE. If $M = F(2, n - 1)$, then $\dim M = 4n - 4$, $M \subseteq R^{8n-9}$, $\not\subseteq R^{6n-6}$.

2. $F(2,4)$. Let $GL(n) \rightarrow F(q, n - q)$ be a principal $GL(q, n - q)$ -bundle, λ , where $GL(n)$ is the general linear group with entries in the field of complex numbers and $GL(q, n - q)$ is a subgroup of $GL(n)$ consisting of all matrices of the form

$$A = \left[\begin{array}{c|c} A' & B \\ \hline 0 & A'' \end{array} \right],$$

where $A' \in GL(q)$, $A'' \in GL(n - q)$ and B is any $q \times (n - q)$ -matrix. Let λ', λ'' be the subbundle and quotient-bundle associated with λ . Then λ' is a canonical q -plane bundle over $F(q, n - q)$. By the splitting principle, λ' splits into line bundles ξ_0, \dots, ξ_{q-1} such that the total Chern class of λ' is given by

$$c(\lambda') = \prod_{i=0}^{q-1} c(\xi_i) = \prod_{i=0}^{q-1} (1 + \gamma_i).$$

Thus

$$c_h(\lambda') = \sigma_h(\gamma_0, \dots, \gamma_{q-1}), \quad h = 1, \dots, q,$$

where $\sigma_h = \sigma_h(\gamma_0, \dots, \gamma_{q-1})$ denotes the h th elementary symmetric function in $\gamma_0, \dots, \gamma_{q-1}$. We also denote by $\bar{\sigma}_h = \bar{\sigma}_h(\gamma_0, \dots, \gamma_{q-1})$ the h th complete symmetric function in $\gamma_0, \dots, \gamma_{q-1}$.

The Whitney sum $\lambda' \oplus \lambda''$ of the subbundle and quotient-bundle associated with λ is the trivial n -plane bundle over $F(q, n - q)$. Therefore, $c(\lambda')c(\lambda'') = 1$. Thus

$$c(\lambda'') = \sum_{i=0}^{\infty} (-1)^i \bar{\sigma}_i(\gamma_0, \dots, \gamma_{q-1}),$$

and since λ'' is an $(n - q)$ -plane bundle, then

$$\bar{\sigma}_i = \bar{\sigma}_i(\gamma_0, \dots, \gamma_{q-1}) = 0, \quad i > n - q.$$

We also, therefore, have that

$$\sum_{i=0}^q \sigma_i \sum_{j=0}^{\infty} (-1)^j \bar{\sigma}_j = 1$$

from which follow the relations:

$$(2.1) \quad \sum_{i=0}^k (-1)^i \sigma_i \bar{\sigma}_{k-i} = 0, \quad k > 0.$$

Now, from [1, p. 522], we have that the σ_r , ($1 \leq r \leq q$) generate $H^*(F(q, n - q), Z)$ subject to the relations

$$(2.2) \quad \bar{\sigma}_{n-q+i} = 0, \quad i \geq 0.$$

Also, from [9, §1.3] and [5, pp. 354–359], the Chern classes, c_r , of $F(2,4)$ are given by

$$\begin{aligned}
c_1 &= 6\sigma_1, \\
c_2 &= 16\bar{\sigma}_2 + 18\sigma_2, \\
c_3 &= 26\bar{\sigma}_3 + 58\sigma_2\sigma_1, \\
c_4 &= 31\bar{\sigma}_4 + 91\bar{\sigma}_2\sigma_2 + 67\sigma_2^2, \\
c_5 &= 90\sigma_2\bar{\sigma}_3 + 120\sigma_2^2\sigma_1, \\
c_6 &= 105\sigma_2^2\bar{\sigma}_2 + 65\sigma_2^3, \\
c_7 &= 60\sigma_2^3\sigma_1, \\
c_8 &= 15\sigma_2^4.
\end{aligned}$$

If $F(2,4)$ immerses in codimension 8, its stable normal bundle, ν , is an $SO(8)$ bundle. If d_i denotes the i th Chern class of ν , then using the identities

$$\left(\sum_{i=0}^8 c_i \right) \left(\sum_{i=0}^8 d_i \right) = 1, \quad \sum_{i=0}^k c_{k-i} d_i = 0, \quad k > 0,$$

and relations (2.1) and (2.2) when $q = 2$, $n = 6$, we obtain the following values of d_i , $1 \leq i \leq 8$.

$$\begin{aligned}
d_1 &= -6\sigma_1, \\
d_2 &= 20\bar{\sigma}_2 + 18\sigma_2, \\
d_3 &= -50\bar{\sigma}_3 - 82\sigma_2\sigma_1, \\
d_4 &= 105\bar{\sigma}_4 + 237\sigma_2\bar{\sigma}_2 + 129\sigma_2^2, \\
d_5 &= -546\sigma_2\bar{\sigma}_3 - 504\sigma_2^2\sigma_1, \\
d_6 &= 1323\sigma_2^2\bar{\sigma}_2 + 595\sigma_2^3, \\
d_7 &= -2100\sigma_2^3\sigma_1, \\
d_8 &= 2100\sigma_2^4.
\end{aligned}$$

Then from [4, Theorem 4.5.1], the Pontryagin class, $p_4(\nu)$, of ν can be computed from the formula

$$p_4(\nu) = 2d_8 - 2d_7d_1 + 2d_6d_2 - 2d_5d_3 + d_4^2$$

to give that $p_4(\nu) = -81\sigma_2^4$.

Now let $\chi = a\bar{\sigma}_4 + b\sigma_2\bar{\sigma}_2 + c\sigma_2^2$ be an element of $H^8(F(2,4); Z)$; then

$$\chi^2 = (a^2 + b^2 + c^2)\sigma_2^4.$$

Then identity (1.1) implies that $a^2 + b^2 + c^2 = -81$. This quadratic equation has no integral solution. Thus nonimmersion in codimension 8 is established. Therefore, together with [3], we have

PROPOSITION 2.3. *The sixteen dimensional manifold $F(2,4)$ does not immerse in codimension 8 but does immerse in codimension 15.*

3. $F(2,5)$.

PROPOSITION 3.1. *The twenty dimensional manifold $F(2,5)$ does not immerse in codimension 10 but does immerse in codimension 19.*

PROOF. The method of proof is completely the same as for Proposition 2.3 and the details are, therefore, omitted. Note that the Chern classes of $F(2,5)$ are contained in [6, Proposition 3.3] and if $F(2,5)$ immerses in codimension 10, one finds that the Pontryagin class $p_5(\nu)$ of its stable normal bundle ν is equal to $-504\sigma_2^5$.

REFERENCES

1. A. Borel and F. Hirzebruch, *Characteristic classes and homogeneous spaces*. I, Amer. J. Math. **80** (1958), 458–538. MR **21** #1586.
2. Fred J. Connell, *Nonimmersions of low dimensional flag manifolds*, Proc. Amer. Math. Soc. **44** (1974), 474–478. MR **49** #8037.
3. M. Hirsch, *Immersions of manifolds*, Trans. Amer. Math. Soc. **93** (1959), 242–276. MR **22** #9980.
4. F. Hirzebruch, *Topological methods in algebraic geometry*, 3rd ed., Springer-Verlag, Berlin and New York, 1966. MR **34** #2573.
5. W. V. D. Hodge and D. Pedoe, *Methods of algebraic geometry*, Vol. II, Cambridge Univ. Press, New York, 1952. MR **13**, 972.
6. S. A. Ilori, *Canonical systems on flag manifolds*, Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. **17** (1974), 10–17.
7. Séminaire C. Chevalley, 2e année, *Anneaux de Chow et applications*, Secrétariat math., Paris, 1958. MR **22** #1572.
8. J. Tornehave, *Immersions of complex flag manifolds*, Math. Scand. **23** (1968), 22–26 (1969). MR **40** #4970.
9. E. Vesentini, *Construction géométrique des classes de Chern de quelques variétés de Grassmann complexes*, Colloque de topologie algébrique, Louvain, 1956, 97–120; Thone, Liège; Masson, Paris, 1957. MR **21** #1585.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IBADAN, IBADAN, NIGERIA