

ON ANALYTIC FUNCTIONS INTO l^p -SPACES

J. GLOBEVNIK¹

ABSTRACT. Let Δ be the open unit disc in the complex plane and let $1 \leq p < \infty$. We construct an example of a discontinuous function from Δ into l^p which is analytic as a function from Δ into $l^{p'}$ for any $p' > p$ and whose derivatives (into $l^{p'}$) of all orders are functions into l^p .

Throughout, we denote by Δ the open unit disc in the complex plane and by $\{e_i\}$ the canonical basis in the spaces l^p ($1 \leq p < \infty$). Instead of $\sum_{i=1}^{\infty}$ we write \sum . We denote by N (resp. N_0) the set of all positive (resp. nonnegative) integers.

Suppose that $\zeta \mapsto f(\zeta) = \sum \varphi_i(\zeta) e_i = (\varphi_i(\zeta))$ is a function from Δ into l^p . In general the analyticity of all φ_i is not sufficient for analyticity of f , as shown by R. Aron and J. Cima [1]. They ask whether the analyticity of all φ_i together with uniform boundedness of φ_i implies analyticity of f .

In the paper we construct a discontinuous function from Δ into l^p ($1 \leq p < \infty$) which is analytic on Δ as a function into $l^{p'}$ for any $p' > p$ and whose derivatives (into $l^{p'}$) of all orders are functions into l^p . In particular, this gives the negative answer to the question mentioned above.

LEMMA 1. *Let $1 \leq p < \infty$. There exists a sequence $\{\psi_i\}$ of complex-valued polynomials such that*

- (i) $(\psi_i^{(n)}(\zeta)) \in l^p$ ($\zeta \in \Delta, n \in N_0$);
- (ii) *the sum of the series $\sum |\psi_i(\zeta)|^p$ is not locally bounded at the point $\frac{1}{2}$.
(As usual $\psi^{(n)}$ ($n \geq 1$) are the derivatives of a function ψ and $\psi^{(0)} = \psi$.)*

PROOF. (See [1]; the idea about derivatives is due to Professor R. Aron.) Let $\{\theta_n\}$, $\theta_n < \frac{1}{2}$ ($n \in N$) be a strictly decreasing sequence of positive numbers, converging to 0. Denote $B_n = \{re^{i\theta} : \frac{1}{4} < r < 1, \theta_n < \theta < \theta_{n+1}\}$ ($n \in N$). It is clear that it is possible to choose the sequence θ_n in such a way that for each $n \in N$ the distance between B_n and the real axis is greater than $\frac{1}{2^n}$. Now choose a sequence of points $\{\zeta_n\}$, $\lim \zeta_n = \frac{1}{2}$, such that $\zeta_n \in B_n$ ($n \in N$). By the Runge approximation theorem there exists for each $i \in N$ a polynomial ψ_i with the property that

$$\begin{aligned} |\psi_i(\zeta_i)| &\geq i, \\ |\psi_i(\zeta)| &\leq 1/i! \end{aligned} \quad (\zeta \in \Delta, \zeta \notin B_i, i \in N).$$

Received by the editors September 23, 1974.

AMS (MOS) subject classifications (1970). Primary 30A96, 46E30.

¹ This work was supported in part by the Boris Kidrič Fund, Ljubljana, Yugoslavia.

Copyright © 1977, American Mathematical Society

Let ζ be any point in Δ which does not lie on the real axis. Then there exists a disc, say of radius $\delta > 0$, centered at ζ and contained in Δ , which intersects at most two domains, say B_{i_0}, B_{i_0+1} . So if $n \in N_0$ we have by the Cauchy estimates

$$\begin{aligned} |\psi_i^{(n)}(\zeta)| &\leq \frac{n!}{\delta^n} \sup_{|\zeta-\eta|<\delta} |\psi_i(\eta)| \\ &\leq (n!/8^n) \cdot (1/i!) \quad (i \in N, i \neq i_0, i \neq i_0 + 1). \end{aligned}$$

Hence for each such ζ the series $\sum |\psi_i^{(n)}(\zeta)|^p$ converges for all $n \in N_0$. Now let ζ lie on the real axis. For each $i \in N$ the disc $\{\eta: |\zeta - \eta| < \frac{1}{2}\}$ does not intersect B_i by the assumption. By the Cauchy estimates it follows that

$$\begin{aligned} |\psi_i^{(n)}(\zeta)| &\leq \frac{n!}{(\frac{1}{2})^n} \sup_{|\zeta-\eta|<\frac{1}{2}} |\psi_i(\eta)| \\ &\leq n! \cdot 2^{in} \cdot (1/i!) \quad (i \in N, n \in N_0) \end{aligned}$$

which shows that the series $\sum |\psi_i^{(n)}(\zeta)|^p$ converges for all $n \in N_0$. Hence (i) is satisfied. Further, since $\sum |\psi_i(\zeta_k)|^p \geq k^p$ ($k \in N$) and since $\lim \zeta_k = \frac{1}{2}$ it follows that the sum of the series $\sum |\psi_i(\zeta)|^p$ is not locally bounded at $\frac{1}{2}$, hence (ii) is also satisfied. Q.E.D.

LEMMA 2. *Let $\sum a_i, \sum b_i$ be two series with real nonnegative terms. Let $\{p_i\}, \{q_i\}$ be two increasing sequences of positive integers. Suppose that $\sum_{j=1}^{p_i} a_j = \sum_{j=1}^{q_i} b_j$ ($i \in N$). If $\sum a_j$ converges, then $\sum b_j$ also converges and both series have the same sum.*

This is easily established by standard reasoning.

DEFINITION. Let $\{a_i\}$ be a sequence of complex numbers and let $\{p_i\}$ be a sequence of positive integers. Define the sequence $\{b_i\}$ as follows:

$$b_i = \frac{a_n}{p_n} \quad \left(\sum_{j=1}^{n-1} p_j < i \leq \sum_{j=1}^n p_j, n \in N \right).$$

Then we say that the sequence $\{b_i\}$ is obtained by $\{p_i\}$ -splitting of the sequence $\{a_i\}$ and we write $\{b_i\} = \text{spl}(\{a_i\}, \{p_i\})$.

LEMMA 3. *Let $1 \leq p < \infty$. There exists a sequence $\{\varphi_i\}$ of complex-valued polynomials such that*

- (i) $|\varphi_i^{(n)}(\zeta)| \leq 1$ ($i \in N, n \in N_0, \zeta \in \Delta$);
- (ii) $(\varphi_i^{(n)}(\zeta)) \in l^p$ ($n \in N_0, \zeta \in \Delta$);
- (iii) the sum of the series $\sum |\varphi_i(\zeta)|^p$ is not locally bounded at $\frac{1}{2}$.

PROOF. Let $\{\psi_i\}$ be the sequence of polynomials given by Lemma 1. Since ψ_i is a polynomial for each $i \in N$ it follows that there exists a sequence of numbers $\{M_i\}$ such that

$$|\psi_i^{(n)}(\zeta)| \leq M_i \quad (i \in N, n \in N_0, \zeta \in \Delta).$$

It is clear that there exists a sequence of numbers $\{N_i\}$ such that

- (i') $N_i \geq 1$, $N_i \geq M_i$ ($i \in N$),
(ii') the series $\sum (M_i/N_i)^p$ converges.

For each $i \in N$ let k_i be the largest integer less than or equal to N_i^p . Since $N_i \geq 1$ ($i \in N$) we have $k_i \geq 1$ ($i \in N$). Define

$$\{\varphi_i(\xi)\} = \text{spl}((\{k_i/N_i\}\psi_i(\xi)), \{k_i\}) \quad (\xi \in \Delta).$$

It is clear that all φ_i are analytic on Δ and by (i') we have $|\varphi_i^{(n)}(\xi)| \leq 1$ ($i \in N, n \in N_0, \xi \in \Delta$), hence (i) is satisfied. Consider the series $\sum |\varphi_i^{(n)}(\xi)|^p$. We have

$$\sum_{j=1}^m k_i |\varphi_j^{(n)}(\xi)|^p = \sum_{i=1}^m \frac{k_i}{N_i^p} |\psi_i^{(n)}(\xi)|^p \quad (m \in N, n \in N_0, \xi \in \Delta).$$

Now, $\sum |\psi_i^{(n)}(\xi)|^p$ converges for each $n \in N_0, \xi \in \Delta$. Since $0 < k_i/N_i^p \leq 1$ ($i \in N$) it follows that the series $\sum (k_i/N_i^p)|\psi_i^{(n)}(\xi)|^p$ also converges for each $n \in N_0, \xi \in \Delta$. By the last equality Lemma 2 implies that the series $\sum |\varphi_i^{(n)}(\xi)|^p$ converges for each $n \in N_0, \xi \in \Delta$, hence (ii) is satisfied. By the same equality Lemma 3 also implies that

$$\sum |\varphi_i(\xi)|^p = \sum \frac{k_i}{N_i^p} |\psi_i(\xi)|^p \quad (\xi \in \Delta).$$

Now we have

$$\begin{aligned} \left| \sum |\psi_i(\xi)|^p - \sum |\varphi_i(\xi)|^p \right| &= \sum \left(1 - \frac{k_i}{N_i^p} \right) |\psi_i(\xi)|^p \\ &= \sum \frac{N_i^p - k_i}{N_i^p} |\psi_i(\xi)|^p \leq \sum \frac{1}{N_i^p} \cdot M_i^p \quad (\xi \in \Delta), \end{aligned}$$

where the last series converges by (ii'). Since the sum of $\sum |\psi_i(\xi)|^p$ is not locally bounded at $\frac{1}{2}$, it follows that also the sum of $\sum |\varphi_i(\xi)|^p$ is not locally bounded at $\frac{1}{2}$, hence (iii) is also satisfied. Q.E.D.

THEOREM. Let $1 \leq p < \infty$. There exists a function from Δ into l^p with the following properties:

- (i) it is not continuous;
- (ii) it is analytic on Δ as a function into $l^{p'}$ for any $p' > p$;
- (iii) its derivatives (into any $l^{p'}$) of all orders are functions from Δ into l^p .

PROOF. Let $\{\varphi_i\}$ be the sequence of functions from Lemma 3. Define

$$\{\psi_i(\xi)\} = \text{spl}((2^{i(1-1/p)}\varphi_i(\xi)), \{2^i\}) \quad (i \in N, \xi \in \Delta).$$

Clearly the functions ψ_i are again analytic on Δ . Let $\alpha > 0$. We have

$$\sum_{j=1}^n 2^j |\psi_j(\xi)|^{p+\alpha} = \sum_{i=1}^n 2^{-i\alpha/p} |\varphi_i(\xi)|^{p+\alpha} \quad (n \in N, \xi \in \Delta).$$

Since $|\varphi_i(\xi)| \leq 1$ ($\xi \in \Delta$) it follows that the convergent series $\sum 2^{-i\alpha/p}$ is majorant for $\sum 2^{-i\alpha/p} |\varphi_i(\xi)|^{p+\alpha}$ as $\xi \in \Delta$. Consequently the latter series

converges and its sum is bounded on Δ . By Lemma 2 the last equality implies that the series $\sum |\psi_i(\xi)|^{p+\alpha}$ converges to the same sum. Hence the sum of $\sum |\psi_i(\xi)|^{p+\alpha}$ is bounded on Δ and by Aron-Cima theorem (see [1]) $\xi \mapsto (\psi_i(\xi))$ is an analytic function from Δ into $l^{p'}$ for all $p' > p$. Next we show that $(\psi_i^{(n)}(\xi)) \in l^p$ ($n \in N_0, \xi \in \Delta$). By the definition of ψ_i we have

$$\sum_{j=1}^{\sum_{i=1}^m 2^i} |\psi_j^{(n)}(\xi)|^p = \sum_{j=1}^m |\varphi_j^{(n)}(\xi)|^p \quad (m \in N, n \in N_0, \xi \in \Delta)$$

and the conclusion follows by Lemma 3 from the fact that $(\varphi_i^{(n)}(\xi)) \in l^p$ ($n \in N_0, \xi \in \Delta$). Finally by Lemma 2 we also have

$$\sum |\psi_i(\xi)|^p = \sum |\varphi_i(\xi)|^p \quad (\xi \in \Delta)$$

which shows that the function $\xi \mapsto (\psi_i(\xi))$ is not continuous from Δ into l^p since the sum of $\sum |\varphi_i(\xi)|^p$ is not locally bounded at $\frac{1}{2}$. Hence the function $\xi \mapsto (\psi_i(\xi))$ has all required properties. Q.E.D.

REMARK. By observing that $\sum |\psi_i(\xi)|^{p'}$ above converges uniformly on Δ if $p' > p$ it is easy to see that the Theorem still holds if we replace $l^{p'}$ ($p' \geq p$) by $L^{p'}(S, \Sigma, \mu)$ ($p' \geq p$) provided that Σ contains a disjoint sequence $\{E_n\}$ of sets of measure 1. In this case we define a sequence $\{x_i\}$ of unit vectors in $L^{p'}(S, \Sigma, \mu)$ ($p' \geq p$) by

$$x_i(s) = \begin{cases} 1 & \text{if } s \in E_i, \\ 0 & \text{if } s \notin E_i \end{cases}$$

and define our function as $\xi \mapsto \sum \psi_i(\xi) x_i$.

ACKNOWLEDGEMENT. The result presented in the paper was obtained during the author's stay at the University of Kentucky. The author wishes to express his thanks for several useful discussions to Professors L. Harris, M. Schottenloher, T. Suffridge and specially to Professor R. Aron who also read the manuscript and gave some helpful remarks.

REFERENCES

1. R. Aron and J. A. Cima, *A theorem on holomorphic mappings into Banach spaces with basis*, Proc. Amer. Math. Soc. **36** (1972), 289–292. MR **47** # 5594.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KENTUCKY, LEXINGTON, KENTUCKY 40506

INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, UNIVERSITY OF LJUBLJANA, LJUBLJANA,
YUGOSLAVIA (Current address)