

## SCALAR CURVATURES ON $O(M)$ , $G_2(M)$

ALCIBIADES RIGAS<sup>1</sup>

**ABSTRACT.** We show that every  $C^\infty f: G_2(M) \rightarrow \mathbf{R}$ ,  $M^n$  a compact connected riemannian manifold  $n \geq 3$ , is the scalar curvature function of some complete riemannian metric on  $G_2(M)$ , the grassmann bundle of 2 planes over  $M$ , except possibly when  $K = \text{constant} \geq 0$ . A similar result holds for  $O(M)$  bundle of orthonormal frames on  $M$ .

This note is an application of Theorem A of [3] and O'Neill's formula for the curvature of a riemannian submersion [5], [4] and [1]. Theorem C of [3] gives an affirmative answer to the question described in the abstract if  $f(P) < 0$  for at least one 2-plane section  $P$  tangent to  $M$ .

**Preliminaries.** Let  $(M^n, ds^2)$  be a compact riemannian manifold and let  $O(M)$  be the principal  $O(n)$  bundle of  $ds^2$ -orthonormal frames on  $M$ . Choose a connection of  $\pi: OM \rightarrow M$ .

We assume the setting of [1, Section 1]. (See also [4].) Briefly, let  $\langle, \rangle$  be the bi-invariant metric on  $O(n)$  defined via the positive definite  $-B$  (killing form) on the lie algebra  $\hat{G}$  of  $O(n)$ , and  $\gamma$  the connection form for  $\pi$ . Split each  $X \in T(OM)$  into horizontal and vertical components relative to  $\gamma$ :  $X = X^H + X^V$ . For  $t > 0$ ,  $g_t = \pi^* ds^2 + t^2 \langle \gamma, \gamma \rangle$  is a family of  $O(n)$ -right invariant complete riemannian metrics on  $P$  and relative to each one  $\pi$  is a riemannian submersion on  $(M, ds^2)$ . Fix a  $t > 0$  and let  $g \equiv g_t$ . Let  $U$  be open in  $M$  and  $X'_1, \dots, X'_n$  be a  $ds^2$ -orthonormal (o.n. from now on) frame on  $U$ . Consider reductive decomposition of  $\hat{G} = \hat{N} + \hat{H}$ , orthogonal relative to  $-B$  where  $\hat{H}$  is the lie algebra of  $O(2) \times O(n-2)$ , and  $\hat{N}$  is the space of skew symmetric matrices of the form:

$$\begin{pmatrix} 0 & 0 & -\xi^t \\ 0 & 0 & -\eta^t \\ \xi & \eta & 0 \end{pmatrix}, \quad \xi, \eta \text{ column vectors in } \mathbf{R}^{n-2},$$

(see [2, vol. II, p. 280]).

Let  $e'_1, e'_2, \dots, e'_n$ ,  $r, s = 1, \dots, n-2$ , be the obvious  $(-B)$ -o.n. basis of  $\hat{N}$ , where

Received by the editors December 29, 1975.

AMS (MOS) subject classifications (1970). Primary 53C20; Secondary 55F10.

**Key words and phrases.** Riemannian metric, bundle of orthonormal frames, scalar and sectional curvature, grassmann 2-plane bundle.

<sup>1</sup> Research partially supported by an NSF grant.

Copyright © 1977, American Mathematical Society

$e_1^r$  is the matrix we obtain above if  $\xi$  is the  $r$ th standard basis element of  $R^{n-2}$  and  $\eta = 0$ . Similarly with  $e_2^s$ . Let  $\hat{H} = \hat{H}_1 \dot{+} \hat{H}_{n-2}$  with  $\hat{H}_1 = \{(\begin{smallmatrix} 0 & -r \\ r & 0 \end{smallmatrix}), r \in R\}$  the lie algebra of  $O(2)$  and  $\hat{H}_{n-2}$  the lie algebra of  $O(n-2)$ .

Let  $X'_{n+1}, \dots, X'_{3n-4}; X'_{3n-3}; X'_{3n-2}, \dots, X'_{n+r}$  be notation for an o.n. basis of  $\hat{N} \dot{+} \hat{H}_1 \dot{+} \hat{H}_2$ , where  $r = \dim O(n)$ . If  $A \rightarrow A^*$  denotes the lie algebra monomorphism  $\hat{G} \rightarrow \mathfrak{X}(OM)$ , the vector fields of  $O(M)$ , determined by the free  $O(n)$  action, we have that  $X_1, \dots, X_n; t^{-1}X_{n+1}^*, \dots, t^{-1}X_{n+r}^*$  form a local  $g$ -o.n. basis on  $\pi^{-1}(U)$ , where  $X_i$  is the  $\gamma$ -horizontal lift of  $X'_i$ .

Consider now the following

$$\begin{array}{ccc} O(M), & g \xrightarrow{p} & \frac{O(M)}{O(2) \times O(n-2)} \equiv G_2(M), \hat{g} \\ \pi \downarrow & & \downarrow q \\ M, ds^2 & = & M, ds^2 \end{array}$$

where  $\hat{g}$  is the metric on  $G_2(M)$  with respect to which  $p$  is a riemannian submersion. Let  $u \in \pi^{-1}(U) \subset O(M)$ ,  $p(u) \in G_2(M)$ ,  $x = \pi(u) \in M$ . We want to calculate  $S_{\hat{g}}(p(u))$ : the scalar curvature at  $p(u)$  relative to  $\hat{g}$ .

A  $\hat{g}$ -o.n. frame at  $p(u)$  is obtained by projecting a  $g$ -o.n.,  $p$ -horizontal (i.e., normal to the  $p$ -fibre relative to  $g$ ) frame at  $u$  of  $O(M)$ . From  $qp = \pi$  follows that  $\text{Horiz}(p) = \text{Horiz}(\pi) \dot{+} \text{span}\{X_{n+1}, \dots, X_{3n-4}\}$ ,  $X_{n+s} = t^{-1}X_{n+s}^*$  (sum orthogonal relg). From now on let  $1 \leq \alpha, \beta \leq n+r$ ,  $3n-3 \leq a \leq n+r$ ,  $n+1 \leq b, \eta, \theta \leq n+r$ ,  $n+1 \leq \lambda, \mu \leq 3n-4$ ,  $1 \leq i, j, k \leq n$ .

Let  $\bar{K}_{\alpha\beta} \equiv K_{\hat{g}}(p_*(X_\alpha), p_*(X_\beta))$  where  $X_\alpha$  is one of the  $X_i$ 's or  $t^{-1}X_\lambda^*$ 's and  $K_{\hat{g}}$  is the sectional curvature relative to  $\hat{g}$ . By O'Neill's formula for the curvature of a riemannian submersion ([5], [4], [1]),

$$(1) \quad \bar{K}_{\alpha\beta} = K_g(X_\alpha, X_\beta) + \frac{3}{4} \|[X_\alpha, X_\beta]^V\|^2,$$

where  $V$  stands for "vertical part" or " $g$ -orthogonal projection onto the  $p$ -fibre" in any  $T_u(OM)$ . Notice that

$$\begin{aligned} [X_i, X_j]^V &= \sum_a g([X_i, X_j], X_a) X_a = \sum_a t^2 \langle \gamma[X_i, X_j], \gamma(X_a) \rangle X_a \\ &= \sum_a \langle \gamma[X_i, X_j], X'_a \rangle X_a^*. \end{aligned}$$

If  $\| \cdot \|$  is the length relative to  $g$ ,

$$\|[X_i, X_j]^V\|^2 = t^2 \sum_a \langle \gamma_{ij}, X'_a \rangle^2,$$

where  $\gamma_{ij} = \gamma[X_i, X_j] \in \hat{G}$ .

Let  $\langle \gamma_{ij}, X'_a \rangle(u) \equiv 2H_{ij}^a(u)$  and obtain

$$(2) \quad \bar{K}_{ij} = K_{ij} + 3t^2 \sum_a (H_{ij}^a)^2$$

Similarly  $\bar{K}_{\lambda\lambda} \equiv K_g(p_*(X_\lambda), p_*(X_\lambda)) = K(X_\lambda, X_\lambda) + \frac{3}{4} \|[X_i, X_\lambda]^V\|^2$ . Here  $X_i$  is  $\pi$ -

horizontal and  $X_\lambda$  is a fundamental  $p$ -vertical.  $\therefore [X_i, X_\lambda]$  is zero.

$$(3) \quad \bar{K}_{i\lambda} = K(X_i, X_\lambda).$$

Now,  $\bar{K}_{\lambda,\mu} = K(X_\lambda, X_\mu) + \frac{3}{4} \|[X_\lambda, X_\mu]^V\|^2$ . Recall that  $X_\lambda = t^{-1} X_\lambda^*$  and the basis  $X'_\lambda$  was exactly the  $e_1^r$ 's and  $e_2^s$ 's above. It is  $[e_1^r, e_1^s] = [e_2^r, e_2^s] = 0$  and

$$[e_1^r, e_2^s] = \delta_{rs} \begin{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \\ 0 \end{pmatrix} \quad \text{in } \hat{H}_1 \subset \hat{G}.$$

If  $-B$  is so normalized as to have  $e_1^r, e_2^s$  of length one then  $[e_1^r, e_2^s]$  is also of length 1.

$$[X_\lambda, X_\mu] = [t^{-1} X_\lambda^*, t^{-1} X_\mu^*] = t^{-2} [X_\lambda^*, X_\mu^*].$$

From the above discussion  $\|[X_\lambda, X_\mu]^V\|^2 = (t^{-2})^2 t^2 = 1$  or it is zero depending on  $\lambda, \mu$ . The number of nonzero ones is  $n - 2$ .

From this and (2), (3) we have:

$$(4) \quad \begin{aligned} \sum_{\alpha, \beta} \bar{K}_{\alpha, \beta} &= \sum_{i < j} \bar{K}_{ij} + \sum_{i, \lambda} \bar{K}_{i\lambda} + \sum_{\lambda < \mu} \bar{K}_{\lambda, \mu} \\ &= \sum_{i < j} \left\{ K_{ij} + 3t^2 \sum_a (H_{ij}^a)^2 \right\} + \sum_{i, \lambda} K_{i\lambda} \\ &\quad + \sum_{\lambda < \mu} K_{\lambda\mu} + \frac{3}{4} t^{-2} (n - 2). \end{aligned}$$

Now from O'Neill's theorem [5] on the submersion  $\pi$  one can express  $K_{ij}, K_{i\lambda}, K_{\lambda\mu}$  relative to the curvature of  $M$  and the  $\pi$ -fibre  $O(n)$ .

$$K_{ij} \equiv K(X_i, X_j) = K_{ds^2}(X'_i, X'_j) - \frac{3}{4} \|[X_i, X_j]^W\|^2$$

where  $W$  denotes the vertical component relative to the  $\pi$ -fibre:

$$\begin{aligned} [X_i, X_j]^W &= \sum_b g([X_i, X_j], X_b) X_b \\ &= \sum_b \langle \gamma[X_i, X_j], X'_b \rangle X_b^* = 2 \sum_b H_{ij}^b X_b^*, \\ \|[X_i, X_j]^W\|^2 &= 4t^2 \sum_b (H_{ij}^b)^2, \quad (H_{ij}^b \equiv H_{ij}^b(u)). \end{aligned}$$

$$(5) \quad K_{ij} = K_{ds^2}(X'_i, X'_j) - 3t^2 \sum_b (H_{ij}^b)^2.$$

$K_{i\lambda} = t^2 \|(\nabla_{X_i} X_\lambda)^{\text{hor}}\|^2$  ([5], [4]) where  $\nabla$  is the riemannian connection corresponding to  $g$ , and hor stands for  $\pi$ -horizontal component. It is easy to see from [5, Lemma 2, p. 446] or [4], that

$$\begin{aligned}
 \|\langle \nabla_{X_i} X_\lambda \rangle^{\text{hor}}\|^2 &= \sum_j \frac{1}{2} g([X_i, X_j], X_\lambda)^2 \\
 (6) \qquad \qquad \qquad &= \frac{1}{4} \sum_j (2H_{ij}^\lambda)^2, \\
 K_{i\lambda} &= t^2 \sum_j (H_{ij}^\lambda)^2.
 \end{aligned}$$

*Note.* This is the same result as in Proposition 5 of [1] where  $K_{i\lambda} \equiv R_{i\lambda i\lambda} \cdot K_{\lambda\mu} = K(X_\lambda, X_\mu)$  and it must be the curvature of the same section considered as tangent to the totally geodesic  $\pi$ -fibre, i.e.,

$$K_{\lambda\mu} = \frac{1}{4} t^{-2} \|\langle X_\lambda^*, X_\mu^* \rangle\|^2$$

where  $\|\cdot\|$  is the original  $O(n)$  norm. (This agrees with Proposition 5 of [1]:

$$K_{\lambda\mu} = R_{\lambda\mu\lambda\mu} = \frac{1}{4} t^{-2} \sum_f (C_{\mu f}^\lambda)^2,$$

where  $C_{st}^r = \langle [X_s, X_t], X_r \rangle = C_{rt}^s$ , etc.) But  $\|\langle X_\lambda^*, X_\mu^* \rangle\| = 1$  or  $0$  as above and there are exactly  $(n - 2)$  combinations  $(\lambda < \mu)$  that give us 1:

$$(7) \qquad \qquad \qquad \sum_{\lambda < \mu} K_{\lambda\mu} = \frac{1}{4} t^{-2} (n - 2).$$

From (4), (5), (6), (7) we have:

$$\begin{aligned}
 \sum_{\alpha, \beta} \bar{K}_{\alpha\beta} &= \sum_{i < j} K_{ds^2}(X'_i, X'_j) - \sum_{i < j} 3t^2 \sum_b (H_{ij}^b)^2 \\
 &+ \sum_{i < j} 3t^2 \sum_a (H_{ij}^a)^2 + \sum_{i, \lambda} t^2 \sum_j (H_{ij}^\lambda)^2 \\
 &+ \frac{1}{4} t^{-2} (n - 2) + \frac{3}{4} t^{-2} (n - 2).
 \end{aligned}$$

Therefore,

$$S_{\hat{g}}(p(u)) = S_{ds^2}(\pi(u)) - 2t^2 \sum_\lambda \sum_{i < j} (H_{ij}^\lambda(u))^2 + t^{-2}(n - 2)$$

after collecting terms and observing the ranges of the indices  $a, b$ , and  $\lambda$ . ( $S_{ds^2}(\pi(u))$  is the scalar curvature of  $ds^2$  on  $M$  at  $\pi(u)$ .)

$$\Lambda(U) \equiv 2 \sum_\lambda \sum_{i < j} (H_{ij}^\lambda(u))^2 \geq 0 \quad \text{for all } u \in \pi^{-1}(U).$$

Let  $S$  be the notation for the scalar curvature of  $ds^2$  on  $M$ ,  $S: M \rightarrow R$  and  $\Lambda: O(M) \rightarrow R$  but it factors through  $M$  by the  $O(n)$ -invariance of  $g$ . So, instead of  $\Lambda(u)$  we write  $\Lambda(x)$ ,  $x = \pi(u)$ .

We proved:

**PROPOSITION.** *The scalar curvature of  $\hat{g}$  at  $p(u)$  is equal to  $S(x) - t^2 \Lambda(x) + t^{-2}(n - 2)$  where  $x = \pi(u)$ , and  $\Lambda(x) \geq 0$ .*

**THEOREM.** *A  $C^\infty$  function  $f: G_2(M) \rightarrow R$  is a scalar curvature function for some riemannian metric on  $G_2(M)$  except perhaps when it is a nonnegative constant.*

The proof of this theorem is an application of

**THEOREM A ([3]).** *Let  $(N, g)$  be a smooth compact riemannian manifold with gaussian (resp. scalar if  $\dim N \geq 3$ ) curvature  $S$  and let  $f \in C^\infty(N)$ . If there is a constant  $c > 0$  such that*

$$\min cf < S(x) < \max cf$$

*for all  $x \in N$ , then there is a smooth metric  $g_1$  on  $N$  with gaussian (resp. scalar) curvature  $f$ .*

Recall that  $2H_{ij}^\lambda = \langle \gamma[X_i, X_j], X'_\lambda \rangle$  and  $X_i, X_j$  are  $\gamma$ -horizontal vectors in  $T(OM)$  of unit length relative to  $g \equiv g_t$  and therefore of unit length relative to every  $g_t, t > 0$ . The length of  $\gamma[X_i, X_j]$  in  $\hat{G}$  is bounded independent of  $t$ , i.e.,  $0 \leq \Lambda_m \leq \Lambda(x) \leq \Lambda_M$  for all  $x \in M$  with  $\Lambda_m, \Lambda_M$  constants independent of  $t$ .

Since  $S(x)$  is bounded, it follows that  $\hat{S}_m(t)/\hat{S}_M(t) \rightarrow 1$  as  $t \rightarrow 0$ , where  $\hat{S}_m(t)$  and  $\hat{S}_M(t)$  are the maximum and minimum of the scalar curvature of  $\hat{g}_t$ , and the proof is complete.

*Special case.* If  $f = \text{constant} > 0$  is the sectional curvature of  $(M, ds^2)$  then  $M = S^n/\Gamma, \Gamma \subset O(n + 1)$  and  $O(M) = O(n + 1)/\Gamma$ , where  $\Gamma$  is a finite subgroup of  $O(n + 1)$  [6, p. 69]. If  $M$  is 1-connected,  $M = S^n$  of radius  $f^{-1/2}$  and therefore  $G_2(M) = O(n + 1)/O(2) \times O(n - 2)$ , which admits a constant scalar curvature function as a homogeneous space of  $O(n + 1)$ . Any  $C^\infty$  function from  $G_2(M) \rightarrow R$  will then be a scalar curvature for some riemannian metric on  $G_2(M)$ , by Theorem C of [3].

The following is proved exactly the same way as the above theorem.

**PROPOSITION.** *A  $C^\infty$   $f: O(M) \rightarrow R$  is a scalar curvature function for some riemannian metric on  $O(M)$ , except perhaps when  $f = \text{constant} \geq 0$ .*

**PROOF.** Using the same notation conventions,

$$K_{ij} = K_{ds^2}(X'_i, X'_j) - 3t^2 \sum_{\eta} (H_{ij}^\eta)^2,$$

$$K_{i\theta} = t^2 \sum_j (H_{ij}^\theta)^2,$$

$$K_{\eta\theta} = \frac{1}{4}t^{-2} \langle [X_\eta^*, X_\theta^*], [X_\eta^*, X_\theta^*] \rangle = t^{-2} K_0(X'_\eta, X'_\theta)$$

where  $K_0$  is the sectional curvature of the fibre  $O(n)$  in its original metric  $\langle , \rangle$ . Therefore,  $S_g(u) = S(x) - t^2 \Lambda_1(u) + t^{-2}c$  where  $S_g(u)$  is the scalar curvature of  $O(M)$  relative to the metric  $g \equiv g_t$  at  $u$ ,

$$\Lambda_1(u) = 2 \sum_{i < j} \sum_{\eta} (H_{ij}^\eta(u))^2 \geq 0$$

bounded independent of  $t$  and  $c$  is the constant positive scalar curvature of  $(O(m), \langle \cdot, \cdot \rangle)$ .

If  $0 \leq f_m < f_M$  by the exact same procedure as in the theorem we obtain that  $f$  is a scalar curvature on  $O(M)$ .

In the particular case that  $f$  is  $K \circ p$  with  $K$  the constant positive curvature of  $M$ , then  $M = S^n/\Gamma$  and  $O(M) = O(n+1)/\Gamma$  is a homogeneous space of a compact lie group that admits a metric with positive constant scalar curvature. By Theorem C of [3], all  $C^\infty$  functions on  $O(M)$  are scalar curvatures and in particular  $f$ .

#### REFERENCES

1. G. Jensen, *Einstein metrics on principal fibre bundles*, J. Differential Geometry **8** (1973), 599–614. MR **50** #5694.
2. S. Kobayashi and K. Nomizu, *Foundations of differential geometry*, Vol. II, Interscience, New York, 1969. MR **38** #6501.
3. J. Kazdan and F. Warner, *A direct approach to the determination of gaussian and scalar curvature functions*, Invent. Math. **28** (1975), 227–230.
4. H. B. Lawson and S. T. Yau, *Scalar curvature, non-abelian group actions, and the degree of symmetry of exotic spheres*, Comment. Math. Helv. **49** (1974), 232–244. MR **50** #11300.
5. B. O. O'Neill, *The fundamental equations of a submersion* Michigan Math. J. **13** (1966), 459–469. MR **34** #751.
6. J. Wolf, *Spaces of constant curvature*, 3rd ed., Publish or Perish, Cambridge, Mass., 1974.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WESTERN ONTARIO, LONDON, ONTARIO, CANADA

*Current address:* IMECC, UNICAMP, Campinas, S.P., Brazil