

A NOTE ON KC WALLMAN COMPACTIFICATIONS

DARRELL W. HAJEK AND ANGEL E. JIMÉNEZ

ABSTRACT. In a previous paper, D. W. Hajek showed that if a space X is a T_3 space and A is a compact subset of WX , the Wallman compactification of X , then $X \cap A$ is a closed subset of X . This raises the question of whether this "closed intersection" property characterizes the T_3 spaces among the Hausdorff spaces. In the present paper, the authors show this conjecture is false by giving an example of a nonregular Hausdorff space whose Wallman compactification is a KC (compact closed)-space, and, hence, trivially satisfies this "closed intersection" property.

All spaces considered in this paper will be T_1 spaces. In [1] it was shown that if X is a T_3 space and if A is a compact subset of WX , the Wallman compactification of X , then $A \cap X$ is a closed subset of X . It is then natural to ask whether T_3 spaces are the only Hausdorff spaces which have this property. In this paper we give an example of a nonregular Hausdorff space whose Wallman compactification is a KC-space (i.e. is a space in which every compact subset is closed).

Recall that for any space X the Wallman compactification WX consists of the set $\{\mu: \mu \text{ is an ultrafilter in the lattice of all closed subsets of } X\}$ with topology generated by $\{C(A) = \{\mu \in : A \in \mu\}: A \text{ is a closed subset of } X\}$ as a base for the closed subsets of WX . With this topology WX is a compact T_1 space which is Hausdorff if and only if X is T_2 , and the function $\varphi_X: X \rightarrow WX$ defined by $\varphi_X(x) = \{A: A \text{ is closed in } X \text{ and } x \in A\}$ is a dense embedding of X in WX . When no ambiguity can result it is common practice to ignore the distinction between X and $\varphi_X[X]$ and to speak of X as a subset of WX . We note that it is easily shown that if A is a closed compact subset of X then $\text{cl}_{WX}(A) = A$ and that for any closed subset A of X , $\text{cl}_{WX}(A) = C(A)$.

We denote by T the space whose elements are the points in the unit interval $[0, 1]$ and whose topology is generated by the usual open subsets of $[0, 1]$ together with $[0, 1] \sim \{1/n: n \text{ a natural number}\}$. It is clear that T is a Hausdorff space and that any open set containing the closed set $\{1/n\}$ must intersect every neighborhood of 0; so T is not regular (see [3, p. 86]).

Received by the editors August 4, 1975 and, in revised form, November 1, 1975.

AMS (MOS) subject classifications (1970). Primary 54D35; Secondary 54D10, 54D20, 54G20.

Key words and phrases. Wallman compactification, KC (compact closed)-space, "closed intersection" property, nonregular Hausdorff space.

Copyright © 1977, American Mathematical Society

THEOREM 1. *If A is a compact subset of WT then $A \cap T$ is a closed subset of T .*

PROOF. Suppose $A \cap T$ is not closed in T . Then $A \cap T$ has a limit point x in $T \sim A$. If $x \neq 0$ there is some number K such that $n > K$ implies that $(0, 1] \cap [x - 1/n, x + 1/n]$ is a closed compact subset of T , and, hence, of WT . This then implies that $\{A \cap [x - 1/n, x + 1/n]: n > K\}$ is a collection of closed subsets of A which has the finite intersection property and empty intersection. Since A is compact no such collection can exist; so A can have no limit points in T other than 0. We note that any element of $WT \sim T$ is clearly in $C(\{1/n\})$, and thus for any $\epsilon > 0$ the set $[0, \epsilon) \sim \{1/n\}$ is open both in T and in WT ; so if 0 is a limit point of A , there must be a decreasing sequence $\{a_n\}$ of elements of A converging to 0, none of the elements of which are in $\{1/n\}$. It is clear, however, that if $0 \notin A$, then $\{a_n\}$ is a closed subset of A which is not compact.

With this theorem the following corollary is immediate from [2, Theorem 1].

COROLLARY. *If $f: X \rightarrow T$ is a Wallman extendible function, then the Wallman extension is unique.*

Theorem 1 tells us that the T_3 spaces are not the only spaces which have closed intersections with compact subsets of their Wallman compactifications. However we promised more.

THEOREM 2. *Every compact subset of WT is closed.*

PROOF. Suppose A is a compact subset of WT which is not closed. Then A must have a limit point μ which is not in A . From Theorem 1 we know that $\mu \notin T$, and again we note that the only elements of $WT \sim T$ are in $C(\{1/n\}) \sim \{1/n\}$. We also note that the relative topology on $\{1/n\}$ is the discrete topology. Since $\{C(E): E \in \mu, E \subseteq \{1/n\}\}$ is a collection of closed subsets of WT whose intersection is $\{\mu\}$, it is immediate that there is some subset B of $\{1/n\}$ such that $B \in \mu$ and $C(B)$ is disjoint from A . But since B is an infinite subset of a sequence, B itself must be a sequence. Then, as $A \cap T$ is closed in T , for each $x \in B$ there is an open subset \mathcal{U}_x of T such that \mathcal{U}_x is disjoint from A and $\mathcal{U}_x \cap \{1/n\} = \{x\}$. It is then elementary to show that $C(B) \cup (\cup \mathcal{U}_x)$ is an open subset of WT which contains μ and is disjoint from A .

We note that this process can be utilized in any compact Hausdorff space having a nonisolated first countable point to obtain the same results. It is not clear, however, whether a compact Hausdorff space with no first countable points must have a nonclosed subset which, when added to the closed sets yields a space with KC Wallman compactification or even the "closed intersection" property. The proof to Theorem 2 makes clear, however, that if a space is constructed by adding a set A to the closed subsets of a compact Hausdorff space and if the Wallman compactification of the subspace A is a KC-space, then the "closed intersection" property and having a KC Wallman compactification are equivalent.

We also note that [3, pp. 92–93] contains another example of a nonregular Hausdorff space which can be shown to have a KC Wallman compactification. Finally, the Euclidian plane with topology generated by the usual open sets together with all sets of the form $\{(0, 0)\} \cup \mathcal{U}$ where \mathcal{U} is obtained by deleting the positive y -axis from an open set of which it is a subset is an example of a Hausdorff space whose Wallman compactification does not have the “closed intersection” property.

BIBLIOGRAPHY

1. D. Hajek, *A characterization of T_3 spaces*, Indiana Univ. Math. J. **23** (1973/74), 23–25. MR **48** #4997.
2. ———, *Functions with continuous Wallman extensions*, Czechoslovak Math. J. **24** (99) (1974), 40–43.
3. L. A. Steen and J. A. Seebach, Jr., *Counterexamples in topology*, Holt, Rinehart and Winston, New York, 1970. MR **42** #1040.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PUERTO RICO, MAYAGUEZ,
PUERTO RICO 00708

DEPARTMENT OF MATHEMATICS, BOSTON UNIVERSITY, BOSTON, MASSACHUSETTS 02215