

ON A PROBLEM OF BRUCKNER AND CEDER  
CONCERNING  
THE SUM OF DARBOUX FUNCTIONS

J. S. LIPÍŃSKI

**ABSTRACT.** The main purpose of this paper is to show that for some continuous function  $f$  and any preassigned, countable and dense set  $D$  of real numbers there exists a measurable function  $d$  which takes on every real value in every interval such that the range of  $f + d$  is  $D$ .

A real valued function  $f$  defined on an interval  $I$  is said to have the intermediate value property if whenever  $x_1$  and  $x_2$  are in  $I$ , and  $y$  is any number between  $f(x_1)$  and  $f(x_2)$ , there is a number  $x_3$  between  $x_1$  and  $x_2$  such that  $f(x_3) = y$ . Because of Darboux's work on this property, one now usually calls a function having the intermediate value property a Darboux function. A function  $f$  is Darboux if and only if  $f$  maps any connected subset of  $I$  onto a connected set.

Since the sum of Darboux functions may fail to be a Darboux function, some mathematicians have examined how badly it can fail. Bruckner and Ceder [1] have recently shown that corresponding to each Darboux function  $f$  which is not constant on any subinterval, there exists a function  $d$  which takes on every real value in every subinterval of  $I$  such that the range of  $f + d$  is a preassigned countable and dense set. Certainly  $d$  is a Darboux function and the range of  $f + d$  is totally disconnected.

The technique used by Bruckner and Ceder in constructing the function  $d$  does not imply the measurability of  $d$ . Indeed Bruckner and Ceder have shown that  $d$  must be nonmeasurable whenever  $f$  is absolutely continuous. They asked whether  $d$  had to be nonmeasurable if we weaken absolute continuity of  $f$  to continuity of  $f$ .

This paper answers this question affirmatively for certain continuous functions  $f$ . This is a consequence of the following.

**THEOREM.** *Let  $g$  be a Darboux function which is not constant on any subinterval of its domain  $I$  such that the set*

$$A = \{\alpha: \alpha \in R, g^{-1}(\alpha) \text{ is perfect}\}$$

---

Received by the editors March 2, 1976.

AMS (MOS) subject classifications (1970). Primary 26A15; Secondary 26A21, 28A20.

Key words and phrases. Darboux functions.

© American Mathematical Society 1977

is dense in  $g(I)$ . Then there exists a homeomorphism  $h$  from  $I$  onto itself such that for every countable, dense set  $D \subset R$  there exists a function  $d$  which takes on every real value in every subinterval of  $I$  such that  $D$  is the range of  $(g \circ h) + d$  and  $(g \circ h) + d$  is constant almost everywhere.

It is known that there exist continuous functions  $g$  which are not constant on any interval such that  $A = g(I)$ . See for example [4]. (The properties of the set  $A$  for a Darboux function  $g$  are discussed in [2] and [3].) Let  $g$  be such a function. Now define  $f$  by  $f = g \circ h$  where  $h$  is the homeomorphism of the Theorem above. Then  $f$  is continuous and from this Theorem the range of  $f + d$  is  $D$ . By the Theorem  $f + d$  is constant almost everywhere. Thus  $d$  is measurable.

In order to prove the Theorem we shall use the type of construction suggested by Theorem 1 of [1]. But first we need the following.

**LEMMA.** Suppose  $F \subset I$ , and for every interval  $J \subset I$  there exists a perfect set  $P \subset F \cap J$ . Then there exists a homeomorphism  $\chi$  from  $I$  onto itself such that  $\chi(F)$  is measurable and  $|\chi(F)| = |I|$ .

**PROOF.** The Lemma is trivial if  $F = I$ . Suppose that  $I \setminus F \neq \emptyset$ . Let  $P^0$  be a perfect subset of  $F$ . Choose an increasing homeomorphism  $h_1$  from  $I$  onto itself so that  $|h_1(P^0)| > \frac{1}{2}|I|$ . Let  $\{(a_i, b_i)\}$  be a sequence which may be finite, of disjoint open intervals, such that  $I \setminus (\{a\} \cup \{b\} \cup h_1(P^0)) = \cup_i (a_i, b_i)$ . Choose a sequence  $\{I_i\}$  of nonoverlapping intervals so that  $\cup_i I_i = \cup_i \langle a_i, b_i \rangle$  and

$$(1) \quad |I_i| < \frac{1}{2}|I|.$$

Now choose for each  $i$  a perfect set  $P_i \subset h_1^{-1}(I_i) \cap F$ . Next choose a strictly increasing continuous function  $h_2$  from  $I$  onto itself such that

$$(2) \quad x \in h_1(P_0) \text{ implies } h_2(x) = x,$$

$$(3) \quad h_2(I_i) = I_i$$

and  $|h_2(h_1(P_i))| > \frac{1}{2}|I_i|$ .

By (2)  $x \in P^0$  implies  $h_1(x) = h_2(h_1(x))$ . Now put  $P^1 = P^0 \cup \cup_i P_i$ . Then  $P^1$  is a perfect set,  $P^0 \subset P^1 \subset F$ ,  $|h_2(h_1(P^1))| > \frac{3}{4}|I|$  and by (2), (3) and (1),  $|x - h_2(x)| < \frac{1}{2}|I|$ .

Using this method we can define by induction the sequences  $\{P^n\}$  of perfect sets and  $\{h_n\}$  of strictly increasing continuous functions from  $I$  onto itself such that

$$\begin{aligned} x = h_2(x) = h_3(x) = \dots & \quad \text{for } x \in h_1(P^0), \\ x = h_3(x) = h_4(x) = \dots & \quad \text{for } x \in h_2(h_1(P^1)) \end{aligned}$$

and generally,

$$(4) \quad x = h_{n+1}(x) = h_{n+2}(x) = \dots \quad \text{for } x \in h_n(h_{n-1}(\dots h_1(P^{n-1}) \dots)),$$

$$(5) \quad |x - h_n(x)| < (1/2^{n-1})|I|,$$

$$(6) \quad |h_n(h_{n-1}(\dots h_1(P^{n-1}) \dots))| > (2^n - 1/2^n)|I|.$$

By (5) we have  $|h_n(x) - h_{n+1}(h_n(x))| < (1/2^n)|I|$ . Hence

$$|h_n(h_{n-1}(\dots h_1(x) \dots)) - h_{n+1}(h_n(\dots h_1(x) \dots))| < (1/2^n)|I|.$$

By this inequality the sequence of superpositions  $\{h_n(h_{n-1}(\dots h_1(x) \dots))\}$  converges uniformly to a continuous function. Denote this limit function by  $\chi$ . All  $h_n$  are strictly increasing so  $\chi$  is increasing. By (4),

$$(7) \quad \chi(x) = h_n(h_{n-1}(\dots h_1(x) \dots)) \text{ for } x \in P^{n-1}.$$

Hence  $\chi$  is strictly increasing on  $P^{n-1}$  and also on the set  $\cup_{n=1}^\infty P^n$  which is dense in  $I$ . Thus  $\chi$  is strictly increasing. By (7)

$$\chi(P^{n-1}) = h_n(h_{n-1}(\dots h_1(P^{n-1}) \dots)).$$

Hence, since  $\chi(P^{n-1}) \subset \chi(F) \subset I$  by (6) we have  $|\chi(E)| > ((2^n - 1)/2^n)|I|$ . Hence  $|\chi(F)| = |I|$ .

This completes the proof of the Lemma.

**PROOF OF THE THEOREM.** Let  $\{I_n\}$  be a sequence of all the open intervals with rational endpoints contained in  $I$ . Since  $g(I_1)$  is a subinterval of  $g(I)$  and  $A$  is dense in  $g(I)$ , there exists a point  $\alpha_1 \in A \cap g(I_1)$ . Let  $F_1$  be a perfect set such that  $F_1 \subset I_1 \cap g^{-1}(\alpha_1)$  and  $F_1$  is nowhere dense in  $g^{-1}(\alpha_1)$ . Suppose we have chosen for each  $i = 1, 2, \dots, k$  perfect sets  $F_i$  and numbers  $\alpha_i$  such that  $F_i \subset I_i, g(F_i) = \alpha_i \in A, F_i$  is nowhere dense in  $g^{-1}(\alpha_i)$  and the sets  $F_i$  are disjoint. Let  $I^*$  be an interval such that  $I^* \subset I_{k+1}$  and  $I^* \cap (\cup_{i=1}^k F_i) = \emptyset$ . Choose a perfect set  $F_{k+1} \subset I^*$  such that  $g(F_{k+1}) = \alpha_{k+1} \in A, F_{k+1}$  is nowhere dense in  $g^{-1}(\alpha_{k+1})$ . Then  $F_{k+1} \cap F_i = \emptyset$  for  $i \neq k + 1$ . Clearly the sum  $F = \cup_{k=1}^\infty F_k$  satisfies the conditions of the Lemma. Then there is a homeomorphism  $\chi$  from  $I$  onto itself such that  $|\chi(F)| = |I|$ . Denote  $H = I \setminus \chi(F), h = \chi^{-1}$  and  $\Gamma = \{(x, y): x \in H, y + g(h(x)) \in D\}$ . Certainly  $\chi(F)$  is a set of the first category of Baire. Thus  $H$  is dense in  $I$ .

If  $B$  is a subset of  $H \times R$  we shall denote its  $x$ -projection, i.e. the set  $\{x: \exists y, (x, y) \in B\}$  by  $\text{dom } B$ , and its  $y$ -projection, i.e. the set  $\{y: \exists x, (x, y) \in B\}$  by  $\text{rng } B$ . For any  $t \in H$  we shall denote  $B_t = \{(x, y): x = t, (x, y) \in B\}$  and for any  $s \in R$  we shall denote  $B^s = \{(x, y): y = s, (x, y) \in B\}$ . Since  $\Gamma$  is the union of the graphs of all functions  $z_n - g(h(x))$  where  $z_n \in D$ , it follows that  $\text{rng } \Gamma = R$ .

We shall say that two points  $A(x', y')$  and  $B(x'', y'')$  of  $\Gamma$  are equivalent,  $A \sim B$ , if there exists a finite sequence of points  $A_i(x_i, y_i) \in \Gamma, i = 1, 2, \dots, n$ , such that  $A = A_1, B = A_n$ , and for each  $i$  we have either  $x_i = x_{i+1}$  or  $y_i = y_{i+1}$ . Denote by  $\Gamma/\sim$  the family of all equivalence classes for this equivalence relation. It is clear that for each  $t \in H, \text{rng } \Gamma_t$  contains a translation of  $D$ . Then  $\text{rng } \Gamma_t$  is a dense set. It follows from the definition of

equivalence that if  $G$  is an equivalence class, then  $G \cap \Gamma_i \neq \emptyset$  implies  $G \supset \Gamma_i$ . In the same way if  $G \cap \Gamma^s \neq \emptyset$  then  $G \supset \Gamma^s$ .

Now we shall prove that all  $\text{dom } \Gamma^s$  are dense in  $H$ . Since  $g$  is not constant on any interval then for any open interval  $I^* \subset I$  there exist points  $x'$  and  $x''$  in  $I^*$  for which  $g(h(x')) \neq g(h(x''))$ . Since  $D$  is dense there exists an  $r \in D$  such that  $r - s$  lies between  $g(h(x'))$  and  $g(h(x''))$ . Since  $g \circ h$  is Darboux there exists an  $x \in I^*$  for which  $g(h(x)) = r - s$ . If  $x \in H$  then  $(x, s) \in \Gamma$  and therefore  $x \in \text{dom } \Gamma^s \cap I^*$ .

If  $x \notin H$  then  $x \in \chi(F)$ . Thus  $r - s = g(h(x)) = g(\chi^{-1}(x)) = \alpha_i \in A$ . The level set  $g^{-1}(\alpha_i)$  is a perfect set and  $h(F) \cap g^{-1}(\alpha_i) = h(F_i) \cap g^{-1}(\alpha_i)$ . The set  $h(F_i)$  is nowhere dense in  $g^{-1}(\alpha_i)$ . It follows that there exists a point  $x^* \in I^* \cap [g^{-1}(\alpha_i) \setminus h(F)]$ . Thus  $x^* \in H \cap I^*$  and  $\alpha_i = g(h(x^*)) = r - s$ . Hence  $(x^*, s) \in \Gamma$  and  $x^* \in \text{dom } \Gamma^s \cap I^*$ . This shows that  $\text{dom } \Gamma^s$  is dense in  $H$ .

Suppose  $A(x', y') \sim B(x'', y'')$ . Then

$$y'' - y' = \sum_{i=1}^{n-1} (y_{i+1} - y_i) = \sum_{i'} (y_{i'+1} - y_{i'})$$

where  $1 \leq i' \leq n$  and  $y_{i'+1} - y_{i'} \neq 0$ . This inequality implies  $x_{i'} = x_{i'+1}$ . Then

$$y_{i'+1} - y_{i'} = y_{i'+1} - g(h(x_{i'+1})) - (y_{i'} - g(h(x_{i'})))$$

So  $y_{i'+1} - y_{i'}$  is the difference of two members of  $D$ . The set of all such differences is countable. The set of finite sums of these differences is countable too. Each difference  $y'' - y'$  is such a sum of this form. Therefore the set of all differences of ordinates of points belonging to an equivalence class  $G$  is countable. Thus  $\text{rng } G$  is countable. Let us enumerate members of  $\text{rng } G$  as  $\{y_m\}$  and all rational subintervals of  $I$  as  $\{J_n\}$ . Now enumerate all pairs  $(y_m, J_n)$  as  $\{(y_{m_i}, J_{n_i})\}$ . Since  $\text{dom } \Gamma^{y_{m_i}}$  is dense in  $H$ , there exists a point  $t_i \in J_{n_i} \cap \text{dom } \Gamma^{y_{m_i}}$ . Let  $t_1, t_2, \dots, t_k$  be chosen such that  $t_i \in J_{n_i} \cap \text{dom } \Gamma^{y_{m_i}}$  and  $t_i \neq t_j$  for  $1 \leq i < j \leq k$ . Take  $t_{k+1} \in J_{n_{k+1}} \cap \text{dom } \Gamma^{y_{m_{k+1}}}$ . Each set  $\text{dom } \Gamma^{y_{m_i}}$  contains a dense set of members of the sequence  $\{t_n\}$ . Set  $d(x) = y_{m_i}$  for  $x = t_i$ . Then  $\{d(x) : x = t_n \in J_{m_i}\} = \text{dom } G$ . Since  $\bigcup_{G \in \Gamma/\sim} \text{rng } G = R$  the function  $d$  defined on  $\bigcup_{G \in \Gamma/\sim} \{t_n\}$  maps each intersection  $H \cap J_m$  onto  $R$ . If  $G' \neq G''$  then  $\text{dom } G' \cap \text{dom } G'' = \emptyset$ . Therefore there is no contradiction in our extension of  $d$  from the subset of  $\text{dom } G$  to the sum  $\bigcup \text{dom } G$ .

At present  $d$  is defined on a subset of  $H$ . This subset is dense in  $I$ , therefore each next extension of  $d$  on  $I$  must be Darboux. Let us fix a point  $c \in D$ . Set  $d(x) = c - g(h(x))$  for each point  $x$  at which we have not defined  $d(x)$ . Since  $|H| = 0$ ,  $g(h(x)) + d(x) = c$  almost everywhere.

We still have to prove that  $d - (g \circ h)$  maps  $I$  onto  $D$ . Let  $G^0$  be the equivalence class containing  $\Gamma^0$ . Then  $D \in \text{rng } G^0$ . For each  $x = t_n$  the point  $(x, d(x))$  belongs to  $\Gamma$ . Thus  $d(x) + g(h(x)) \in D$ . For each  $x \neq t_n$  we have

$d(x) = c - g(h(x))$  where  $c \in D$  so that  $d(x) + g(h(x)) \in D$ . This completes the proof of Theorem.

## REFERENCES

1. A. M. Bruckner and J. Ceder, *On the sum of Darboux functions*, Proc. Amer. Math. Soc. **51** (1975), 97–102.
2. K. M. Garg, *Monotonicity, continuity and level sets of Darboux functions*, Colloq. Math. **28** (1973), 91–103. MR **48** #2317.
3. J. S. Lipiński, *On level sets of Darboux functions*, Fund. Math. **86** (1974), 193–199. MR **50** #10170.
4. A. N. Singh, *On some new types of non-differentiable functions*, Ann. of Math. (2) **28** (1927), 472–476.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF GDAŃSK, 80-952 GDAŃSK, POLAND