

## CONTRACTIONS OF CONVEX SETS

ROBERT E. JAMISON

**ABSTRACT.** In this paper it is shown that, in a vector space over any ordered field, a noninfinitesimal contraction of a convex set  $K$  can be written as an intersection of translates of  $K$ .

A subset  $K$  of a vector space over a totally ordered field is called *convex* provided  $\lambda x + (1 - \lambda)y$  is in  $K$  whenever  $x$  and  $y$  are in  $K$  and  $\lambda$  is a scalar such that  $0 \leq \lambda \leq 1$ . Recalling that any ordered field  $F$  has characteristic zero, and hence contains a copy of the rational numbers, we shall say that a positive element  $\mu$  in  $F$  is *infinitesimal* if  $\mu < r$  for all positive rational numbers  $r$ . (For further discussion and examples of ordered fields, see [1, Chapter 13].) In this note we shall prove the following intersection theorem:

**THEOREM.** *Suppose  $K$  is a convex set in a vector space over an ordered field and  $\mu$  is a positive scalar less than 1. If  $\mu$  is not infinitesimal, then, for some set  $T$  of vectors,*

$$\mu K = \bigcap \{K + t : t \in T\}.$$

It is easy to see that this result is plausible by considering either a square or a triangle in the plane, or in fact, any closed convex set. Difficulties arise, however, in the case of a convex set which includes only a portion of its boundary—say, the open unit disk together with the points of its circumference with rational  $x$ -coordinate.

**PROOF OF THE THEOREM.** Let  $\mu$  be a noninfinitesimal positive scalar less than 1. Set  $P = \mu K$  and suppose  $q$  is a point not in  $P$ . To prove the theorem, we must find a vector  $t$  such that  $P \subseteq K + t$  but  $q \notin K + t$ . We distinguish two cases:

*Case I.* For all  $x$  in  $P$ , we have  $q + \mu(x - q) \in P$ . Let  $t = (1 - \mu^{-1})q$  so that

$$K + t = \mu^{-1}P + (1 - \mu^{-1})q = q + \mu^{-1}(P - q).$$

The vectors in  $P - q$  are all nonzero since  $q$  is not in  $P$ . Thus  $q$  is not in  $K + t$ . But if  $x$  is in  $P$ , then  $q + \mu(x - q) \in P$  by the case hypothesis, so that  $x \in q + \mu^{-1}(P - q) = K + t$ . Thus  $P \subseteq K + t$ .

---

Received by the editors August 5, 1975 and, in revised form, March 24, 1976.

*AMS (MOS) subject classifications* (1970). Primary 52A05; Secondary 15A03, 12J15.

*Key words and phrases.* Contraction, convex set, intersection theorem, ordered field, translation, vector space.

© American Mathematical Society 1977

*Case II.* There is a point  $p$  in  $P$  such that  $q + \mu(p - q) \notin P$ . Note that if  $0 < \lambda < \mu$ , then the segment from  $q + \lambda(p - q)$  to  $q + (p - q)$  contains  $q + \mu(p - q)$ . Since  $q + (p - q) = p \in P$  and  $P$  is convex, the assumption on  $p$  forces  $q + \lambda(p - q) \notin P$ .

Now since  $\mu$  is not infinitesimal, the positive integral powers of  $1 - \mu$  become ultimately smaller than any preassigned positive rational number and, hence, smaller than any preassigned positive noninfinitesimal. Hence,  $(1 - \mu)^n < \mu$  for some sufficiently large positive integer  $n$ . By the preceding note,  $q + (1 - \mu)^n(p - q)$  cannot belong to  $P$ .

Let  $m$  be the smallest positive integer such that  $q + (1 - \mu)^m(p - q) \notin P$ , and set  $v = q + (1 - \mu)^{m-1}(p - q)$ . (Here  $v = p$  if  $m = 1$ .) Then  $v$  belongs to  $P$ . If  $x$  is any point in  $P$ , then  $(1 - \mu)v + \mu x \in P$  since  $P$  is convex. Whence

$$x \in \mu^{-1}(P - (1 - \mu)v) = K + (1 - \mu^{-1})v.$$

Thus letting  $t = (1 - \mu^{-1})v$ , we have  $P \subseteq K + t$ . But

$$\mu q + (1 - \mu)v = q + (1 - \mu)(v - q) = q + (1 - \mu)^m(p - q) \notin P.$$

Consequently,  $q$  does not belong to

$$\mu^{-1}(P - (1 - \mu)v) = K + (1 - \mu^{-1})v = K + t.$$

This completes the proof of the theorem.

We conclude with a simple one-dimensional example to show that the result cannot be extended to include infinitesimal contractions. Let  $F$  be an ordered field which contains a positive infinitesimal element  $\delta$  [1, p. 70]. Take the convex set  $K = \{x \in F: x \geq 0 \text{ and for some integer } n, x \leq n\}$ . If  $\lambda \in F$  such that  $\delta K \subseteq K - \lambda$ , then  $0 \in K - \lambda$  since  $0 \in \delta K$ . Thus  $\lambda \in K$ , so  $\lambda + 1$  must also be in  $K$ . Whence  $1 \in K - \lambda$ . Thus 1 belongs to every translate of  $K$  containing  $\delta K$ , but 1 does not belong to  $\delta K$  since  $\delta$  is infinitesimal.

**ACKNOWLEDGEMENT.** The author would like to acknowledge the referee's suggestions of refinements in the proof of the theorem.

#### BIBLIOGRAPHY

1. L. Gillman and M. Jerison, *Rings of continuous functions*, Van Nostrand, Princeton, N. J., 1960. MR 22 #6994.

DEPARTMENT OF MATHEMATICAL SCIENCES, CLEMSON UNIVERSITY, CLEMSON, SOUTH CAROLINA 29631

*Current address:* Mathematisches Institut, Universität Erlangen, 852 Erlangen, Federal Republic of Germany