

## A TWO-CARDINAL THEOREM AND A COMBINATORIAL THEOREM

SAHARON SHELAH<sup>1</sup>

ABSTRACT. We prove a new two-cardinal theorem, e.g.  $(\aleph_\omega, \aleph_0) \rightarrow (2^{\aleph_0}, \aleph_0)$ .  
 For this we prove a combinatorial theorem.

This is a sequel of Shelah [S1], and solves the main problem there. This problem also appears in Chang and Keisler [CK] and Friedman [Fr, Problem 30]. Our result is:

**THEOREM 1.** (A) *If for every  $n < \omega$ , the first order theory  $T$  has a model type  $(\aleph_{\alpha+n}, \aleph_\alpha)$  then whenever  $|T| \leq \mu < \lambda < \text{Ded}^* \mu$ ,  $T$  has a model type  $(\lambda, \mu)$ .*

(B) *If  $\aleph_{\alpha+\omega} < \text{Ded}^* \aleph_\alpha$  then  $(\aleph_{\alpha+\omega}, \aleph_\alpha)$  is  $\aleph_\alpha$ -compact and is complete.*

**REMARK.**  $\text{Ded}^* \mu$  is the first cardinal  $\chi$  such that no tree with  $\leq \mu$  nodes has  $\geq \chi$  branches of the same height. Note that  $\text{Ded}^* \aleph_0 = (2^{\aleph_0})^+$ , for every  $\lambda \lambda^+ < \text{Ded}^* \lambda \leq (2^\lambda)^+$ , and it is consistent with ZFC that  $\text{Ded}^* \aleph_1 \leq 2^{\aleph_1}$ .

This leads to many conjectures whose difficulty is not known to me; a sample is:

**CONJECTURE 2.** (A)  $(\aleph_{\alpha+\omega+\omega}, \aleph_{\alpha+\omega}, \aleph_\alpha) \rightarrow (\lambda, \mu, \chi)$  whenever  $\chi < \mu < \lambda < \text{Ded}^* \chi$ .

(B) *If a countable theory  $T$  has a  $\lambda$ -like model,  $\lambda$  a limit cardinal, and  $|T| \leq \mu < \lambda_1 < \text{Ded}^* \mu$ ,  $\lambda_1$  a singular cardinal then  $T$  has a  $\lambda_1$ -like model. If  $\lambda$  is  $M_\omega$ -Mahlo weakly inaccessible cardinal, we can remove the singularity of  $\lambda_1$ .*

(C) *If  $\psi \in L_{\omega_1, \omega}$  has a model of cardinality  $\aleph_{\omega_1}$ , then  $\psi$  has a model of cardinality  $2^{\aleph_0}$ .*

**NOTATION.** Let  $I$  denote a well-ordered set. A  $(\lambda, n)$ -box  $B$  is  $\prod_{l < n} I_l$  where  $I_l$  has order type  $\lambda$ ;  $\lambda, \mu, \chi$  denote infinite cardinals, elements of boxes will be denoted by  $\eta, \tau, \nu$ , and  $\eta = \langle \eta(0), \dots, \eta(n-1) \rangle$ . For a  $(\lambda, n)$ -box  $B$ , and  $\eta_l \in B$  ( $l < n$ ) we say  $\langle \eta_0, \dots, \eta_{n-1} \rangle$  is proper for  $B$  if  $k \neq l < n \Rightarrow \eta_k(l) < \eta_l(l)$ .

Let  $\lambda^{+0} = \lambda, \lambda^{+(k+1)} = (\lambda^{+k})^+ =$  the successor of  $\lambda^{+k}$ .

A  $B$ -indexed set is  $\{a_\eta : \eta \in B\}$  such that  $\eta \neq \tau \rightarrow a_\eta \neq a_\tau$ . Under those conditions  $\langle a_{\eta_0}, \dots \rangle$  is proper, iff  $\langle \eta_0, \dots \rangle$  is proper. A  $(\lambda, n)$ -indexed set is a  $B$ -indexed set for some  $(\lambda, n)$ -box  $B$ .

**LEMMA 3.** *Suppose that  $f_\alpha : (\lambda^+)^2 \rightarrow \lambda$  for each  $\alpha < \lambda$ . Then there exist*

Received by the editors November 27, 1974 and, in revised form, May 1, 1975.

AMS (MOS) subject classifications (1970). Primary 02H05, 04A20.

Key words and phrases. Two-cardinal theorem, partition calculus.

<sup>1</sup> I would like to thank the referee for detecting many errors, and rewriting Lemma 3—Theorem 5.

$s, t < \lambda^+$  such that for each  $\alpha < \lambda$  there exist  $a, b, c, d$  for which  $s < a < b < \lambda^+, t < c < d < \lambda^+$  and  $f_\alpha(s, t) = f_\alpha(a, d) = f_\alpha(b, c)$ .

**PROOF.** For every  $u < \lambda^+$  let  $t_u < \lambda^+$  be such that whenever  $\alpha < \lambda$  and  $t \geq t_u$ , then

$$|\{v < \lambda^+ : f_\alpha(u, t) = f_\alpha(u, v)\}| = \lambda^+.$$

Let  $X_u = \{(\alpha, \beta) : f_\alpha(u, t) = \beta \text{ for some } t \geq t_u\}$ . Now let  $s < \lambda^+$  be such that whenever  $u \geq s$  and  $(\alpha, \beta) \in X_u$ , then

$$|\{v < \lambda^+ : (\alpha, \beta) \in X_v\}| = \lambda^+.$$

Let  $t = t_s$ . It is clear that this  $s$  and  $t$  work.

**LEMMA 4.** Let  $f: A^n \rightarrow \lambda$  where  $A$  is any  $(\lambda^+, n + 1)$ -indexed set and  $k < n$ . Then there is a  $(\lambda, n)$ -indexed set  $A^* \subset A$  such that:

(\*) For any proper sequence  $\langle a_0, \dots, a_{n-1} \rangle$  from  $A^*$  there is a proper sequence  $\langle b_0, \dots, b_n \rangle$  from  $A$  such that

$$f(a_0, \dots, a_{n-1}) = f(b_0, \dots, b_{n-1}) = f(b_0, \dots, b_{k-1}, b_n, b_{k+1}, \dots, b_{n-1}).$$

**PROOF.** Let  $A$  be a  $B$ -indexed set where  $B = \prod_{l < n+1} I_l$  and  $A = \{a_\eta : \eta \in B\}$ . For notational simplicity let  $k = n - 1$  and each  $I_l = \lambda^+$ . Now we define  $\langle s_\alpha : \alpha < \lambda \rangle$  and  $\langle t_\alpha : \alpha < \lambda \rangle$  by induction on  $\alpha$  such that:

- (i)  $s_\alpha, t_\alpha < \lambda^+$ .
- (ii)  $\langle s_\alpha : \alpha < \lambda \rangle$  and  $\langle t_\alpha : \alpha < \lambda \rangle$  are increasing.
- (iii) Whenever  $\eta_0, \dots, \eta_{n-2} \in B$  and  $\tau \in \lambda^{n-1}$  are such that for each  $i, l < n - 1$  there is  $\beta < \alpha$  such that  $\eta_i(l) < \lambda, \eta_i(n - 1) = s_\beta$  and  $\eta_i(n) = t_\beta$ , then there are  $a, b, c, d$  such that  $s_\alpha < a < b < \lambda^+, t_\alpha < c < d < \lambda^+$  and

$$\begin{aligned} f(a_{\eta_0}, \dots, a_{\eta_{n-2}}, a_{\tau \wedge \langle s_\alpha, t_\alpha \rangle}) &= f(a_{\eta_0}, \dots, a_{\eta_{n-2}}, a_{\tau \wedge \langle a, d \rangle}) \\ &= f(a_{\eta_0}, \dots, a_{\eta_{n-2}}, a_{\tau \wedge \langle b, c \rangle}). \end{aligned}$$

Suppose we have defined  $s_\beta$  and  $t_\beta$  for  $\beta < \alpha$ . For each  $\eta_0, \dots, \eta_{n-2}, \tau$  satisfying the conditions of (iii), there is a function  $g: (\lambda^+)^2 \rightarrow \lambda$  defined by

$$g(x, y) = f(a_{\eta_0}, \dots, a_{\eta_{n-2}}, a_{\tau \wedge \langle x, y \rangle}).$$

There are  $\leq \lambda$  such functions  $g$ . So we can apply Lemma 3 to get  $s_\alpha, t_\alpha$  such that  $s_\alpha > t_\beta$  and  $t_\alpha > t_\beta$  for each  $\beta < \alpha$ , and for each such  $g$  there are  $a, b, c, d$  such that  $s_\alpha < a < b < \lambda^+, t_\alpha < c < d < \lambda^+$  and  $g(s_\alpha, t_\alpha) = g(a, d) = g(b, c)$ .

Now we define the  $(\lambda, n)$ -indexed set  $A^*$ . For each  $\tau \in \lambda^{n-1}$  let  $b_{\tau \wedge \langle \alpha \rangle} = a_{\tau \wedge \langle s_\alpha, t_\alpha \rangle}$ . Then let  $A^* = \{b_\eta : \eta \in \lambda^n\}$ . Now it is easy to check that (\*) holds.

**THEOREM 5.** Let  $f_l: (\lambda^{+n})^l \rightarrow \lambda$  whenever  $0 < l \leq n$ , and let  $h: (n + 1) \rightarrow n$  be such that  $h(l) < l$  whenever  $0 < l \leq n$ . Then there are distinct  $a_0, \dots, a_n < \lambda^{+n}$  such that

$$f_l(a_0, \dots, a_{l-1}) = f_l(a_0, \dots, a_{h(l)-1}, a_l, a_{h(l)+1}, \dots, a_{l-1})$$

whenever  $0 < l \leq n$ .

**PROOF.** We let  $\lambda^{+n}$  be  $(\lambda^{+n}, n + 1)$ -indexed. Now we prove the theorem by induction on  $n$ . For  $n = 0$  there is nothing to prove. For  $n + 1$  we use Lemma 4, and the induction hypothesis on  $A^*$ .

**PROOF OF THEOREM 1.** Clear from [S1, §3] and Lemma 4.

**REMARKS.** (1) The following theorem is clear.

**THEOREM .** *If every finite subset of  $T$  has, for each  $n$ , a model of type  $(\lambda_m, \dots, \lambda_0)$  where  $\lambda_0^{+n} < \lambda_1, [(\lambda_{l+1})^{(\lambda_l^{+n})}]^{+n} < \lambda_{l+2}$  ( $l < n - 1$ ) and  $|T| \leq \mu_0 \leq \mu_1 \leq \dots \leq \mu_m < \text{Ded}^* \mu_0$  then  $T$  has a model of type  $(\mu_m, \dots, \mu_0)$ . (The parallel theorem in [S1] was noted by Papageorgiou.)*

(2) We can prove the main theorem of [S1] in a way similar to the proof here.

(3) In the notation of Shelah [S2, §3], we have proved in Lemma 4 that for  $m = 2^n, r < \omega, \lambda^{+m} \xrightarrow{w} (n)_\lambda^r$ . This answers positively question 3 from [S2]. But it is still unknown whether we have the best results.

(4) Halperin and Levi [H Le] used an indiscernibility similar to the one used in [S1], and Halperin and Lauchli proved the necessary combinatorial theorem. We have not succeeded in generalizing their proof. However, we can use our method to prove a weaker variant of their theorem, which is sufficient to prove that if  $T \subset T_1, |T_1| = \aleph_0, T$  is complete and there are  $> \aleph_0$  complete  $L(T)$ -types consistent with  $T, \lambda > \aleph_0$  then  $T$  has  $\geq \min\{2^\lambda, 2^{2^{\aleph_0}}\}$  nonisomorphic  $L(T)$ -reducts of models of  $T_1$ . (This will appear in [S3].)

(5) To see the connection note that by [S3] it follows by Theorem 5 that

**THEOREM .** *Let  $f_j: (\lambda^{+n})^l \rightarrow \lambda$  whenever  $0 < l \leq n, n = 2^m - 1$ . Then there are distinct  $a_\eta < \lambda^{+n}$  for  $\eta \in 2^m$  (i.e.  $\eta$  is a sequence of ones and zeros of length  $m$ ) such that: if  $k < m, 0 < l \leq n, \tau_1, \dots, \tau_l$  are distinct members of  $2^k$ , and for  $1 \leq i \leq l, 0 \leq j \leq 1, \eta_i^j \in 2^m$  and  $\tau_i$  is an initial segment of  $\eta_i^j$  then  $f_l(\eta_1^0, \dots, \eta_l^0) = f_l(\eta_1^1, \dots, \eta_l^1)$ .*

(6) In Lemma 4 and Theorem 5 we can consider  $\mu$  such functions, provided that  $\mu \leq x, \lambda^\mu = \lambda$ , resp.

(7) We can use only a  $(B, n)$ -indexed set for a fixed  $n$  in (4), but then in Remark (3)  $m$  will become bigger.

REFERENCES

[CK] C. C. Chang and H. J. Keisler, *Model theory*, North-Holland, Amsterdam, 1973.  
 [H Le] J. D. Halpern and A. Lévy, *The Boolean prime ideal theorem does not apply the axiom of choice*, Proc. Sympos. Pure Math., vol. 13, part 1, Amer. Math. Soc., Providence, R.I., 1971, pp. 83-134. MR 44 #1557.  
 [H La] J. D. Halpern and H. Lauchli, *A partition theorem*, Trans. Amer. Math. Soc. 124 1966, 360-367. MR 34 #71.  
 [S1] S. Shelah, *A two-cardinal theorem*, Proc. Amer. Math. Soc. 48 (1975), 207-213.  
 [S2] ———, *Coloring without triangles and partition relation*, Israel Math. 20 (1975), 1-12.  
 [S3] ———, *Stability and number of non-isomorphic models* (in preparation).

INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY, JERUSALEM, ISRAEL