

AN EXAMPLE OF A WEIGHT WITH TYPE III CENTRALIZER

UFFE HAAGERUP

ABSTRACT. We give an example of a weight ϕ on the hyperfinite type II_∞ factor R , such that the centralizer M_ϕ of ϕ is a type III von Neumann algebra.

Introduction. Let ϕ be a normal, semifinite, faithful weight on a von Neumann algebra M , and σ_t^ϕ the modular automorphism group associated with ϕ . The centralizer M_ϕ is the fixed point-algebra under σ_t^ϕ , i.e.,

$$M_\phi = \{x \in M \mid \sigma_t^\phi(x) = x, t \in R\}.$$

The restriction of ϕ to M_ϕ is a trace. Hence if ϕ is *strictly semifinite* (i.e. the restriction of ϕ to M_ϕ is semifinite), then M_ϕ is a semifinite von Neumann algebra (cf. [1]). In particular, if ϕ is bounded, then M_ϕ is finite. Recently A. Connes and M. Takesaki have proved that M_ϕ is semifinite for a great class of non strictly semifinite weights, namely the *integrable* weights, i.e. those weights ϕ , for which the set $\{x \in M_+ \mid \int_{-\infty}^\infty \sigma_t^\phi(x) dt \in M_+\}$ is σ -weakly dense in M_+ (cf. [3]).

In this paper we prove that M_ϕ is not, in general, semifinite. At the same time we find that the relative commutant $N' \cap M$ of a semifinite von Neumann algebra N in a semifinite von Neumann algebra M is not, in general, semifinite (Lemma 2). That is the negative solution to a problem considered in [2].

LEMMA 1. *On a separable Hilbert space H , there exists a hyperfinite type II_1 factor M and an abelian von Neumann algebra $A \subseteq M'$, such that the von Neumann algebra $(M \cup A)''$ is of type III.*

PROOF. Let $B = \bigotimes_{n=1}^\infty F_n$ be the uniformly hyperfinite C^* -algebra obtained as tensor product of a sequence of type I_2 -factors (cf. [5, Chapter 1, §23]). B has a unique normalized trace τ , and $\pi_\tau(B)''$ is the hyperfinite type II_1 -factor. Let U_n be the unitary group in F_n , and put $G = \prod_{n=1}^\infty U_n$. Then G is a compact group with normalized Haar measure dg . We define an action of G as $*$ -automorphisms on B in the following way: For $g \in G$, $g = (u_n)_{n \in \mathbb{N}}$, we put

$$\sigma_g = \bigotimes_{n=1}^\infty \text{ad}(u_n).$$

Received by the editors February 2, 1976.

AMS (MOS) subject classifications (1970). Primary 46L10.

Key words and phrases. von Neumann algebras, Tomita-Takesaki theory, weights.

© American Mathematical Society 1977

It is easy to check that $g \rightarrow \sigma_g$ is a homeomorphism, and that $(g, x) \rightarrow \sigma_g x$ is jointly continuous. Let ϕ be a state on B , and put

$$\omega(x) = \int_G \phi \cdot \sigma_g(x) dg.$$

Let ω_n (resp. τ_n) be the restriction of ω (resp. τ) to $F_1 \otimes F_2 \otimes \dots \otimes F_n$. By definition ω_n is invariant under $\text{ad}(u_1) \otimes \dots \otimes \text{ad}(u_n)$, $u_k \in U_k$, $k = 1, \dots, n$. From this it follows that $\omega_n = \tau_n$ for any $n \in N$, and thus $\omega = \tau$. Hence for any state ϕ on B ,

$$\tau(x) = \int_G \phi \cdot \sigma_g(x) dg.$$

In the following we let ϕ be a fixed type III state on B . On the C^* -tensor product $B \otimes C(G) = C(G, B)$ we regard the state

$$\tilde{\phi}(x(g)) = \int_G \phi \cdot \sigma_g(x(g)) dg, \quad x \in B \otimes C(G).$$

$\tilde{\phi}$ is a type III state, because

$$\tilde{\phi}(x) = (\phi \otimes dg)\tilde{\sigma}(x), \quad x \in B \otimes C(G),$$

where $\tilde{\sigma}$ is the $*$ -automorphism of $B \otimes C(G)$ given by $\tilde{\sigma}(x)(g) = \sigma_g x(g)$. Hence $N = \pi_{\tilde{\phi}}(B \otimes C(G))''$ is of type III. However, N is generated by $\pi_{\tilde{\phi}}(B \otimes I)''$ and $\pi_{\tilde{\phi}}(I \otimes C(G))''$. The first is isomorphic to the hyperfinite type II_1 -factor $\pi_{\tau}(B)''$ because $\tilde{\phi}(x \otimes I) = \tau(x)$, $x \in B$, and the latter is commutative. Moreover the two algebras commute. Since $B \otimes C(G)$ is a separable C^* -algebra, the Hilbert space $H_{\tilde{\phi}}$ associated with $\pi_{\tilde{\phi}}$ is separable. This completes the proof.

LEMMA 2. *Let R be the hyperfinite factor of type II_{∞} . There exists an abelian sub von Neumann algebra $A \subseteq R$, such that the relative commutant $R \cap A'$ is of type III.*

PROOF. Let M and A be as in Lemma 1. Put $R = M'$. Then $R \cap A' = (M \cup A)'$ is of type III. Since R contains a type III von Neumann algebra, it is of infinite multiplicity. Hence $R \simeq M \otimes F_{\infty}$ where F_{∞} is a type I_{∞} factor.

THEOREM. *There exists a normal, semifinite faithful weight on the hyperfinite factor of type II_{∞} , such that the centralizer M_{ϕ} is of type III.*

PROOF. Let A and R be as in Lemma 2. Since A is an abelian von Neumann algebra, which can be represented faithfully on a separable Hilbert space, A is the von Neumann algebra generated by a single selfadjoint element h . We may assume that $h \geq 1$. Put $\phi = \tau(h \cdot)$ where τ is the trace on R . Then

$$\sigma_t^{\phi}(x) = h^{it} x h^{-it}, \quad x \in R \quad (\text{cf. [4]}).$$

Hence $M_{\phi} = R \cap \{h\}' = R \cap A'$ which is of type III.

REFERENCES

1. F. Combes, *Poids et espérances conditionnelles dans les algèbres de von Neumann*, Bull. Soc. Math. France **99** (1971), 73–112. MR **44** #5785.
2. F. Combes and C. Delaroche, *Groupe modulaire d'une espérance conditionnelle dans une algèbre de von Neumann*, Université d'Orléans, 1974 (preprint).
3. A. Connes and M. Takesake, *Flow of weights on factors of type III*, 1975 (preprint).
4. G. K. Pedersen and M. Takesaki, *The Radon-Nikoȳm theorem for von Neumann algebras*, Acta Math. **130** (1973), 53–87.
5. Shōichirō Sakai, *C*-algebras and W*-algebras*, Springer-Verlag, Berlin, 1971.

ODENSE UNIVERSITET, ODENSE, DENMARK