

## MULTIPLIERS ON DUAL $A^*$ -ALGEBRAS

B. J. TOMIUK

**ABSTRACT.** Let  $A$  be an  $A^*$ -algebra which is a dense  $*$ -ideal of a  $B^*$ -algebra  $\mathfrak{A}$ . We use tensor products and the algebra  $M_l(A)$  of left multipliers on  $A$  to obtain a characterization of duality in  $A$ . We show, moreover, that if  $A$  is dual then  $M_l(A)$  is algebra isomorphic to the second conjugate space  $\mathfrak{A}^{**}$  of  $\mathfrak{A}$  when  $\mathfrak{A}^{**}$  is given Arens product.

**1. Introduction.** Let  $A$  be an  $A^*$ -algebra which is a dense  $*$ -ideal of a  $B^*$ -algebra  $A$ . In [8] a necessary and sufficient condition was given for  $A$  to be dual which was expressed in terms of the weak operator topology on  $M_r(A)$ , the algebra of right multipliers on  $A$ , and a certain property of  $A$  called property (P2). In this paper we give several characterizations of property (P2) and then use some of them to give conditions for duality in  $A$ . Our presentation here differs somewhat from that in [8]. We use the tensor product approach as developed in [3] and [6]. Particularly in §2 we follow closely the presentation given in [3].

We shall use the notation of [8]. An  $A^*$ -algebra  $A$  is said to be of the first kind if it is an ideal of its completion  $\mathfrak{A}$  in the auxiliary norm  $|\cdot|$ . It follows that there exists a constant  $k > 0$  such that  $\|xy\| \leq k\|x\| \|y\|$  for all  $x \in A, y \in \mathfrak{A}$  [4, Lemma 4, p. 18]. If  $A$  is a modular annihilator  $A^*$ -algebra then  $|\cdot|$  is unique [1, (1.3), p. 6] so that  $\mathfrak{A}$  is also unique.

**2. The property (P2).** Let  $A$  be a Banach algebra,  $A^*$  and  $A^{**}$  its first and second conjugate spaces. Let  $A \hat{\otimes} A^*$  be the projective tensor product of  $A$  and  $A^*$  [7, pp. 92–95]. Then  $A \hat{\otimes} A^*$  is a Banach space with elements of the form  $\sum_{k=1}^{\infty} a_k \hat{\otimes} f_k$  such that  $\sum_{k=1}^{\infty} \|a_k\| \|f_k\| < \infty, a_k \in A, f_k \in A^*$ , and the norm given by

$$\|h\| = \inf \left\{ \sum_{k=1}^{\infty} \|a_k\| \|f_k\| : h = \sum_{k=1}^{\infty} a_k \hat{\otimes} f_k \right\}.$$

For  $a \in A, f \in A^*$ , let  $af \in A^*$  be given by  $(af)x = f(xa), x \in A$ . This makes  $A^*$  into a left Banach  $A$ -module. We note that if  $\sum_{k=1}^{\infty} a_k \hat{\otimes} f_k \in A \hat{\otimes} A^*$ , then  $\sum_{k=1}^{\infty} a_k f_k \in A^*$ . Let  $\psi$  be the continuous linear map of  $A \hat{\otimes} A^*$  into  $A^*$  given by

---

Received by the editors June 17, 1976.

AMS (MOS) subject classifications (1970). Primary 46K10, 46L20; Secondary 47B05.

Key words and phrases. Dual  $A^*$ -algebra, multipliers, Arens product, projective tensor product, Banach  $A$ -module.

© American Mathematical Society 1977

$$\psi(a \hat{\otimes} f) = af \quad (a \in A, f \in A^*).$$

For  $a \in A, F \in A^{**}$ , let  $Fa \in A^{**}$  be given by  $(Fa)f = F(af), f \in A^*$ . Then  $A^{**}$  is a right Banach  $A$ -module. (See [6] for the definition and properties of Banach  $A$ -modules.)

Let  $A \circ A^*$  be the Banach space  $A \hat{\otimes} A^*/\ker(\psi)$  with the usual quotient norm, where  $\ker(\psi)$  is the kernel of  $\psi$ . Then  $(A \circ A^*)^*$  consists of all those  $\mathcal{F} \in (A \hat{\otimes} A^*)^*$  which vanish on  $\ker(\psi)$ . Now, for each  $F \in A^{**}$ , let  $\mathcal{F}_F \in (A \hat{\otimes} A^*)^*$  be given by

$$\mathcal{F}_F(a \hat{\otimes} f) = F(\psi(a \hat{\otimes} f)) = F(af) \quad (a \in A, f \in A^*).$$

Each  $\mathcal{F}_F$  vanishes on  $\ker(\psi)$ , so that  $\{\mathcal{F}_F: F \in A^{**}\}$  may be identified as a subspace of  $(A \circ A^*)^*$ . Moreover if  $Fa = 0$  for all  $a \in A$  implies  $F = 0$ , then  $F \rightarrow \mathcal{F}_F$  is a one-to-one map of  $A^{**}$  into  $(A \hat{\otimes} A^*)^*$ . Thus in this case  $F \rightarrow \mathcal{F}_F$  identifies  $A^{**}$  as a subspace of  $(A \circ A^*)^*$ .

Let  $\sigma$  denote the  $w^*$ -topology of  $(A \hat{\otimes} A^*)^*$ .

LEMMA 2.1.  $(A \circ A^*)^*$  is the  $\sigma$ -closure of  $\{\mathcal{F}_F: F \in A^{**}\}$ .

PROOF. We have

$$\begin{aligned} \ker(\psi) &= \left\{ \sum_{k=1}^{\infty} a_k \hat{\otimes} f_k \in A \hat{\otimes} A^*: \sum_{k=1}^{\infty} a_k f_k = 0 \right\} \\ &= \left\{ \sum_{k=1}^{\infty} a_k \hat{\otimes} f_k \in A \hat{\otimes} A^*: F\left(\sum_{k=1}^{\infty} a_k f_k\right) = 0, F \in A^{**} \right\}. \end{aligned}$$

Thus  $\ker(\psi) = \cap [\ker(\mathcal{F}_F): F \in A^{**}]$ , which means that  $\ker(\psi)$  is the polar of  $\{\mathcal{F}_F: F \in A^{**}\}$ . Therefore, by the Bipolar Theorem [7, p. 126],  $(A \circ A^*)^*$  is the  $\sigma$ -closure of  $\{\mathcal{F}_F: F \in A^{**}\}$ . This completes the proof.

We observe that if  $A^2 = (0)$ , then  $\ker(\mathcal{F}_F) = A \hat{\otimes} A^*$ , for every  $F \in A^{**}$ , so that  $\ker(\psi) = A \hat{\otimes} A^*$  and consequently  $(A \circ A^*)^* = (0)$ .

Let  $\mathfrak{B}(A, A^{**})$  be the Banach space of all bounded linear operators  $T: A \rightarrow A^{**}$  normed with the operator bound norm. For each  $\mathcal{F} \in (A \hat{\otimes} A^*)^*$ , let  $T_{\mathcal{F}}$  be the map on  $A$  into  $A^{**}$  given by

$$(f, T_{\mathcal{F}}(a)) = \mathcal{F}(a \hat{\otimes} f) \quad (a \in A, f \in A^*).$$

Then clearly  $T_{\mathcal{F}} \in \mathfrak{B}(A, A^{**})$  for every  $\mathcal{F} \in (A \hat{\otimes} A^*)^*$ , and it is easy to check that the map  $\phi: \mathcal{F} \rightarrow T_{\mathcal{F}}$  is an isometric isomorphism of  $(A \hat{\otimes} A^*)^*$  onto  $\mathfrak{B}(A, A^{**})$ . Give  $\mathfrak{B}(A, A^{**})$  the image of the  $\sigma$  topology by the map  $\phi$ . Following Máté [3], we shall refer to this topology as the ultraweak topology on  $\mathfrak{B}(A, A^{**})$ .

Now consider  $A^{**}$  as a right Banach  $A$ -module and let  $\text{Hom}_A(A, A^{**})$  be the set of all  $T \in \mathfrak{B}(A, A^{**})$  such that  $T(ab) = T(a)b, a, b \in A$ . The canonical map  $\pi: A \rightarrow A^{**}$  belongs to  $\mathfrak{B}(A, A^{**})$  since  $\pi(ab)f = f(ab) = \pi(a)(bf)$  for all  $a, b \in A$  and  $f \in A^*$ . For each  $F \in A^{**}$ , let  $T_F: A \rightarrow A^{**}$  be given by  $T_F(a) = Fa, a \in A$ . Then  $T_F \in \text{Hom}_A(A, A^{**})$ , and we have  $\phi(\mathcal{F}_F)$

$= T_F$  for all  $F \in A^{**}$ . In view of Lemma 2.1 and the fact that the ultraweak closure of  $\{T_F: F \in A^{**}\} \subseteq \text{Hom}_A(A, A^{**})$  we have

LEMMA 2.2.  $\phi((A \circ A^*)^*) \subseteq \text{Hom}_A(A, A^{**})$  and is the ultraweak closure of  $\{T_F: F \in A^{**}\}$ .

If  $\phi$  maps  $(A \circ A^*)^*$  onto  $\text{Hom}_A(A, A^{**})$ , we shall write  $(A \circ A^*)^* \cong \text{Hom}_A(A, A^{**})$ . In this case, for every  $T \in \text{Hom}_A(A, A^{**})$ ,  $\mathfrak{F}_T \in (A \hat{\otimes} A^*)^*$ , given by  $\mathfrak{F}_T(a \hat{\otimes} f) = (f, T(a))$ , belongs to  $(A \circ A^*)^*$ . In particular,

$$\mathfrak{F}_\pi(a \hat{\otimes} f) = f(a), \text{ for all } a \in A, f \in A^*.$$

We recall that a Banach algebra  $A$  is said to have property (P2) if:  $a_k \in A, f_k \in A^*, \sum_{k=1}^\infty \|a_k\| \|f_k\| < \infty$  and  $\sum_{k=1}^\infty a_k f_k = 0$  implies that  $\sum_{k=1}^\infty f_k(a_k) = 0$ . (This is the left-hand version of the definition given in [8].)

THEOREM 2.3. Let  $A$  be a Banach algebra. Then the following statements are equivalent:

- (i)  $A$  has property (P2).
- (ii) For  $h = \sum_{k=1}^\infty a_k \hat{\otimes} f_k \in \ker(\psi)$  we have  $\sum_{k=1}^\infty f_k(a_k) = 0$ .
- (iii)  $\mathfrak{F}_\pi$  vanishes on  $\ker(\psi)$ .
- (iv)  $\mathfrak{F}_\pi \in (A \circ A^*)^*$ .
- (v)  $\text{Hom}_A(A, A^{**}) \cong (A \circ A^*)^*$ .
- (vi) There exists a net  $\{u_\alpha\}$  in  $A$  such that  $\{\mathfrak{F}_{\pi(u_\alpha)}\}$  converges to  $\mathfrak{F}_\pi$  in the  $w^*$ -topology on  $(A \hat{\otimes} A^*)^*$ .

PROOF. (i)  $\Leftrightarrow$  (ii) and (iii)  $\Leftrightarrow$  (iv) are clear.

(iv)  $\Rightarrow$  (vi). Suppose (iv) holds. Then  $\ker(\mathfrak{F}_\pi) \supset \ker(\psi)$ . We have

$$\mathfrak{F}_\pi(h) = \mathfrak{F}_\pi\left(\sum_{k=1}^\infty a_k \hat{\otimes} f_k\right) = \sum_{k=1}^\infty \mathfrak{F}_\pi(a_k \hat{\otimes} f_k) = \sum_{k=1}^\infty f_k(a_k),$$

for all  $h = \sum_{k=1}^\infty a_k \hat{\otimes} f_k \in A \hat{\otimes} A^*$ . Since  $\{\mathfrak{F}_F: F \in A^{**}\}$  is  $\sigma$ -dense in  $(A \circ A^*)^*$ , there exists a net  $\{F_\alpha\}$  in  $A^{**}$  such that  $\mathfrak{F}_{F_\alpha}(h) \rightarrow \mathfrak{F}_\pi(h)$  for all  $h \in A \hat{\otimes} A^*$ . Since  $\pi(A)$  is  $w^*$ -dense in  $A^{**}$  and  $\sigma$  is weaker than the  $w^*$ -topology on  $A^{**}$ , it follows that  $\{\mathfrak{F}_{\pi(a)}: a \in A\}$  is  $\sigma$ -dense in  $(A \circ A^*)^*$ . Hence there exists a net  $\{u_\alpha\}$  in  $A$  such that  $\{\mathfrak{F}_{\pi(u_\alpha)}\}$   $\sigma$ -converges to  $\mathfrak{F}_\pi$ . We have

$$\begin{aligned} \mathfrak{F}_{\pi(u_\alpha)}(h) &= \pi(u_\alpha)(\psi(h)) = \sum_{k=1}^\infty \pi(u_\alpha)(a_k f_k) \\ &= \left(\sum_{k=1}^\infty a_k f_k\right)(u_\alpha) = \sum_{k=1}^\infty (a_k f_k)(u_\alpha) \\ &= \sum_{k=1}^\infty f_k(u_\alpha a_k). \end{aligned}$$

Thus

$$(1) \quad \lim_{\alpha} \mathfrak{F}_{\pi(u_{\alpha})}(h) = \lim_{\alpha} \sum_{k=1}^{\infty} f_k(u_{\alpha} a_k) = \sum_{k=1}^{\infty} f_k(a_k) = \mathfrak{F}_{\pi}(h),$$

for all  $h = \sum_{k=1}^{\infty} a_k \hat{\otimes} f_k \in A \hat{\otimes} A^*$ .

(vi)  $\Rightarrow$  (v). We have  $\phi((A \circ A^*)^*) \subseteq \text{Hom}_A(A, A^{**})$ . We need only show that  $\text{Hom}_A(A, A^{**}) \subseteq \phi((A \circ A^*)^*)$ . Let  $T \in \text{Hom}_A(A, A^{**})$  and let  $\mathfrak{F}_T$  be the corresponding element of  $(A \hat{\otimes} A^*)^*$ . Then, using (1), we obtain (identifying  $A$  as a subset of  $A^{**}$  and  $A^*$  as a subset of  $A^{***}$ ):

$$\begin{aligned} \mathfrak{F}_T \left( \sum_{k=1}^{\infty} a_k \hat{\otimes} f_k \right) &= \sum_{k=1}^{\infty} \mathfrak{F}_T(a_k \hat{\otimes} f_k) = \sum_{k=1}^{\infty} (f_k, T(a_k)) \\ &= \sum_{k=1}^{\infty} (T^* f_k)(a_k) = \lim_{\alpha} \sum_{k=1}^{\infty} (T^* f_k)(u_{\alpha} a_k) \\ &= \lim_{\alpha} \sum_{k=1}^{\infty} f_k(T(u_{\alpha} a_k)) = \lim_{\alpha} \sum_{k=1}^{\infty} f_k(T(u_{\alpha}) a_k) \\ &= \lim_{\alpha} \sum_{k=1}^{\infty} (T^* a_k f_k)(u_{\alpha}) = \sum_{k=1}^{\infty} T^*(a_k f_k), \end{aligned}$$

where  $T^*$  is the conjugate of  $T$ . Hence if  $\sum_{k=1}^{\infty} a_k f_k = 0$  then  $\mathfrak{F}_T(\sum_{k=1}^{\infty} a_k \hat{\otimes} f_k) = 0$ , so that  $\ker(\psi) \subset \ker(\mathfrak{F}_T)$ . Thus  $\text{Hom}_A(A, A^{**}) \subseteq \phi((A \circ A^*)^*)$  and so  $\text{Hom}_A(A, A^{**}) \cong (A \circ A^*)^*$ .

(v)  $\Rightarrow$  (iv). This is clear since  $\pi \in \text{Hom}_A(A, A^{**})$ .

**3. Dual  $A^*$ -algebras.** Let  $A$  be a Banach algebra. A map  $T: A \rightarrow A$  is called a left (resp. right) multiplier if  $T(ab) = T(a)b$  (resp.  $T(ab) = aT(b)$ ), for all  $a, b \in A$ . Let  $M_l(A)$  (resp.  $M_r(A)$ ) be the set of all bounded linear left (resp. right) multipliers on  $A$ .  $M_l(A)$  and  $M_r(A)$  are Banach algebras under the usual operations for operators and the operator bound norm. We observe that if  $T \in M_l(A)$  then the composite map  $\pi \circ T \in \text{Hom}_A(A, A^{**})$ . Let  $\phi_{\pi}$  be the map of  $M_l(A)$  into  $\text{Hom}_A(A, A^{**})$  given by

$$\phi_{\pi}(T) = \pi \circ T \quad (T \in M_l(A)).$$

For any Banach space  $X$ , let  $\mathfrak{S}(X)$  denote the closed unit ball of  $X$ . It follows from the proof of [8, Theorem 4.7, p. 286] that if  $A$  is a dual  $A^*$ -algebra of the first kind then  $\mathfrak{S}(M_l(A))$  is  $\tau_l$ -compact, where  $\tau_l$  is the weak operator topology on  $M_l(A)$ . (We take the left-hand version of the arguments in [8, p. 286].)

**THEOREM 3.1.** *Let  $A$  be an  $A^*$ -algebra of the first kind. Then the following statements are equivalent:*

- (i)  $A$  is dual.
- (ii)  $\phi_{\pi}(M_l(A))$  is the ultraweak closure of  $\{T_F: F \in A^{**}\}$ .

**PROOF.** (i)  $\Rightarrow$  (ii). Suppose  $A$  is dual. Then, by [8, Theorem 4.7, p. 286], it has property (P2) and therefore, by Theorem 2.3,  $(A \circ A^*)^* \cong \text{Hom}_A(A, A^{**})$ . Hence  $\text{Hom}_A(A, A^{**})$  is the ultraweak closure of  $\{T_F: F \in$

$A^{**}$ ). Now  $T_F(a)f = (Fa)f = (F * \pi(a))f$  and, by [9, Theorem 5.2, p. 830],  $\pi(A)$  is an ideal of  $A^{**}$  when  $A^{**}$  is given either Arens product, so that  $T_F(a) \in \pi(A)$  for all  $a \in A$ . Therefore  $T_F = \pi \circ T$ , for some  $T \in M_l(A)$ , and so  $\{T_F: F \in A^{**}\} \subseteq \phi_\pi(M_l(A))$ . Let  $Q \in \text{Hom}_A(A, A^{**})$ . Then, by Lemma 2.2 and the fact that  $(A \circ A^*)^* \cong \text{Hom}_A(A, A^{**})$ , there exists a net  $\{F_\alpha\}$  in  $A^{**}$  such that  $(f, T_{F_\alpha}(a)) \rightarrow (f, Q(a))$  for all  $a \in A, f \in A^*$ . Let  $T_\alpha \in M_l(A)$  be such that  $T_{F_\alpha} = \pi \circ T_\alpha$ , for all  $\alpha$ . Then  $(f, T_{F_\alpha}(a)) = f(T_\alpha(a))$  since  $T_\alpha(a) \in A$ . But, by [8, Theorem 4.7, p. 286],  $M_l(A)$  is  $\tau_l$ -complete. Hence there exists  $T \in M_l(A)$  such that  $f(T_\alpha(a)) \rightarrow f(T(a))$ , for all  $a \in A, f \in A^*$ . This shows that  $\pi(T(a))f = (f, Q(a))$ , for all  $a \in A, f \in A^*$ , or equivalently,  $\pi(T(a)) = Q(a)$ , for all  $a \in A$ , i.e.,  $Q = \pi \circ T$ . Thus  $Q \in \phi_\pi(M_l(A))$  and so  $\text{Hom}_A(A, A^{**}) = \phi_\pi(M_l(A))$ . Since  $(A \circ A^*)^* \cong \text{Hom}_A(A, A^{**})$  and since  $(A \circ A^*)^*$  is the  $\sigma$ -closure of  $\{\mathfrak{F}_F: F \in A^{**}\}$ , it follows that  $\phi_\pi(M_l(A))$  is the ultraweak closure of  $\{T_F: F \in A^{**}\}$ .

(ii)  $\Rightarrow$  (i). Suppose (ii) holds. Then, in view of Lemma 2.2,  $M_l(A)$  is isometrically isomorphic to  $(A \circ A^*)^*$ . From Lemma 2.1 we obtain  $\mathfrak{S}((A \circ A^*)^*)$  is  $\sigma$ -compact, and since  $\tau_l$  is weaker than the ultraweak topology on  $M_l(A)$ , it follows that  $\mathfrak{S}(M_l(A))$  is  $\tau_l$ -compact and therefore  $\tau_l$ -complete. Let  $I$  be the identity element of  $M_l(A)$ . Since  $\{T_F: F \in A^{**}\}$  is ultraweak dense in  $\phi_\pi(M_l(A))$ , there exists a net  $\{F_\alpha\}$  in  $A^{**}$  such that  $T_{F_\alpha}$  converges ultraweakly to  $\pi \circ I = \pi$ , or equivalently,  $\mathfrak{F}_{F_\alpha}$   $\sigma$ -converges to  $\mathfrak{F}_\pi$ . Since  $\pi(A)$  is  $w^*$ -dense in  $A^{**}$  and the  $w^*$ -topology is stronger than the  $\sigma$ -topology on  $A^{**}$ , it follows that there exists a net  $\{u_\alpha\}$  in  $A$  such that  $\{\mathfrak{F}_{\pi(u_\alpha)}\}$   $\sigma$ -converges to  $\mathfrak{F}_\pi$  and so, by Theorem 2.3,  $A$  has property (P2). Therefore, by [8, Theorem 4.7, p. 287],  $A$  is dual.

**COROLLARY 3.2.** *Let  $A$  be an  $A^*$ -algebra of the first kind. Then  $A$  is dual if and only if  $\phi_\pi(M_l(A)) = \phi((A \circ A^*)^*)$ .*

**COROLLARY 3.3.** *Let  $A$  be a modular annihilator  $A^*$ -algebra of the first kind. If  $\text{Hom}_A(A, A^{**})$  is the ultraweak closure of  $\{T_F: F \in A^{**}\}$  then  $A$  is dual.*

**PROOF.** By [9, Theorem 5.2, p. 830],  $\pi(A)$  is an ideal of  $A^{**}$  so that  $T_F$  maps  $A$  into  $\pi(A)$  for every  $F \in A^{**}$ . Hence if  $\text{Hom}_A(A, A^{**})$  is the ultraweak closure of  $\{T_F: F \in A^{**}\}$ , then  $\text{Hom}_A(A, A^{**}) = \phi_\pi(M_l(A))$  by the proof above. Therefore  $A$  is dual by Theorem 3.1.

**4. A realization of the algebra  $M_l(A)$ .**

**THEOREM 4.1.** *Let  $A$  be a dual  $A^*$ -algebra of the first kind and let  $\mathfrak{A}$  be its completion. Let  $\pi_{\mathfrak{A}}$  be the canonical map of  $\mathfrak{A}$  into  $\mathfrak{A}^{**}$ . Then  $\pi_{\mathfrak{A}}(A)$  is an ideal of  $\mathfrak{A}^{**}$  when  $\mathfrak{A}^{**}$  is given Arens product.*

**PROOF.** Let  $x \in A, F \in \mathfrak{A}^{**}$  and let  $\{e_\alpha\}$  be a maximal orthogonal family of selfadjoint minimal idempotents in  $A$ . By [4, Theorem 16, p. 30],  $\sum_\alpha e_\alpha x$  is summable to  $x$  in the norm  $\|\cdot\|$ , and hence there exists only a countable number of  $e_\alpha$  for which  $e_\alpha x \neq 0$ , say  $e_{\alpha_1}, e_{\alpha_2}, \dots$ . Since  $A$  and  $\mathfrak{A}$  have the

same socle and  $\pi_{\mathfrak{A}}(\mathfrak{A})$  is an ideal of  $\mathfrak{A}^{**}$ , it follows that  $F * \pi_{\mathfrak{A}}(e_{\alpha_i}) \in \pi_{\mathfrak{A}}(A)$  for  $i = 1, 2, \dots$ . Let  $m, n$  be positive integers,  $m < n$ . By [4, Lemma 4, p. 18], we have

$$\begin{aligned} & \left\| \sum_{i=1}^n F * \pi_{\mathfrak{A}}(e_{\alpha_i}) \pi_{\mathfrak{A}}(x) - \sum_{i=1}^m F * \pi_{\mathfrak{A}}(e_{\alpha_i}) * \pi_{\mathfrak{A}}(x) \right\| \\ &= \left\| \left( \sum_{i=m+1}^n F * \pi_{\mathfrak{A}}(e_{\alpha_i}) \right) \left( \sum_{i=m+1}^n \pi_{\mathfrak{A}}(e_{\alpha_i}) * \pi_{\mathfrak{A}}(x) \right) \right\| \\ &\leq k \left\| \left( \sum_{i=m+1}^n F * \pi_{\mathfrak{A}}(e_{\alpha_i}) \right) \left( \sum_{i=m+1}^n \pi_{\mathfrak{A}}(e_{\alpha_i}) * \pi_{\mathfrak{A}}(x) \right) \right\| \\ &\leq k|F| \left\| \sum_{i=m+1}^n \pi_{\mathfrak{A}}(e_{\alpha_i}) \right\| \left\| \sum_{i=m+1}^n \pi_{\mathfrak{A}}(e_{\alpha_i}) * \pi_{\mathfrak{A}}(x) \right\| \\ &\leq k|F| \left\| \sum_{i=m+1}^n \pi_{\mathfrak{A}}(e_{\alpha_i}) * \pi_{\mathfrak{A}}(x) \right\|, \end{aligned}$$

where  $|F|$  denotes the norm of  $F$  in  $\mathfrak{A}^{**}$  and  $k$  is a positive constant. Thus  $\{\sum_{i=1}^n F * \pi_{\mathfrak{A}}(e_{\alpha_i}) * \pi_{\mathfrak{A}}(x)\}$  is a Cauchy sequence in  $\pi_{\mathfrak{A}}(A)$  with respect to the norm  $\|\cdot\|$ , and so there exists  $z \in A$  such that  $\pi_A(z) = \sum_{i=1}^{\infty} F * \pi_{\mathfrak{A}}(e_{\alpha_i}) * \pi_{\mathfrak{A}}(x)$ . Since  $\sum_{i=1}^{\infty} F * \pi_{\mathfrak{A}}(e_{\alpha_i}) * \pi_{\mathfrak{A}}(x)$  also converges to  $\pi_{\mathfrak{A}}(z)$  and to  $F * \pi_{\mathfrak{A}}(x)$  in the norm  $|\cdot|$ , we have  $\pi_{\mathfrak{A}}(z) = F * \pi_{\mathfrak{A}}(x)$ . Hence  $F * \pi_{\mathfrak{A}}(x) \in \pi_{\mathfrak{A}}(A)$ , for all  $x \in A$  and  $F \in \mathfrak{A}^{**}$ . Similarly we can show that  $\pi_{\mathfrak{A}}(x) * F \in \mathfrak{A}^{**}$ , for all  $x \in A$  and  $F \in \mathfrak{A}^{**}$ . Therefore  $\pi_{\mathfrak{A}}(A)$  is an ideal of  $\mathfrak{A}^{**}$ .

**THEOREM 4.2.** *Let  $A$  be a dual  $A^*$ -algebra of the first kind and  $\mathfrak{A}$  its completion. Then  $M_l(A)$  is algebra isomorphic to  $\mathfrak{A}^{**}$  when  $\mathfrak{A}^{**}$  is given Arens product. This isomorphism is given by the following relation: For each  $T \in M_l(A)$  there exists a unique  $F_T \in \mathfrak{A}^{**}$  such that*

$$\pi_{\mathfrak{A}}(Tx) = F_T * \pi_{\mathfrak{A}}(x) \quad (x \in A).$$

**PROOF.** For each  $x \in A$ , let  $\|x\|'_A = \sup\{\|xy\|: \|y\| \leq 1, y \in A\}$ . Then  $\|\cdot\|'_A$  is a norm on  $A$  which is equivalent to  $|\cdot|$  [4, Theorem 18, p. 31]. Hence if  $T \in M_l(A)$  and  $x \in A$ , then

$$\begin{aligned} \|Tx\|'_A &= \sup\{\|T(x)y\|: \|y\| \leq 1, y \in A\} \\ &= \sup\{\|T(xy)\|: \|y\| \leq 1, y \in A\} \\ &\leq \|T\| \sup\{\|xy\|: \|y\| \leq 1, y \in A\} \\ &\leq k' \|T\| |x|, \end{aligned}$$

where  $k'$  is a constant  $> 0$ . Thus  $|Tx| \leq k''|x|$  for all  $x \in A$  and some constant  $k'' > 0$ . Since  $A$  is dense in  $\mathfrak{A}$ , it follows that  $T$  has a unique bounded extension  $T'$  to  $\mathfrak{A}$ . Clearly  $T' \in M_l(\mathfrak{A})$ . By [2, Corollary 3.2, p. 509], there exists a unique  $F_T \in \mathfrak{A}^{**}$  such that  $\pi_{\mathfrak{A}}(Tx) = F_T * \pi_{\mathfrak{A}}(x)$  for all  $x \in A$ .

Since, by Theorem 4.1,  $\pi_{\mathfrak{A}}(A)$  is an ideal of  $\mathfrak{A}^{**}$ , we have that  $T \rightarrow F_T$  is an algebra isomorphism of  $M_I(A)$  onto  $\mathfrak{A}^{**}$ .

## REFERENCES

1. B. A. Barnes, *Subalgebras of modular annihilator algebras*, Proc. Cambridge Philos. Soc. **66** (1969), 5–12. MR **41** #4236.
2. B. D. Malviya and B. J. Tomiuk, *Multiplier operators on  $B^*$ -algebras*, Proc. Amer. Math. Soc. **31** (1972), 505–510. MR **46** #4215.
3. L. Máté, *On representation of module-homomorphisms (multipliers)*, Studia Sci. Math. Hungar. **8** (1973), 187–192. MR **51** #1286.
4. T. Ogasawara and K. Yoshinaga, *Weakly completely continuous Banach\*-algebras*, J. Sci. Hiroshima Univ. Ser. A **18** (1954), 15–36. MR **16**, 1126.
5. C. E. Rickart, *General theory of Banach algebras*, Van Nostrand, Princeton, N.J., 1960. MR **22** #5903.
6. M. Rieffel, *Induced Banach representations of Banach algebras and locally compact groups*, J. Functional Analysis **1** (1967), 443–491. MR **36** #6544.
7. H. H. Schaefer, *Topological vector spaces*, Macmillan, New York, 1966. MR **33** #1689.
8. B. J. Tomiuk, *Multipliers and duality in  $A^*$ -algebras*, Proc. Amer. Math. Soc. **50** (1975), 281–288. MR **51** #2834.
9. Pak-ken Wong, *Modular annihilator  $A^*$ -algebras*, Pacific J. Math. **37** (1971), 825–834. MR **46** #4231.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OTTAWA, OTTAWA, ONTARIO, CANADA K1N 6N5