

DENSE SUBGROUPS OF LIE GROUPS

DAVID ZERLING

ABSTRACT. Let G be a dense analytic subgroup with compact center of an analytic group L . Then there exist closed vector subgroups W and U of G and a (CA) closed normal analytic subgroup M of G , which contains the center of G , such that $G = MWU$, $MW \cap U = M \cap W = \{e\}$, and WU is a closed vector subgroup of G . Moreover, $L = MW\bar{U}$, where MW is a closed normal analytic subgroup of L and \bar{U} is a toral group, such that $MW \cap \bar{U}$ is finite.

1. Introduction. By an analytic group and an analytic subgroup of a Lie group, we mean a connected Lie group and a connected Lie subgroup, respectively. If G and H are Lie groups and φ is a one-to-one (continuous) homomorphism from G into H , φ will be called an immersion. φ will be called closed or dense, as $\varphi(G)$ is closed or dense in H . G_0 and $Z(G)$ will denote the identity component group and center of G , respectively.

If G is an analytic group, $A(G)$ will denote the Lie group of all (bicontinuous) automorphisms of G , topologized with the generalized compact-open topology. G will be called (CA) if $I(G)$, the Lie group of all inner automorphisms of G , is closed in $A(G)$. It is well known that G is (CA) if and only if its universal covering group is (CA).

If G is a normal analytic subgroup of an analytic group H , then each element h of H induces an automorphism of G , namely, $g \mapsto hgh^{-1}$. We will denote this homomorphism from H into $A(G)$ by ρ_{GH} . $I_H(h)$ will denote the inner automorphism of H determined by $h \in H$. More generally, if A is a subset of H , $I_H(A)$ will denote the set of all inner automorphisms of H determined by elements of A . $I_H(H)$ will be written as $I(H)$, and the mapping $h \mapsto I_H(h)$ of H onto $I(H)$ will be denoted by I_H .

If N is an analytic group and H is an analytic subgroup of $A(N)$, then $N \circledast H$ will denote the semidirect product of N and H . On the other hand, if G is an analytic group containing a closed normal analytic subgroup N and a closed analytic subgroup H , such that $G = NH$, $N \cap H = \{e\}$, and such that the restriction of ρ_{NG} to H is one-to-one, we will frequently identify G with $N \circledast \rho_{NG}(H)$ and H with $\rho_{NG}(H)$, that is, we may write $G = N \circledast H$.

In Zerling [3] we proved the following theorem.

MAIN STRUCTURE THEOREM. *Let G be a non-(CA) analytic group. Then there exist a (CA) analytic group M , a toral group T in $A(M)$, and a dense vector*

Presented to the Society, January 27, 1977; received by the editors July 27, 1976.

AMS (MOS) subject classifications (1970). Primary 22E15; Secondary 57E05.

© American Mathematical Society 1977

subgroup V of T , such that:

- (i) $H = M \otimes T$ is a (CA) analytic group.
- (ii) G is isomorphic to the dense analytic subgroup $M \otimes V$ of H .
- (iii) $Z(G)$ is contained in M .
- (iv) $Z_0(G) = Z_0(H)$, and $\pi(Z(H))$ is finite, where π is the natural projection of H onto T . Moreover, if $G/Z(G)$ is homeomorphic to Euclidean space, then $Z(G) = Z(H)$.
- (v) Each automorphism σ of G can be extended to an automorphism $\varepsilon(\sigma)$ of H , such that $\varepsilon: A(G) \rightarrow A(H)$ is a closed immersion.

We will now use this theorem in §2 to obtain our main results.

2. Main results.

LEMMA. *Let us maintain the notation of the main structure theorem and let $f: G \rightarrow L$ be a dense immersion of G into an analytic group L . If $Z(G)$ is compact, then $f(M)$ is closed in L .*

PROOF. Since G is non-(CA) we can appeal to Goto [1]: Let N be a maximal analytic subgroup of $I(G)$, which contains the commutator subgroup of $I(G)$ and is closed in $A(G)$. Then there is a closed vector subgroup V' of $I(G)$, such that $I(G) = NV'$, $N \cap V' = \{e\}$, and $\overline{I(G)} = N \cdot \overline{V'}$, where $T' = \overline{V'}$ is a toral group. Moreover, $N \cap T'$ is finite, and the space of $\overline{I(G)}$ is diffeomorphic to the product space $N \times T'$.

In the proof of the main structure theorem in Zerling [3], H is constructed in such a way that $\rho_{GH}(M) = N$, $\rho_{GH}(V) = V'$, and $\rho_{GH}(T) = T'$. Also, ρ_{GH} is 1-1 on T .

Since $Z(G)$ is of finite index in $Z(H)$ from Zerling [4, Lemma 2.1], $Z(H)$ is also compact. Consider the normal analytic subgroup $\overline{f(M)} \cdot f(V)$ of L . Since the inner automorphic action of $f(V)$ on $\overline{f(M)}$ is effective, we have the Lie group $P = \overline{f(M)} \otimes V$. The image of each one-parameter subgroup of V under ρ_{GP} is not closed in $A(G)$. Therefore, since G is dense in P , we see from Lemma 3.1 of Zerling [4] that the closure of V in $A(\overline{f(M)})$ is a toral group, which we will denote by T_1 .

Now let $Q = \overline{f(M)} \otimes T_1$. Then $\rho_{GQ}(T_1) = T'$, and since $\tau_1 \cdot (m, v) \cdot \tau_1^{-1} = (\tau_1(m), v)$ for all (m, v) in G , we see that ρ_{GQ} is 1-1 on T_1 . Since $\rho_{GH}(T) = T'$, and ρ_{GH} is 1-1 on T , we have that $\rho_{GQ}^{-1} \circ \rho_{GH}$ is an isomorphism of T onto T_1 . Hence, $H = M \otimes T$ is a dense (CA) analytic subgroup of Q . Since $Z(H)$ is compact, we may appeal to van Est [2, Theorem 2.2.1] to conclude that $H \cong Q$. Hence, $f(M) = \overline{f(M)}$ and our theorem is proved.

THEOREM. *Let $f: G \rightarrow L$ be a proper dense immersion of an analytic group G into an analytic group L . Suppose $Z(G)$ is compact. Then there exist closed vector subgroups W and U of G and a (CA) closed normal analytic subgroup M of G , which contains $Z(G)$, such that:*

- (i) $G = MWU$, $MW \cap U = M \cap W = \{e\}$, and WU is a closed vector subgroup of G .

(ii) $L = f(MW) \cdot \overline{f(U)}$, where $f(MW)$ is a closed normal analytic subgroup of L and $\overline{f(U)}$ is a toral group. Moreover, $f(MW) \cap \overline{f(U)}$ is finite, and $f(W) \cap (f(M) \cdot \overline{f(U)}) = f(W) \cap Z(L) = \{e\}$.

(iii) If L is (CA) and $Z(L)$ is compact, then $W = \{e\}$.

PROOF. Since $Z(G)$ is compact, we can conclude from van Est [2] that G is non-(CA). We will now maintain the notation of the main structure theorem, as well as the notation in the proof of the above lemma.

Since $Z(G)$ is compact, we know from the above lemma that there exists a maximal analytic subgroup J of G which contains M , such that $f(J)$ is closed in L . Then from Goto [1] there exists a closed vector subgroup U of G such that $G = JU$, $J \cap U = \{e\}$. Moreover, $L = f(J) \cdot \overline{f(U)}$, where $\overline{f(U)}$ is a toral group and $f(J) \cap \overline{f(U)}$ is finite.

In the proof of Goto's theorem [1], applied to $I(G)$, T' is a closed central subgroup of an arbitrarily fixed maximal compact subgroup K of $I(G)$. We will assume that K has been selected so that it contains $\rho_{GL}(\overline{f(U)})$.

Now let $\pi: G \rightarrow V$ be the natural projection and let $W = \pi(J)$. Then W is a closed vector subgroup of V and since J contains M we see that $J = M \cdot W$, $M \cap W = \{e\}$. Therefore,

$$L = f(J) \cdot \overline{f(U)} = f(M) \cdot f(W) \cdot \overline{f(U)} = f(M) \cdot \overline{f(U)} \cdot f(W),$$

where $f(M) \cdot \overline{f(U)}$ is a closed normal analytic subgroup of L . Since $\overline{f(U)} \cap f(J)$ is finite and contained in $f(G)$, it is contained in $f(M)$. Hence, if $(f(M) \cdot \overline{f(U)}) \cap f(W) \neq \{e\}$, then $f(w) = f(m) \cdot x$, $x \in \overline{f(U)}$, and so $x = f(m)^{-1} \cdot f(w)$. Hence, $x \in \overline{f(U)} \cap f(J)$, which is contained in $f(M)$. By the uniqueness of the decomposition in J , we have $w = e$. So $L = (f(M) \cdot \overline{f(U)}) \cdot f(W)$, $(f(M) \cdot \overline{f(U)}) \cap f(W) = \{e\}$. Moreover, $f(W) \cap Z(L) = \{e\}$, since $f(J) \cap Z(L)$ is contained in $f(M)$, and $W \cap M = \{e\}$. Hence, $L = (f(M) \cdot \overline{f(U)}) \otimes f(W)$.

We now show that WU is a closed vector subgroup of $G = MWU$. Let $\varphi: W \rightarrow A(MU)$ be given by $\varphi(w)(mu) = w(mu)w^{-1}$. Since $W \cap Z(G) = \{e\}$, φ is an immersion and so $G = MU \otimes W$. Since the image of each one-parameter subgroup of W under I_G is not closed in $A(G)$, we see from Zerling [4] that $\overline{\varphi(W)}$ is a toral group.

Let $u \in U$. Then $I_G(\varphi(w) \cdot u) = I_G(wuw^{-1}) = I_G(u)$, since $I_G(w) \in T'$ commutes with $I_G(u) \in \rho_{GL}(\overline{f(U)})$ because of our selection of K above. Hence $\sigma(u) \cdot u^{-1} \in Z(G)$ for all $\sigma \in \overline{\varphi(W)}$. Since $Z(G)$ is a closed central subgroup of MU and each element of $\overline{\varphi(W)}$ keeps $Z(G)$ elementwise fixed, we see from Zerling [3, Lemma 2.2] that $\sigma(u) = u$ for all $\sigma \in \overline{\varphi(W)}$. Therefore, UW is a closed vector subgroup of G .

Now suppose that L is (CA) and $Z(L)$ is compact. As we did above, we can show that the closure of $f(W)$ in $A(f(M) \cdot \overline{f(U)})$ is a toral group, call it T_2 . Then L is properly dense in $(f(M) \cdot \overline{f(U)}) \otimes T_2$. This is a contradiction from van Est [2]. Hence $W = \{e\}$. This completes the proof of our theorem.

3. **An example.** The following example shows that in the above theorem W need not be $\{e\}$, even if L is (CA). Let $G = MV$ be any non-(CA) analytic group and suppose that the dimension of V is at least two. Such a group G can easily be obtained by a slight modification of the example in Zerling [3].

We continue with our previous notation and let $S = G \otimes T'$. Since V is dense in T we can select an element $v_0 \in V$ such that v_0 generates a dense subgroup of T . Then $v'_0 = I_G(v_0)$ generates a dense subgroup of T' . Let D denote the subgroup of S generated by $(v_0, v'_0)^{-1}$. Then D is a free discrete central subgroup of S and $L = S/D$ is a (CA) analytic group for which $g \mapsto (g, e)D$ is a dense immersion f of G into L ; see Zerling [4, the proof of Theorem 2.2].

Next let V_λ be the one-dimensional vector subgroup of V containing v_0 , and let V_μ be a vector subgroup of V such that $V = V_\lambda \cdot V_\mu$, $V_\lambda \cap V_\mu = \{e\}$. Let $\delta: S \rightarrow L$ be the canonical projection. Then $G = MV_\mu V_\lambda$, $f(MV_\mu)$ is closed in L and $\overline{f(V_\lambda)} = f(V_\lambda) \cdot \delta(T')$ is a toral group. $L = f(MV_\mu) \cdot \overline{f(V_\lambda)}$, and $f(MV_\mu) \cap \overline{f(V_\lambda)}$ is trivial. Hence $V_\mu = W \neq \{e\}$.

BIBLIOGRAPHY

1. M. Goto, *Analytic subgroups of $GL(n, \mathbf{R})$* , Tôhoku Math. J. (2) **25** (1973), 197–199. MR **48** #463.
2. W. T. van Est, *Dense imbeddings of Lie groups*, Nederl. Akad. Wetensch. Proc. Ser. A **54** = Indag. Math. **13** (1951), 321–328. MR **13**, 432.
3. D. Zerling, *Some theorems on (CA) analytic groups*, Trans. Amer. Math. Soc. **205** (1975), 181–192. MR **51** #802.
4. ———, *Some theorems on (CA) analytic groups. II*, Tôhoku Math. J. (to appear).

DEPARTMENT OF MATHEMATICS AND PHYSICS, PHILADELPHIA COLLEGE OF TEXTILES AND SCIENCE, PHILADELPHIA, PENNSYLVANIA 19144.