

## INDECOMPOSABLE COMPACT PERTURBATIONS OF THE BILATERAL SHIFT

DOMINGO A. HERRERO

**ABSTRACT.** Recent results of M. Radjabalipour and H. Radjavi assert that the sum of a normal operator  $N$  with spectrum on a smooth Jordan curve and a compact operator  $K$  in the Macaev ideal  $\mathfrak{S}_\omega$  is decomposable provided the spectrum of  $N + K$  does not fill the interior of the curve. Examples are given to show that this result cannot be essentially improved by taking  $K$  in a larger ideal.

1. Let  $\mathcal{L}(\mathcal{H})$  be the algebra of all (bounded linear) operators on a complex Hilbert space  $\mathcal{H}$ . The *Macaev ideal*  $\mathfrak{S}_\omega$  is the set of all compact operators  $K$  in  $\mathcal{L}(\mathcal{H})$  such that  $\sum_1^\infty \mu(n)/n < \infty$ , where  $\mu(1), \mu(2), \dots$ , are the eigenvalues of  $(K^*K)^{1/2}$  arranged in decreasing order and repeated according to multiplicity.

Compact perturbations of normal operators with spectrum on a smooth Jordan curve by an operator  $K \in \mathfrak{S}_\omega$  are known to have a rich family of invariant subspaces. To make this precise, several definitions will be necessary:

An invariant subspace  $\mathfrak{M}$  of  $T \in \mathcal{L}(\mathcal{H})$  is a *maximal spectral subspace* of  $T$  if  $\mathfrak{N} \subset \mathfrak{M}$  for all invariant subspaces  $\mathfrak{N}$  of  $T$  such that the spectrum  $\Lambda(T|\mathfrak{N})$  of the restriction of  $T$  to  $\mathfrak{N}$  is contained in  $\Lambda(T|\mathfrak{M})$ .  $T$  is *decomposable* (in the sense of [1]) if for every finite covering  $G_j, j = 1, 2, \dots, n$ , of  $\Lambda(T)$  there exists a set of maximal spectral subspaces  $\mathfrak{H}_j, j = 1, 2, \dots, n$ , of  $T$  such that  $\Lambda(T|\mathfrak{H}_j) \subset G_j, j = 1, 2, \dots, n$ , and  $\mathcal{H} = \mathfrak{H}_1 + \mathfrak{H}_2 + \dots + \mathfrak{H}_n$ . Moreover,  $T$  is called *strongly decomposable* if  $T|\mathfrak{M}$  is decomposable for every maximal spectral subspace  $\mathfrak{M}$ .

M. Radjabalipour and H. Radjavi [10]–[13] have improved a result of V. I. Macaev about compact perturbations of hermitian operators [9] by proving the following:

(i) Let  $T$  be the sum of an operator  $A$  having spectrum on a  $C^2$  Jordan curve  $J$  and a compact operator  $K \in \mathfrak{S}_\omega$ . Assume that  $\|(z - A)^{-1}\| \leq C/d(z)$ , where  $d(z)$  is the distance from  $z$  to  $\Lambda(A)$  and  $C \geq 1$  is a constant

---

Received by the editors April 2, 1975 and, in revised form, June 5, 1975, October 21, 1975 and May 24, 1976.

*AMS (MOS) subject classifications* (1970). Primary 47A15, 47A55.

*Key words and phrases.* Bilateral weighted shifts, decomposable operators, Macaev ideal of compact operators.

© American Mathematical Society 1977

independent of  $z$ , and that  $\Lambda(T)$  does not fill the interior of  $J$ . Then  $T$  is strongly decomposable.

(ii) Assume that  $\Lambda(T)$  is contained in a  $C^2$  Jordan curve  $J$  and that there exist a positive number  $\epsilon$  and a nonincreasing function  $M: (0, \epsilon) \rightarrow (0, \infty)$  such that  $\int_0^\epsilon \log^{(2)} M(t) dt < \infty$  ( $\log^{(m)} x$  denotes the  $m$ th iterated logarithm). If  $\|(z - T)^{-1}\| \leq M[d(z)]$  for  $z \notin J$ , then  $T$  is strongly decomposable. This is true, in particular, if  $M(t) = \exp(\exp t^{-p})$ ,  $0 < p < 1$ .

It will be shown that these two results are essentially sharp by proving that compact perturbations of the (unitary!) bilateral shift  $U$  in  $\ell^2$  (defined by  $Ue_n = e_{n+1}$  for all  $n$  in the set  $\mathbf{Z}$  of all integers, where  $\{e_n\}$  is the canonical basis of  $\ell^2$ ) by an operator in a certain class of ideals which are only "slightly larger" than  $\mathfrak{S}_\omega$  fail to be decomposable.

The author wishes to thank Professor Mehdi Radjabalipour for making available his unpublished papers and the referee for pointing out a mistake in the proof of part (a) of the theorem and for providing a suggestion which made it possible to simplify the proof of part (c).

2. Let  $\{a_n\}$  be a bounded two-sided sequence of positive reals and define the bilateral weighted shift  $B$  in  $\ell^2$  by  $Be_n = a_n e_{n+1}$ ,  $n \in \mathbf{Z}$ . If  $\lim_{n \rightarrow \pm\infty} a_n = 1$ , then  $B - U$  is compact and  $\Lambda(B)$  is the boundary  $\partial D$  of the unit disc  $D = \{z: |z| < 1\}$ .

Define  $w_0 = 1$ ,  $w_n = a_0 \cdot \dots \cdot a_{n-1}$  and  $w_{-n} = (a_{-1} a_{-2} \cdot \dots \cdot a_{-n})^{-1}$  for all positive  $n$ . Then  $B$  is unitarily equivalent to multiplication by  $z$  on the space

$$L^2(\mathbf{w} = \{w_n\}) = \left\{ f(z) = \sum_{n \in \mathbf{Z}} b_n z^n: \|f\|^2 = \sum |b_n w_n|^2 < \infty \right\}$$

of formal Laurent series [3], [6]. Now we are in a position to construct the counterexamples.

**THEOREM.** *Let  $\varphi = \{\varphi_n\}_1^\infty$  be a nonincreasing sequence of positive reals and assume that: (i)  $\{\varphi_n \log en\}$  is nonincreasing, (ii)  $n(\varphi_n - \varphi_{n+1}) \rightarrow 0$ , and (iii)  $\sum \varphi_n/n = \infty$ . Let  $B$  be the bilateral weighted shift defined by  $Be_n = a_n e_{n+1}$ , where  $a_n = w_{n+1}/w_n$ ,  $w_0 = 1$  and  $w_n = w_{-n} = \exp(n\varphi_n)$  for  $n = 1, 2, \dots$ . Then*

(a) *Every nonzero invariant subspace  $\mathfrak{N}$  of  $B$  is either invariant under  $B^{-1}$  and satisfies  $\Lambda(B|\mathfrak{N}) = \partial D = \Lambda(B)$ , or it is not invariant under  $B^{-1}$  and  $\Lambda(B|\mathfrak{N}) = D^-$  is the closed unit disc.*

(b)  *$K = B - U$  is a compact operator such that the eigenvalues of  $(K^*K)^{1/2}$  (ordered as above indicated) satisfy  $\mu_{2n-1}, \mu_{2n} \leq C\varphi_n$ , where  $C$  is a constant independent of  $n$ .*

(c)  $\log^{(2)}\|(z - B)^{-1}\| \leq M/d(z)$ , for a suitable constant  $M$ .

**PROOF.** (a) This will follow by using the same kind of "sandwich theory" as in [4]:

Let  $L^1(\mathbf{w}) = \{f(z) = \sum b_n z^n: \|f\|_1 + \sum |b_n w_n| < \infty\}$  and let  $L^1(\sqrt{\mathbf{w}})$  be similarly defined, with  $w_n$  replaced by  $\sqrt{w_n}$ ,  $n \in \mathbf{Z}$  (the norm of an element of  $L^1(\sqrt{\mathbf{w}})$  will be denoted by  $\|\cdot\|''$ ). Then, for every finite sum  $f(z) = \sum b_n w_n$ , by the Cauchy-Schwarz inequality we have

$$\begin{aligned} \|f\|'' &= \sum |b_n \sqrt{w_n}| = \sum |b_n w_n| / \sqrt{w_n} \\ &\leq \left(\sum |b_n w_n|^2\right)^{1/2} \left(\sum 1/w_n\right)^{1/2} = C \|f\|, \end{aligned}$$

where  $C = (\sum 1/w_n)^{1/2} < \infty$  is a constant independent of  $f$ . It readily follows that  $(1/C)\|f\|'' \leq \|f\| \leq \|f\|_1$  and, therefore, that  $L^1(\mathbf{w}) \subset L^2(\mathbf{w}) \subset L^1(\sqrt{\mathbf{w}})$  and both inclusions are continuous.

It follows from (i) and (ii) that  $\lim_{n \rightarrow \pm\infty} a_n = 1$  and, therefore,  $B$  is a compact perturbation of  $U$  and  $\Lambda(B) = \partial D$  (see [3], [4], [6]). Moreover, it follows from (i)–(iii) and the results of [2], [4] that  $L^1(\mathbf{w})$  and  $L^1(\sqrt{\mathbf{w}})$  are actually algebras (under pointwise multiplication) of quasi-analytic functions defined on  $\partial D$ . I.e.,  $L^1(\sqrt{\mathbf{w}})$  is contained in  $C^\infty(\partial D)$ ; if  $f, g \in L^1(\mathbf{w})$  ( $L^1(\sqrt{\mathbf{w}})$ ), then  $f(z) \cdot g(z) \in L^1(\mathbf{w})$  ( $L^1(\sqrt{\mathbf{w}})$ , resp.) and the vanishing of  $f(z) \in L^1(\sqrt{\mathbf{w}})$  together with all its derivatives at some point  $\lambda \in \partial D$  implies that  $b_n = 0$  for all  $n$ .

Let  $\mathfrak{N}$  be a nonzero invariant subspace of  $B$ . If  $\mathfrak{N}$  is not invariant under  $B^{-1}$ , then  $0 \in \Lambda(B|\mathfrak{N})$  and it follows that  $\Lambda(B|\mathfrak{N}) = D^-$ ; if  $\mathfrak{N}$  is also invariant under  $B^{-1}$ , then  $\Lambda(B|\mathfrak{N}) \subset \partial D$  (these two results easily follow from [7], [8]).

In the second case, the closure  $\mathfrak{N}$  of  $\mathfrak{N}$  in  $L^1(\sqrt{\mathbf{w}})$  is an ideal of this algebra and we can proceed as in [4] in order to show that

$$\mathfrak{N} = \bigcap_{j=1}^m \{f \in L^1(\sqrt{\mathbf{w}}): f(z_j) = f'(z_j) = \dots = f^{(m_j-1)}(z_j) = 0\}$$

for a finite set of points  $z_j \in \partial D$ ,  $j = 1, 2, \dots, m$ , and positive integers  $m_j$ ,  $j = 1, 2, \dots, m$ , so that  $\dim L^1(\sqrt{\mathbf{w}})/\mathfrak{N} = n = \sum_{j=1}^m m_j < \infty$ . Clearly,  $\mathfrak{N} \subset \mathfrak{N} \cap L^2(\mathbf{w})$  and for every  $\lambda \in \partial D \setminus \{z_1, z_2, \dots, z_m\}$  there exists a function  $h_\lambda \in \mathfrak{N}$  such that  $h_\lambda(\lambda) \neq 0$ .

Let  $\mathfrak{N}_0$  ( $\mathfrak{N}''$ ) be the closure of  $(B - \lambda)\mathfrak{N}$  in  $L^2(\mathbf{w})$  (in  $L^1(\sqrt{\mathbf{w}})$ , resp.). Since  $(B - \lambda)f(z) = (z - \lambda)f(z)$ , it readily follows that  $\mathfrak{N}'' = \mathfrak{N} \cap \{f \in L^1(\sqrt{\mathbf{w}}): f(\lambda) = 0\}$  has codimension 1 in  $\mathfrak{N}$ . In particular,  $h_\lambda \in \mathfrak{N} \setminus \mathfrak{N}''$ , a fortiori,  $h_\lambda \notin \mathfrak{N}_0$  ( $\mathfrak{N}_0 \subset \mathfrak{N} \cap \mathfrak{N}''$ ) and, therefore,  $(B - \lambda)\mathfrak{N}$  is not dense in  $\mathfrak{N}$ . Hence,  $\lambda \in \Lambda(B|\mathfrak{N})$ .

Since this holds for all but finitely many  $\lambda$ 's in  $\partial D$ , it is not difficult to conclude that  $\partial D \subset \Lambda(B|\mathfrak{N})$  and, therefore,  $\Lambda(B|\mathfrak{N}) = \partial D$ , as promised.

If  $\mathfrak{N}$  contains a nonzero function  $L^1(\mathbf{w})$ , it is easy to see that  $\mathfrak{N} \cap L^1(\mathbf{w})$  is actually an ideal of finite codimension  $n$  in  $L^1(\mathbf{w})$  and, a fortiori, that  $\mathfrak{N}$  also has codimension  $n$  in  $L^2(\mathbf{w})$ . (The details are left to the reader. The author conjectured that  $\mathfrak{N} \cap L^1(\mathbf{w}) \neq \{0\}$  for every invariant subspace  $\mathfrak{N}$ , but was unable to prove it.)

(b) This follows immediately from  $Ke_n = (B - U)e_n = (a_n - 1)e_{n+1}$  (see [5]).

(c) Since  $U$  is normal,  $\gamma(t) = \max\{\|(z - U)^{-1}\|: d(z) = t\} = 1/t$ . On the other hand (b) and condition (i) show that  $\mu_n \leq k/\log en$ , for some constant  $k > 0$  and for all  $n = 1, 2, \dots$ , and, therefore,  $\nu(t) = \max\{n: \mu_n > 1/t\} \leq e^{kt}$ . Now we are in a position to apply Theorem 1 of [13], whence the result follows.  $\square$

3. The sequences

$$\{\varphi_n = \varphi_n(m) = [\log M_1 n \cdot \log^{(2)} M_2 n \cdot \dots \cdot \log^{(m)} M_m n]^{-1}\},$$

where  $M_1 = e$  and  $M_{k+1} = \exp(M_k)$  for  $k = 1, 2, \dots, m - 1$  ( $m = 1, 2, \dots$ ), are concrete examples of sequences satisfying conditions (i)–(iii). Moreover, if  $m_n \nearrow \infty$  and

$$\varphi_n = \psi_n(m_n) = \varphi_n(m_n) / [\log^{(m_n+1)} M_{m_n+1} n]^2$$

satisfies (iii), then it actually satisfies (i)–(iii).

Given an arbitrary increasing function  $h(n)$  such that  $h(1) = 1$  and  $\lim_{n \rightarrow \infty} h(n) = \infty$ , it is not difficult to find a sequence  $\{m_n\}_{n=1}^\infty$  such that  $\{\psi_n(m_n)\}$  satisfies (i)–(iii), but  $\sum_1^\infty \psi_n(m_n) / [nh(n)] < \infty$ . Now we can give a very precise meaning to our introductory sentence “these two results are essentially sharp”:

**COROLLARY.** *Let  $h(n)$  be an arbitrary increasing function such that  $h(1) = 1$  and  $\lim_{n \rightarrow \infty} h(n) = \infty$ , and let  $\mathfrak{S}_h = \{K \in \mathfrak{L}(\ell^2): K \text{ is compact and } \sum \mu(n) / [nh(n)] < \infty\}$ . Then there exists an operator  $K$  in the ideal  $\mathfrak{S}_h$  such that  $U + K$  is indecomposable, even when its resolvent satisfies the growth conditions of the Theorem.*

The proof follows immediately from the Theorem and the above observations.

**4. An example.** Let  $\varphi_n = (\log en \cdot \log^{(2)} e^2 n)^{-1}$ ; then a straightforward computation shows that  $\mu_n$  is of the same order of magnitude as  $\varphi_n$  ( $n = 1, 2, \dots$ ) and that, in this case,  $\nu(t) \leq \exp(kt/\log t)$ , whence (by applying once again Theorem 1 of [13]) we can improve the estimation (c) of the Theorem to

$$\log^{(2)} \|(z - B)^{-1}\| \leq M / [d(z) \cdot \log 1/d(z)],$$

for a suitable constant  $M$  and for  $0 < d(z) < \frac{1}{2}$ .

REFERENCES

1. I. Colojară and C. Foiaş, *Theory of generalized spectral operators*, Gordon and Breach, New York, 1968.
2. Y. Domar, *Harmonic analysis based on certain commutative Banach algebras*, *Acta Math.* **96** (1956), 1–66. MR **17**, 1228.
3. R. Gellar, *Operators commuting with a weighted shift*, *Proc. Amer. Math. Soc.* **23** (1969), 538–545. MR **41** #4277a.

4. R. Gellar and D. A. Herrero, *Hyperinvariant subspaces of bilateral weighted shifts*, Indiana Univ. Math. J. **23** (1973/74), 771–790. MR **48** #9417.

5. I. C. Gohberg and M. G. Kreĭn, *Introduction to the theory of linear nonselfadjoint operators*, “Nauka”, Moscow, 1965; English transl, Transl. Math. Monographs, vol. 18, Amer. Math. Soc., Providence, R.I., 1969. MR **36** #3137; **39** #7447.

6. P. R. Halmos, *A Hilbert space problem book*, Van Nostrand, Princeton, N. J., 1967. MR **34** #8178.

7. D. A. Herrero, *On analytically invariant subspaces and spectra*, Trans. Amer. Math. Soc. (to appear).

8. D. A. Herrero and N. Salinas, *Analytically invariant and bi-invariant subspaces*, Trans. Amer. Math. Soc. **173** (1972), 117–136. MR **47** #856.

9. V. I. Macaev, *On a class of completely continuous operators*, Dokl. Akad. Nauk SSSR **139** (1961), 548–551 = Soviet Math. Dokl. **2** (1961), 972–975. MR **24** #A1617.

10. M. Radjabalipour, *Growth conditions and decomposable operators*, Canad. J. Math. **26** (1974), 1372–1379.

11. ———, *On decomposability of compact perturbations of operators*, Proc. Amer. Math. Soc. **53** (1975), 159–164.

12. M. Radjabalipour and H. Radjavi, *On decomposability of compact perturbations of normal operators*, Canad. J. Math. **27** (1975), 725–735.

13. ———, *On invariant subspaces of compact perturbations of operators* (to appear).

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD SIMÓN BOLÍVAR, CARACAS, VENEZUELA

*Current address:* Departamento de Matemáticas, Instituto Venezolano de Investigaciones Científicas, Caracas, Venezuela