

## ON CALABI'S INHOMOGENEOUS EINSTEIN-KAEHLER MANIFOLDS<sup>1</sup>

JOSEPH A. WOLF

**ABSTRACT.** We use some information on Lie groups to replace a long computation of Calabi, proving that certain complete Einstein-Kaehler manifolds are not locally homogeneous, and finding their isometry groups.

E. Calabi [1] constructed a complete Einstein-Kaehler metric on the tube domain

$$M = \{z = x + iy \in \mathbf{R}^n + i\mathbf{R}^n: \|y\| < r\} \subset \mathbf{C}^n$$

which is invariant under the natural action  $(v, g): x + iy \mapsto v + gx + igy$  of the proper euclidean group  $\mathbf{E}(n) = \mathbf{R}^n \cdot SO(n)$ . He used a rather complicated calculation to show that  $M$  is not homogeneous with that metric. We are going to replace his calculation by a simple group-theoretic argument and obtain a slightly stronger result:

**THEOREM.** *If  $n \geq 2$  and  $ds^2$  is an  $\mathbf{E}(n)$ -invariant Kaehler metric on  $M$ , then  $(M, ds^2)$  cannot be both complete and locally homogeneous, in particular, cannot be homogeneous.*

Here note that Calabi's metric [1] is complete and the flat metric is locally homogeneous.

Finally, we will show that the theorem implies

**COROLLARY.** *If  $n \geq 2$  and  $ds^2$  is an  $\mathbf{E}(n)$ -invariant Kaehler metric on  $M$ , then  $\mathbf{E}(n)$  is the largest connected group of holomorphic isometries of  $(M, ds^2)$ .*

**PROOF OF THEOREM.** Let  $ds^2$  be an  $\mathbf{E}(n)$ -invariant Kaehler metric on  $M$  and  $T^{1,0}(0)$  the holomorphic tangent space at 0. The curvature transformation of  $(M, ds^2)$  at 0 is a linear transformation of  $\Lambda^2 T^{1,0}(0)$  that commutes with the irreducible action of  $SO(n)$  on  $\Lambda^2 T^{1,0}(0)$ , hence is scalar. So  $(M, ds^2)$  has constant holomorphic sectional curvature at 0.

Suppose that  $(M, ds^2)$  is complete and locally homogeneous. Then  $(M, ds^2)$  is complete and simply connected with some constant holomorphic sectional curvature  $c$ , hence holomorphically isometric to a complex projective space ( $c > 0$ ), a complex euclidean space ( $c = 0$ ), or a complex hyperbolic space

---

Received by the editors September 30, 1976 and, in revised form, October 14, 1976.  
*AMS (MOS) subject classifications* (1970). Primary 32M10, 53C30, 53C55; Secondary 20G20, 22E15.

<sup>1</sup>Research partially supported by NSF Grant MCS 76-01692.

( $c < 0$ ). The first two possibilities cannot occur because  $M$  is noncompact and admits nonconstant bounded holomorphic functions such as  $f(z) = 1/(z_1 - 2ir)$ . Thus  $(M, ds^2)$  has holomorphic isometry group  $SU(1, n)$ /(scalars), and so  $\mathbf{E}(n)$  is contained in that group locally isomorphic to  $SU(1, n)$ . The next two lemmas show that this is impossible.

LEMMA. *Let  $G$  be a reductive Lie group. If  $E$  is an analytic subgroup of  $G$  then the solvable radical of  $[E, E]$  is a unipotent subgroup of  $G$ . In particular, if  $n \geq 2$  and  $\mathbf{E}(n)$  is a subgroup of  $G$ , then the translation subgroup  $\mathbf{R}^n \subset \mathbf{E}(n)$  is unipotent in  $G$ .*

PROOF. We may cut  $G$  down to its identity component and then divide out its center, so we may assume that  $G$  is a semisimple linear group. One knows [2, Theorem 3.2, p. 128] that a finite dimensional linear representation of a Lie algebra carries the radical of the derived algebra to an algebra of nilpotent transformations. So the radical of  $[E, E]$  is unipotent in  $G$ . Q.E.D.

LEMMA. *If  $n \geq 2$  and  $G$  is locally isomorphic to the special unitary group  $SU(1, n)$  of Lorentz signature, then  $G$  has no subgroup isomorphic to  $\mathbf{E}(n)$ .*

PROOF. It is known [3, §3] that the maximal unipotent subgroups of  $G$  are isomorphic to the  $(2n - 1)$ -dimensional Heisenberg group  $H_{2n-1} = \mathbf{R} + \mathbf{C}^{n-1}$  with product  $(z, w)(z', w') = (z + z' + \text{Im } w \cdot w', w + w')$  where  $w \cdot w'$  is the usual  $U(n - 1)$ -invariant hermitian scalar product on  $\mathbf{C}^{n-1}$ . An easy calculation with the real symplectic structure underlying  $\mathbf{C}^{n-1}$  shows that every abelian subgroup of  $H_{2n-1}$  is  $U(n - 1)$ -conjugate, hence [3, §3]  $G$ -conjugate, to  $\mathbf{R} + \mathbf{R}^{n-1}$ . Again by [3, §3], the latter has  $G$ -normalizer locally isomorphic to  $H_{2n-1} \cdot (SO(n - 1) \times \mathbf{R})$ , and the latter has no subgroup locally isomorphic to  $SO(n)$ . Q.E.D.

PROOF OF COROLLARY. Let  $G$  be the largest connected group of holomorphic isometries of  $(M, ds^2)$  and  $K = \{g \in G: g(0) = 0\}$ . Then  $z = x + iy \in M$  has  $\mathbf{E}(n)$ -orbit  $\{z' = x' + iy' \in \mathbf{R}^n + i\mathbf{R}^n: \|y'\| = \|y\|\}$ , which has real codimension 1 whenever  $y \neq 0$ . As  $G(z)$  and  $\mathbf{E}(n)(z)$  are complete riemannian submanifolds, and the Theorem ensures that  $G(z)$  has positive real codimension in  $M$ , now  $G(z) = \mathbf{E}(n)(z)$  for  $y \neq 0$ , and, hence, also for the other orbit  $y = 0$ . Now that other orbit  $\mathbf{R}^n = G/K$ , so  $K$  acts on the tangent space to  $M$  at 0 as a subgroup of  $U(n)$  that stabilizes  $\mathbf{R}^n$ . Thus  $K$  coincides with its subgroup  $SO(n)$ , and so by dimension  $G$  coincides with its subgroup  $\mathbf{E}(n)$ . Q.E.D.

#### REFERENCES

1. E. Calabi, *A construction of nonhomogeneous Einstein metrics*, Proc. Sympos. Pure Math., vol. 27, Part II, Amer. Math. Soc., Providence, R. I., 1975, pp. 17-24. MR 52 #816.
2. G. Hochschild, *The structure of Lie groups*, Holden-Day, San Francisco, Calif., 1965. MR 34 #7696.
3. J. A. Wolf, *Representations of certain semidirect product groups*, J. Functional Analysis 19 (1975), 339-372.