

THE RANGE OF A VECTOR MEASURE HAS THE BANACH-SAKS PROPERTY¹

R. ANANTHARAMAN

ABSTRACT. The above result of Diestel and Seifert is proved using the Banach-Saks theorem for $L^p(\lambda)$, $1 < p < \infty$.

Let X be a real Hausdorff quasi-complete locally convex space (L.C.S.), \mathcal{Q} be a α -algebra of subsets of some set S and $\nu: \mathcal{Q} \rightarrow X$ be a measure. We assume that ν is controlled, viz. that there exists a finite positive measure λ on \mathcal{Q} such that ν and λ have the same null sets. When X is a Banach space, Bartle, Dunford and Schwartz [3] proved every measure $\nu: \mathcal{Q} \rightarrow X$ to be controlled. This has been extended to a class of L.C.S. X , including metrizable ones, by Tweddle [8].

A set $A \subset X$ is said to have the *Banach-Saks property* if every sequence in A has a subsequence $\{x_n\}$ such that the sequence of arithmetic means formed from $\{x_n\}$ converges. Using the methods of Szlenk [6], Diestel and Seifert [5] recently proved that $K = \overline{\text{co}} \nu(\mathcal{Q})$ has the Banach-Saks property when X is a Banach space. Our proof of an extension of this result is based instead on the Banach-Saks theorem for $L^p(\lambda)$, $1 < p < \infty$ [2], [4, p. 78].

Since X is quasi-complete, the set $K = \overline{\text{co}} \nu(\mathcal{Q})$ is weakly compact by Theorem 3 of Tweddle [7]. Further, for each $\phi \in L^\infty(\lambda)$ the weak integral $\int \phi d\nu$ belongs to X (see e.g. [1, Lemma 1]). Let $T: L^\infty(\lambda) \rightarrow X$ be the weak integral map, and let T_0 denote its restriction to the weak*-compact convex set

$$P \equiv \{\phi \in L^\infty(\lambda): 0 \leq \phi(s) \leq 1 \text{ s-a.e.}\}.$$

Then $T_0(P) = K$.

As in Proposition 1 of [1] (proved there for $p = 1$), the next result easily follows from Egoroff's theorem.

1. **PROPOSITION.** *Let $1 < p < \infty$. Then the restriction of T to every norm bounded subset B of $L^\infty(\lambda)$ is continuous relative to the L^p -norm on B and the given topology of X .*

2. **THEOREM.** *Let $\nu: \mathcal{Q} \rightarrow X$ be a controlled measure. Then $K = \overline{\text{co}} \nu(\mathcal{Q})$ has the Banach-Saks property.*

Received by the editors March 31, 1977.

AMS (MOS) subject classifications (1970). Primary 28A45.

¹This research was supported by a grant from NRC, Canada.

PROOF. Let $\{x_n\}$ be a sequence in K . Since K is weakly compact, by Eberlein's theorem we may replace $\{x_n\}$ by a subsequence $\{x_m\}$ that converges weakly to some $x \in K$. Since $T(P) = K$, for every m there exists $\phi_m \in P$ such that $T(\phi_m) = x_m$. Fix p , $1 < p < \infty$. As P is a bounded subset of $L^p(\lambda)$, by the Banach-Saks theorem there exists a subsequence $\{\phi_i\}$ of $\{\phi_m\}$ such that the sequence $\{\psi_i\}$ formed of arithmetic means of $\{\phi_i\}$ converges to some $\psi \in L^p(\lambda)$ in the norm of L^p . Clearly, $\psi \in P$. By Proposition 1 and the linearity of T the sequence $\{T(\phi_m)\}$, viz. $\{x_m\}$ has arithmetic means that converge to $T(\psi) \in T(P) = K$. Since $\{x_m\}$ converges weakly to x , we have $T(\psi) = x$, which completes the proof.

REFERENCES

- [1] R. Anantharaman, *On exposed points of the range of a vector measure*, Vector and Operator Valued Measures and Applications (Proc. Sympos. Snowbird Resort, Alta, Utah, 1972), Academic Press, New York, 1973, pp. 7-22.
- [2] S. Banach and S. Saks, *Sur la convergence forte dans les champs L^p* , Studia Math. **2** (1930), 51-57.
- [3] R. G. Bartle, N. Dunford and J. T. Schwartz, *Weak compactness and vector measures*, Canad. J. Math. **7** (1955), 289-305.
- [4] J. Diestel, *Geometry of Banach spaces—Selected topics*, Springer-Verlag, New York, 1975.
- [5] J. Diestel and C. J. Seifert, *An averaging property of the range of a vector measure*, Bull. Amer. Math. Soc. **82** (1976), 907-909.
- [6] W. Szlenk, *Sur les suites faiblement convergentes dans l'espace L* , Studia Math. **25** (1966), 337-341.
- [7] I. Tweddle, *Weak compactness in locally convex spaces*, Glasgow Math. J. **9** (1968), 123-127.
- [8] _____, *Vector-valued measures*, Proc. London Math. Soc. **20** (1970), 469-489.

DEPARTMENT OF MATHEMATICAL SCIENCES, LAKEHEAD UNIVERSITY, THUNDER BAY, ONTARIO, CANADA