

FINITE SIMPLE GROUPS CONTAINING A SELF-CENTRALIZING ELEMENT OF ORDER 6

JOHN L. HAYDEN AND DAVID L. WINTER

ABSTRACT. By a self-centralizing element of a group we mean an element which commutes only with its powers. In this paper we establish the following result:

THEOREM. *Let G be a finite simple group which has a self-centralizing element of order 6. Assume that G has only one class of involutions. Then G is isomorphic to one of the groups M_{11} , J_1 , $L_3(3)$, $L_2(11)$, $L_2(13)$.*

By a self-centralizing (s.c.) element of a group we mean an element which commutes only with its own powers. The structure of a finite group containing a s.c. element of order 2 has been known for a long time. The classification of finite groups with a s.c. element of order 3 was carried out by Feit and Thompson [2] and is an important tool in this paper. The determination of finite simple groups containing a s.c. element of prime order $p > 3$ is a classical problem as yet unsolved although some special cases have been handled. In [12], M. Suzuki proved that the only finite simple groups containing a s.c. element of order 4 are $L_2(7)$, A_6 and A_7 . Finite simple groups containing a s.c. element of order 8 are studied in [10]. In this paper we establish the following result.

THEOREM. *Let G be a finite simple group which has a self-centralizing element of order 6. Assume that G has only one class of involutions. Then G is isomorphic to one of the groups M_{11} , J_1 , $L_3(3)$, $L_2(11)$, $L_2(13)$.*

The alternating groups A_8 , A_9 are examples of simple groups with a s.c. element of order 6 which has two classes of involutions. As indicated in [8], it may be very difficult to handle groups containing a s.c. element of order 6 having more than one class of involutions.

The smallest Ree group is an example of a nonsimple group containing a s.c. element of order 6 and having only one class of involutions. It is isomorphic to an extension of $L_2(2^3)$ by a field automorphism of order 3.

We shall make use of the following theorem.

(1.1) ([2]). *If Y is a finite group with a s.c. element of order 3, then Y has a normal subgroup N such that one of the following holds:*

- (i) Y/N is cyclic of order 3 and N is nilpotent.
- (ii) Y/N is dihedral of order 6 and N is nilpotent.

Received by the editors December 6, 1976 and, in revised form, March 25, 1977.
AMS (MOS) subject classifications (1970). Primary 20D05.

© American Mathematical Society 1977

- (iii) $Y/N \cong A_5$ and N is an elementary abelian 2-group.
 (iv) $Y/N \cong L_2(7)$ and $N = 1$.

From now on we shall let G denote a group satisfying the hypotheses of our theorem. Then G contains an element x of order 3 and an involution t such that $[t, x] = 1$ and $C(tx) = \langle tx \rangle$. Furthermore every involution of G is conjugate to t .

(1.2) $C(x) = \langle t \rangle L$ where L is a nilpotent normal subgroup of $C(x)$ of odd order. $L/\langle x \rangle$ is abelian and is inverted by t .

PROOF. Let $\bar{C} = C(x)/\langle x \rangle$ and let $\bar{v} \in \bar{C}$ be such that $[\bar{v}, \bar{t}] = \bar{1}$. Then $[\bar{v}, \bar{t}] = \bar{1}$ and so $[v, t] = x_0 \in \langle x \rangle$. Hence $v^{-1}t = x_0 t$. Since t has order 2, $x_0 = 1$ and $v \in C(t) \cap C(x) = \langle xt \rangle$. Hence $\bar{v} \in \langle \bar{t} \rangle$. Thus \bar{t} is a s.c. element of \bar{C} of order 2. The structure of \bar{C} is therefore known and (1.2) follows easily.

(1.3) Let H be a subgroup of G with $tx \in H$. Assume that H contains a normal 2-subgroup U with $t \in U$. Then $C_H(x) = \langle tx \rangle$. Also if $\bar{H} = H/U$, then \bar{x} is a s.c. element of \bar{H} of order 3.

PROOF. By (1.2), $U \cap C(x) = \langle t \rangle \triangleleft C_H(x)$. But by (1.2), $N_{C(x)}(\langle t \rangle) = \langle tx \rangle$. Hence $C_H(x) = \langle tx \rangle$. This implies that $\langle x \rangle$ is a Sylow 3-subgroup of H . It is well known and easily verified that $N(\langle x \rangle) = N_{\bar{H}}(\langle \bar{x} \rangle)$. It is also easily checked from this that $\bar{y} \in \bar{H}$ centralizes \bar{x} if and only if $y \in C_H(x)U$. Therefore $C_{\bar{H}}(\bar{x}) = \langle tx \rangle U = \langle \bar{x} \rangle$.

By setting $H = C(t)$, $U = \langle t \rangle$, we have as an immediate consequence of (1.1) and (1.3)

(1.4) $C(t)$ contains a normal subgroup $N \geq \langle t \rangle$ such that one of the following holds:

- (i) $C(t)/N$ is cyclic of order 3 and $N/\langle t \rangle$ is nilpotent.
 (ii) $C(t)/N$ is dihedral of order 6 and N is nilpotent.
 (iii) $C(t)/N \cong A_5$ and $N/\langle t \rangle$ is an elementary abelian 2-group.
 (iv) $C(t)/N \cong L_2(7)$ and $N = \langle t \rangle$.

The remainder of the proof of the theorem is divided into the four cases determined by (1.4). Assume first that (1.4) (iv) holds. Then $C(t)/\langle t \rangle \cong L_2(7)$. It has been proved by Schur [11, p. 120], that $C(t)$ is either isomorphic to a direct product of $\langle t \rangle$ and $L_2(7)$ or to $SL(2, 7)$. By a result of Janko and Thompson [9], the first case is not possible. In the second case $C(t)$, and hence G , has a quaternion Sylow 2-subgroup. By a well-known result of Brauer and Suzuki, G cannot be simple. Therefore (1.4) (iv) cannot hold.

Assume next that (1.4) (iii) holds with $N = \langle t \rangle$. The argument of the preceding paragraph shows $C(t) = \langle t \rangle \times K$, $K \cong A_5$. Therefore, by [7] G is isomorphic to J_1 .

In the remaining cases $C(t)$ is solvable or $C(t)/N \cong A_5$. In the latter case $|N| > 2$ with $C(N) \subseteq N$. Hence $C(t)$ is 2-constrained in all cases. As G has

only one class of involutions, a theorem of Gorenstein and Goldschmidt [3, p. 74] implies $O(C(t)) = 1$ or $SCN_3(2) = \emptyset$. In the latter case, Corollary 4, [5] shows G is isomorphic to one of the groups $L_2(q)$, $L_3(q)$, $U_3(q)$, q odd, $U_3(4)$, A_7 or M_{11} . The only groups among these satisfying our hypotheses are $L_3(3)$, M_{11} , $L_2(11)$ and $L_2(13)$.

We may now assume $O(C(t)) = 1$. This forces N to be a 2-group and $|C(t)| = 2^a \cdot 3 \cdot 5^b$ for some a, b . We now claim that if M is a 2-local subgroup of G , then each Sylow subgroup of M of odd order is cyclic. Without loss we may assume $t \in Z(O_2(M)) = Z$. Suppose M contains an elementary abelian p -subgroup P of order p^2 , p odd. By a well-known result [4, p. 188], $Z = \prod_{y \in P^*} C_Z(y)$. Again without loss, we can assume $t \in C_Z(y)$ for some $y \in P^*$. Then $y \in C(t)$, $p \in \{3, 5\}$. Now let $y_1 \in P - \langle y \rangle$. Since $Z \triangleleft M$, $t^{y_1} \in Z \cap C(t)$. Also, $(t^{y_1})^{y_1} = (t^y)^{y_1} = t^{y_1}$ so $t^{y_1} \in C(t) \cap C(y)$. If $p = 3$, $|y| = 3$ and, as $\langle ty \rangle$ is conjugate to $\langle tx \rangle$ in $C(t)$, $C(t) \cap C(y) = \langle ty \rangle$. If $p = 5$, we are in case (1.4)(iii). Let $\bar{C} = C(t)/\langle t \rangle$. As \bar{x} is a s.c. element of order 3 in \bar{C} , Theorem 8.2 of [6] implies $\bar{N} = N/\langle t \rangle$ is the direct sum of minimal normal subgroups of \bar{C} of order 16 on which $A_5 \cong SL(2, 4)$ acts in the natural way. Consequently \bar{y} is a s.c. element of order 5 in \bar{C} and $C(t) \cap C(y) = \langle ty \rangle$ in this case as well. In all cases $t^{y_1} \in C(t) \cap C(y) = \langle ty \rangle$ so $t^{y_1} = t$. Hence $y_1 \in C(ty) = \langle ty \rangle$ and $y_1 \in \langle y \rangle$, a contradiction proving the claim. A recent result of M. Aschbacher [1] now gives the possibilities for G . We see that there is no additional group satisfying our conditions. This completes the proof of the theorem.

REFERENCES

1. M. Aschbacher, *Thin finite simple groups*, Bull. Amer. Math. Soc. **82** (1976), 484.
2. W. Feit and J. G. Thompson, *Finite groups which contain a self-centralizing subgroup of order 3*, Nagoya Math. J. **21** (1962), 185–197.
3. D. Gorenstein, *Centralizers of involution of finite simple groups*, Finite Simple Groups, Academic Press, London, 1971.
4. ———, *Finite groups*, Harper and Row, New York, 1968.
5. D. Gorenstein and K. Harada, *Finite groups whose 2-subgroups are generated by at most 4 elements*, Mem. Amer. Math. Soc. No. 147, 1974.
6. G. Higman, *Odd characterizations of finite simple groups*, (Lecture notes, University of Michigan, 1968).
7. Z. Janko, *A new finite simple group with Abelian Sylow 2-subgroups and its characterization*, J. Algebra **3** (1966), 147–186.
8. ———, *A class of simple groups of characteristic 2*, Finite Groups' 72, North-Holland, New York, 1973.
9. Z. Janko and J. G. Thompson, *On a class of finite simple groups of Ree*, J. Algebra **4** (1966), 274–292.
10. L. Schiefelbusch, *Sylow 2-subgroups of simple groups*, J. Algebra **31** (1974), 131–153.
11. I. Schur, *Untersuchungen über die Darstellung der endlichen Gruppen durch gebrochene lineare Substitutionen*, J. für die r. und a. Math. **132** (1907), 85–137.
12. M. Suzuki, *On finite groups containing an element of order four which commutes only with its powers*, Illinois J. Math. **3** (1959), 255–271.

DEPARTMENT OF MATHEMATICS, BOWLING GREEN STATE UNIVERSITY, BOWLING GREEN, OHIO 43403

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN 48824