

## MARKOV PARTITIONS ARE NOT SMOOTH<sup>1</sup>

RUFUS BOWEN

**ABSTRACT.** The boundaries of the sets in a Markov partition for linear Anosov diffeomorphisms of  $T^3$  cannot be smooth.

It will be shown that the boundaries of the rectangles in a Markov partition for an Anosov automorphism of the 3-torus are never smooth. The proof constructs a certain one-dimensional invariant set and then appeals to the result of Franks [2] that such a set cannot contain a smooth arc. The existence of such an invariant set was asked by Smale (see Hirsch [3]); S. G. Hancock has also constructed such a set [7].

Let  $A$  be a  $3 \times 3$  integral matrix with eigenvalues  $\lambda_1, \lambda_2, \lambda_3$  such that

$$\det A = \lambda_1 \lambda_2 \lambda_3 = \pm 1 \text{ and each } |\lambda_i| \neq 1.$$

For definiteness we suppose  $|\lambda_1| < 1, |\lambda_2|, |\lambda_3| > 1$ . Let  $V^s$  be the line in  $R^3$  spanned by the eigenvector for  $\lambda_1$  and  $V^u$  the invariant plane corresponding to  $\lambda_2$  and  $\lambda_3$ ; so  $R^3 = V^s \oplus V^u$ . A compact set of the form  $R = A \oplus B$  is called a *rectangle* provided  $R = \overline{\text{int } R}$ . One writes  $\partial^s R = A \oplus \partial B$  and  $\partial^u R = \partial A \oplus B$ . A rectangle in  $T^3$  is the projection of a rectangle in  $R^3$  of small diameter.

The matrix  $A$  induces a diffeomorphism  $f$  on  $T^s = R^3/Z^3$  by  $f(x + Z^3) = Ax + Z^3$ . A *Markov partition* for  $f$  is a family  $\mathcal{C} = \{R_1, \dots, R_m\}$  of small connected rectangles in  $T^3$  covering  $T^3$  and satisfying

- (i)  $\text{int}(R_i \cap R_j) = \emptyset$  for  $i \neq j$ ;
- (ii)  $f(\partial^s \mathcal{C}) \subset \partial^s \mathcal{C}$  where  $\partial^s \mathcal{C} = \bigcup_{i=1}^m \partial^s R_i$ ;
- (iii)  $f^{-1}(\partial^u \mathcal{C}) \subset \partial^u \mathcal{C}$  where  $\partial^u \mathcal{C} = \bigcup_{i=1}^m \partial^u R_i$ .

The conditions  $|\lambda_i| \neq 1$  means that  $f$  is an Anosov diffeomorphism and hence has Markov partitions (Sinai [6]).

**PROPOSITION.**  $X = \bigcap_{k=0}^{\infty} f^k(\partial^s \mathcal{C})$  is a compact one-dimensional invariant set for  $f$ .

**PROOF.** That  $f(X) = X$  follows from (ii) above. Because  $X \subset \partial^s \mathcal{C}$  is nowhere dense in  $T^3$ ,  $\dim X \leq 2$  and then  $\dim X \leq 1$  by a result of Hirsch and Williams [3].

Next one finds a periodic point  $p \in \partial^s \mathcal{C}$ . This is accomplished by counting the number  $N_n(f)$  of fixed points of  $f^n$  in two different ways. For convenience

---

Received by the editors August 26, 1977 and, in revised form, January 13, 1978.

AMS (MOS) subject classifications (1970). Primary 58F15.

<sup>1</sup>Research partially supported by NSF grant MCS74-19388.A01.

assume  $\lambda_1, \lambda_2, \lambda_3$  are all positive. From algebraic topology

$$N_n(f) = |\det(I - A^n)| = |-\lambda_1^n - \lambda_2^n - \lambda_3^n + (\lambda_1\lambda_2)^n + (\lambda_1\lambda_3)^n + (\lambda_2\lambda_3)^n|.$$

On the other hand one estimates  $N_n(f)$  using symbolic dynamics (Manning [5] has an exact count). There is a subshift of finite type  $\sigma: \Sigma_B \rightarrow \Sigma_B$  and an at most  $c$ -to-1 ( $c < \infty$ ) surjection [1]  $\pi: \Sigma_B \rightarrow T^3$  so that  $\pi \circ \sigma = f \circ \pi$ . The map  $\pi$  is 1-1 over points  $x \in T^3 \setminus \bigcup_{k \in \mathbb{Z}} f^k(\partial^s \mathcal{C} \cup \partial^u \mathcal{C})$ . It follows that

$$|N_n(f) - N_n(\sigma|_{\Sigma_B})| \leq c \cdot \text{card}\{x \in \partial^s \mathcal{C} \cup \partial^u \mathcal{C}: f^n x = x\}.$$

The local unstable manifold  $W_\varepsilon^u(x)$  is defined by

$$W_\varepsilon^u(x) = x + \{v \in V^u: \|v\| < \varepsilon\}.$$

Because  $s = 1$  and  $R_k$  is connected, each

$$\partial^u R_k = (W_\varepsilon^u(x_{1,k}) \cap R_k) \cup (W_\varepsilon^u(x_{2,k}) \cap R_k)$$

for certain  $x_{1,k}, x_{2,k} \in R_k$ . The Markov condition  $f^{-1}(\partial^u \mathcal{C}) \subset \partial^u \mathcal{C}$  implies that  $f^{-1}$  maps each of the sets  $\{W^u(x_{j,k}, R_k)\}_{j,k}$  into another one, by a contraction. It follows that  $f^{-1}$  (and hence  $f$ ) has only finitely many periodic points in  $\partial^u \mathcal{C}$ .

We shall prove that  $\partial^s \mathcal{C}$  contains infinitely many periodic points. Otherwise  $|N_n(f) - N_n(\sigma|_{\Sigma_B})|$  is bounded. Now

$$N_n(\sigma|_{\Sigma_B}) = \text{Tr}(B^n) = \mu_1^n + \dots + \mu_r^n$$

where  $\mu_1, \dots, \mu_r$  are the eigenvalues of  $B$ . Since  $|\lambda_1|, |\lambda_1\lambda_2| = |\lambda_3|^{-1}$  and  $|\lambda_1\lambda_3| = |\lambda_2|^{-1}$  are all less than 1, one would have (recall  $\lambda_2\lambda_3 > 0$  for convenience) that

$$c_n = \mu_1^n + \dots + \mu_r^n + \lambda_2^n + \lambda_3^n - (\lambda_2\lambda_3)^n$$

is bounded. Now the entropy of  $f$  equals

$$h(f) = \log(\lambda_2\lambda_3) \quad (\text{see [8]});$$

also  $h(f) = h(\sigma|_{\Sigma_B}) = \log \mu$  where  $\mu$  is the largest positive eigenvalue of  $B$ , say  $\mu = \mu_r$ . Then  $\lambda_2\lambda_3 = \mu_r$  and so

$$c_n = \mu_1^n + \dots + \mu_{r-1}^n + \lambda_2^n + \lambda_3^n.$$

The function

$$\zeta(z) = \exp \sum_{n=1}^{\infty} \frac{c_n z^n}{n} = \frac{1}{(1 - \lambda_2 z)(1 - \lambda_3 z)(1 - \mu_1 z) \cdots (1 - \mu_{r-1} z)}$$

is analytic on  $|z| < 1$  ( $c_n$  bounded) but has a pole at  $z = \lambda_2^{-1}$ , a contradiction.

Now let  $p \in \partial^s \mathcal{C}$  with  $f^n p = p$ . In [1] certain finite strings of integers  $J_s(x)^*$  and  $J_u(x)^*$  were given which described the position of  $x$  relative to  $\mathcal{C}$  and its boundaries. These strings of integers are constant over the minimal set  $\{p, f(p), \dots, f^{n-1}(p)\}$  [1, p. 916]. From this it follows that (see [1, pp. 910-912]) there are neighborhoods  $U_k$  of  $f^k p$  so that

$$y \in W_\varepsilon^u(f^k p) \cap \partial^s \mathcal{C} \cap U_k \Rightarrow f^{-1} y \in \partial^s \mathcal{C}.$$

Since  $f^{-1} W_\varepsilon^u(f^k p) \subset W_\varepsilon^u(f^{k-1} p)$ , it follows that there is a neighborhood

$U \subset U_0$  of  $p$  so that  $f^{-i}y \in W_\varepsilon^u(f^{-i}p) \cap \partial^s \mathcal{C} \cap U_{-i}$  for every  $i \geq 0$  when  $y \in W_\varepsilon^u(p) \cap \partial^s \mathcal{C} \cap U$ . Hence  $W_\varepsilon^u(p) \cap \partial^s \mathcal{C} \cap U \subset \bigcap_{i=0}^{\infty} f^i(\partial^s \mathcal{C}) = X$ . Now  $W_\varepsilon^u(p) \cap U$  contains a small 2-disk  $D$  about  $p$  in  $W_\varepsilon^u(p)$ . The sets  $D \cap R_k$  ( $k = 1, \dots, m$ ) generate a partition of  $D$  into closed sets with dense interior intersecting only in their boundaries and whose boundaries lie in  $\partial^s \mathcal{C} \cap W_\varepsilon^u(p)$ . There are at least two sets since  $p \in \partial^s \mathcal{C}$ ; thus  $\partial^s \mathcal{C} \cap W_\varepsilon^u(p)$  has dimension 1 since it separates open sets in  $R^2$  [4, p. 46]. Hence  $\dim X = 1$ .  $\square$

**THEOREM.**  $\mathcal{C}$  is not a smooth partition.

**PROOF.** Smoothness here means that the boundaries of the  $R_k$ 's are piecewise smooth submanifolds (with corners). If  $\mathcal{C}$  were smooth, then  $\partial^s \mathcal{C} \cap W_\varepsilon^u(p)$  in the above proof would piecewise be a smooth 1-manifold. But a theorem of Franks [2] says that a one-dimensional invariant subset for  $f$  cannot contain a  $C^1$  arc.  $\square$

#### REFERENCES

1. R. Bowen, *Markov partitions and minimal sets for Axiom A diffeomorphisms*, Amer. J. Math. **92** (1970), 907–918.
2. J. Franks, *Invariant sets of hyperbolic toral automorphisms*, Amer. J. Math. **99** (1977), 1089–1095.
3. M. Hirsch, *On invariant subsets of hyperbolic sets*, Essays in Topology and Related Topics, Springer, Berlin, 1970.
4. W. Hurewicz and H. Wallman, *Dimension theory*, Princeton Univ. Press, Princeton, N.J., 1948.
5. A. Manning, *Axiom A diffeomorphisms have rational zeta functions*, Bull. London Math. Soc. **3** (1971), 215–220.
6. Ya. Sinai, *Construction of Markov partitions*, Functional Anal. Appl. **2** (1968), 70–80.
7. S. G. Hancock, *Orbits and paths under hyperbolic toral automorphisms* (to appear).
8. R. Bowen, *Entropy for group endomorphisms and homogeneous spaces*, Trans. Amer. Math. Soc. **153** (1971), 401–414; erratum, *ibid.* **181** (1973), 509–510.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT BERKELEY, BERKELEY, CALIFORNIA 94720