

APPROXIMATE UNITS IN IDEALS OF GROUP ALGEBRAS

HARALD RINDLER

ABSTRACT. Several properties of locally compact groups can be characterized by the existence of certain approximate units in 3 ideals of the group algebra. We combine methods and results from [8], [7], [5].

Let G be a locally compact group, $M(G)$ the Banach algebra of all bounded complex measures, $L^1(G)$ the ideal consisting of all absolute-continuous measures with respect to left Haar-measure,

$$M^0(G) = \left\{ m \in M(G), \int 1 \, dm = 0 \right\}, \quad L^0(G) = M^0(G) \cap L^1(G).$$

Let B be a Banach algebra. We consider the following properties: (U) B has a unit, (A) there exists a bounded net u_α in B such that $\lim_\alpha u_\alpha * f = f$ for every $f \in B$, (A_c) in addition $u_\alpha \in Z(B)$ the center of B , (A_d) $(u_\alpha * f - f) \in Z(B)$ for every $f \in B$.

We consider the following classes of locally compact groups: [SIN]: G has a basis of neighbourhoods of 1 invariant under all inner automorphisms, [FC]⁻: every conjugacy class in G is precompact, [FIA]⁻: the group of inner automorphisms is precompact in the group of all continuous automorphisms, [FD]⁻, resp. FD: the group generated by all commutators $(xyx^{-1}y^{-1})$ is precompact, resp. finite (see [3]). G is amenable if there exists a left invariant mean on $L^\infty(G)$, the space of all measurable essentially bounded functions (see [2] or [8]).

It is well known that $L^1(G)$ always satisfies (A), and (U) iff G is discrete. Mosak, [7], proved that $L^1(G)$ satisfies (A_c) iff $G \in$ [SIN]. Reiter, [9], proved that $L^0(G)$ satisfies (A) iff G is amenable and Kotzmann, Rindler, [5], that $L^0(G)$ satisfies (A_c) iff $G \in$ [FIA]⁻.

Using their methods and results and well-known structure theorems we easily can complete the results above to the following list of characterizations:

Let G be a locally compact group then

- | | | | |
|----|---------------|-----|--------------------------------|
| 1. | G is discrete | iff | L ¹ (G) satisfies U |
| 2. | G is compact | iff | M ⁰ (G) satisfies U |
| 3. | G is finite | iff | L ⁰ (G) satisfies U |

Received by the editors September 1, 1977.

AMS (MOS) subject classifications (1970). Primary 43A20, 22D05, 22D15.

Key words and phrases. Locally compact group, group algebra, approximate unit, amenable group, compactness conditions.

© American Mathematical Society 1978

- | | | | |
|-----|----------------------------|------|----------------------------|
| 4. | G is arbitrary l.c. | then | $L^1(G)$ satisfies A |
| 5. | G is amenable | iff | $M^0(G)$ satisfies A |
| 6. | G is amenable | iff | $L^0(G)$ satisfies A |
| 7. | $G \in [\text{SIN}]$ | iff | $L^1(G)$ satisfies A_c |
| 8. | $G \in [\text{FC}]^-$ | iff | $M^0(G)$ satisfies A_c |
| 9. | $G \in [\text{FIA}]^-$ | iff | $L^0(G)$ satisfies A_c |
| 10. | G is Abelian or discrete | iff | $L^1(G)$ satisfies A_d |
| 11. | $G \in [\text{FD}]^-$ | iff | $M^0(G)$ satisfies A_d |
| 12. | G is Abelian or discrete | iff | $L^0(G)$ satisfies A_d . |
- and $\in [\text{FD}]$

REMARKS. If the boundedness assumption of the net u_α is dropped then the following remarks apply: If we only assume that for every $f \in B$ and $\varepsilon > 0$ there exists $u = u(f, \varepsilon) \in B$ $\|u * b - u\| < \varepsilon$, then parts 5 and 6 remain open. The case of compact extensions of connected groups was settled in [10, Theorem 1 and Proposition 1]. A counterexample for discrete groups would lead to a nonamenable group not containing the free group of 2 generators, B. E. Johnson, [4]. If $u = u(f, \varepsilon)$ can be taken bounded independent of f and ε then (A) holds (see Altman [1] or Wichmann [12] who also proved that the same bound can be chosen for the net u_α). For $M^0(G)$ and $L^0(G)$ the best possible bound is 2.

PROOFS. If δ is the unit of $M(G)$, λ the normalized Haar measure if G is compact (resp. finite) then $\delta - \lambda$ is a unit of $M^0(G)$ (resp. $L^0(G)$). The proof of the converse in 3 contains proofs of 1 and 2: If $u \in L^0(G)$ satisfies $f * u = f$ for every $f \in L^0(G)$ we obtain for $m = \delta - u$, $\int 1 dm = 1$ and $f * m = 0$, $f \in L^0(G)$. Choosing $f(z) = g(xz) - g(z)$, g continuous with compact support, $x \in G$ arbitrary, it follows from $0 = f * m(e) = \int f(y) dm(y^{-1})$ that m is a right Haar measure. $\int 1 dm = 1$ implies G is compact, m is absolutely continuous and so is $\delta = u + m$ and G is also discrete. If u is a unit for $M^0(G)$ then in the same way we obtain that G is compact. If u is a unit for $L^1(G)$ then $0 = f * (\delta - u)(e)$, for every f continuous with compact support implies that $u = \delta \in L^1(G)$ and we obtain that G is discrete. The converse is clear.

The proof of 6 was given in [9], 5 follows analogously. The proof of 7 is contained in [7], that of 9 in [5].

PROOF OF 8. If $\lim \|u_\alpha * m - m\| = 0$ for every $m \in M^0(G)$, u_α central, then $v_\alpha = \delta - u_\alpha$ is central and $v_\alpha * m \rightarrow 0$ for every $m \in M^0(G)$. If w_α denotes the absolutely continuous part of v_α , then w_α also belongs to the center of $M(G)$ and putting $m = g - \delta$, $g \in L^1(G)$, $\int g = 1$, we obtain $\|w_\alpha - v_\alpha\| \leq \|v_\alpha * (g - \delta)\| \rightarrow 0$ ($v_\alpha * g \in L^1(G)$). This implies $\lim_\alpha \|w_\alpha * m\| = 0$ for every $m \in M^0(G)$, therefore $G \in [\text{FC}]^-$ by [5, Proposition, (c)].

If conversely $G \in [\text{FC}]^-$ then by Robertson's structure theorem ([11], for a proof see [6]) there exists a compact normal subgroup K such that $G/K = V \times D$, V a vector group, D a discrete [FC]-group. This reduces the problem to discrete groups, but then $M^0(G) = L^0(G)$ and the result follows from 9.

To prove the nontrivial part of 10 we assume that $(\delta - u_\alpha) * f$ is central for every $f \in L^1(G)$. Choosing f_β the normalized characteristic functions of U_β , a basis of neighbourhoods of $x \in G$, we obtain that $(\delta - u_\alpha) * \delta_x$ is central for every $x \in G$ (the center being weakly closed). The discrete part of this measure is also central and equal to δ_x if G is not discrete and G is necessarily Abelian.

If $G \in [\text{FD}]^-$ then there exists a compact normal subgroup K such that G/K is Abelian. We obtain a suitable approximate unit for $M^0(G)$ by lifting from $M^0(G/K)$. Conversely we have $(\delta - u_\alpha) * m$ is central for every $m \in M^0(G)$. For a fixed α_0 we put $\mu = \delta - u_{\alpha_0}$. We want to show that $\mu * m$ is central for every $m \in M(G)$. We may assume that $\int 1 dm = 1$, then $(m - u_\alpha) \in M^0(G)$ and $\mu * (m - u_\alpha)$ is central for every α . Furthermore $\lim_\alpha \mu * u_\alpha = \lim_\alpha u_\alpha * \mu = 0$. The center being closed this implies that $\mu * m$ and in particular $\mu * \delta_x$ is central for every $x \in G$. This implies that $\mu = \mu * \delta_{yx^{-1}y^{-1}}$ and $\mu = \mu * \delta_z$ for every $z \in K$ the closure of the commutator subgroup. μ being a bounded measure, this implies that G is compact.

PROOF OF 12. As before, we obtain that $(\delta - u_\alpha) * \delta_x$ is central for all $x \in G$. By 10 we may assume that G is discrete and by 11 G is a [FC]-group. The converse is clear.

REFERENCES

1. M. Altman, *Contracteurs dans les algèbres de Banach*, C. R. Acad. Sci. Paris Sér. A-B **274** (1971), A399–A400. MR **45** #2437.
2. P. F. Greenleaf, *Invariant means of topological groups*, Van Nostrand, New York, 1969. MR **40** #4776.
3. S. Grosser and M. Moskowitz, *Compactness conditions in topological groups*, J. Reine Angew. Math. **246** (1971), 1–40. MR **44** #1766.
4. B. E. Johnson, *Some examples in harmonic analysis*, Studia Math. **48** (1973), 182–188. MR **49** #3456.
5. E. Kozmann and H. Rindler, *Central approximate units in a certain ideal of $L^1(G)$* , Proc. Amer. Math. Soc. **57** (1976), 155–158. MR **53** #8784.
6. J. Liukkonen, *Dual spaces of groups with precompact conjugacy classes*, Trans. Amer. Math. Soc. **180** (1973), 85–108. MR **47** #6937.
7. R. D. Mosak, *Central functions in group algebras*, Proc. Amer. Math. Soc. **29** (1971), 613–616. MR **43** #5323.
8. H. Reiter, *Classical harmonic analysis and locally compact groups*, Clarendon Press, Oxford, 1968. MR **46** #5933.
9. ———, *Sur certain idéaux dans $L^1(G)$* , C. R. Acad. Sci. Paris Sér. A-B **267** (1968), A882–A885. MR **39** #6025.
10. H. Rindler, *Zur Eigenschaft P_1 lokalkompakter Gruppen*, Indag. Math. **35** (1973), 142–147. MR **48** #11921.
11. L. C. Robertson, *A note on the structure of Moore groups*, Bull. Amer. Math. Soc. **75** (1969), 594–599. MR **39** #7027.
12. J. Wichmann, *Bounded approximate units and bounded approximate identities*, Proc. Amer. Math. Soc. **41** (1973), 547–550. MR **48** #2767.

MATHEMATICS INSTITUTE, UNIVERSITY OF VIENNA, A-1090 VIENNA, AUSTRIA