

## ON ROBINSON'S $\frac{1}{2}$ CONJECTURE

ROGER W. BARNARD

**ABSTRACT.** In 1947, R. Robinson conjectured that if  $f$  is in  $S$ , i.e. a normalized univalent function on the unit disk, then the radius of univalence of  $[zf(z)]/2$  is at least  $\frac{1}{2}$ . He proved in that paper that it was at least .38. The conjecture has been shown to be true for most of the known subclasses of  $S$ . This author shows through use of the Grunski inequalities, that the minimum lower bound over the class  $S$  lies between .49 and .5.

**Introduction.** Let  $\mathcal{A}$  denote the class of analytic functions on the unit disk  $U = \{z: |z| < 1\}$ . Let  $S$  denote the univalent functions  $f$  in  $\mathcal{A}$  normalized by  $f(0) = 1 - f'(0) = 0$ . Denote by  $K$ ,  $S^*$ ,  $C$ , and  $Sp$  the standard subclasses of  $S$  consisting of functions that are convex starlike, close to convex and spirallike respectively. For a subclass  $X$  (possibly a singleton) of  $\mathcal{A}$  let  $r_S(X)$  denote the minimum radius of univalence over all functions  $f$  in  $X$ . We use corresponding notation for the other subclasses of  $S$ . For example  $r_{S^*}(X)$  denotes the minimum radius of starlikeness over all functions  $f$  in  $X$ .

For a function  $f$  in  $S$  define the operator  $\Gamma: S \rightarrow \mathcal{A}$  by  $\Gamma f = (zf) \frac{1}{2}$ . In 1947 R. Robinson [10] considered the problem of determining  $r_S[\Gamma(S)]$ . Robinson observed that for each  $f$  in  $S$ ,  $[\Gamma(f)]' \neq 0$  for  $|z| < \frac{1}{2}$ . He also noted that for the Koebe function  $k$ ,  $k(z) = z(1 - z)^{-2}$ ,  $r_S(k) = r_{S^*}(k) = \frac{1}{2}$ , which implies  $r_S[\Gamma(S)] \leq \frac{1}{2}$ . He in fact conjectured that  $r_S[\Gamma(S)] = \frac{1}{2}$ . He was able to show that  $r_{S^*}[\Gamma(S)] > .38$ .

There have been a number of papers (e.g. [2], [3], [6], [7], [8]) on the connection between the operator  $\Gamma$  and various subclasses of  $S$ . In these papers it has been shown that

$$r_K[\Gamma(K)] = r_{S^*}[\Gamma(S^*)] = r_C[\Gamma(C)] = r_{Sp}[\Gamma(Sp)] = \frac{1}{2}$$

and that  $\Gamma$  preserves Rogosinski's class of typically real functions (not necessarily univalent) up to  $|z| < \frac{1}{2}$ . It was observed in [2] that with the exception of the result  $r_{Sp}[\Gamma(Sp)] = \frac{1}{2}$  these results follow directly from the S. Ruscheweyh-T. Sheil-Small theory [11]. They proved that, except for  $Sp$ , convolution by convex functions preserves the above subclasses of  $S$ . In order to obtain the related results in [2] one need only observe that for  $f(z) = \sum a_n z^n$ ,

$$\Gamma[f(z)] = h * f(z) = \sum [(n + 1)/2] z^n * f(z) = \sum [(n + 1)/2] a_n z^n$$

---

Presented to the Society, January 4, 1978; received by the editors November 11, 1977.

AMS (MOS) subject classifications (1970). Primary 30A32.

Key words and phrases. Univalent functions, Grunsky's inequalities.

© American Mathematical Society 1978

and that  $h(z) = (z - z^2/2)(1 - z)^{-2}$  is convex for  $|z| < \frac{1}{2}$ . As was shown in [2] most of the results that had been obtained on generalizations of the operator  $\Gamma$  on subclasses of  $S$  can also be obtained in a similar manner by the appropriate modifications of  $h$ . However, for the entire class  $S$ , if we let  $r_0 = r_C(S) \approx .80$  from [5], it appears that the easily obtained lower bound for  $r_S[\Gamma(S)]$  of  $r_0/2 \approx .41$  is the most that can be obtained from the convolution operator method. It does show that  $.41 < r_S[\Gamma(S)] \leq \frac{1}{2}$ . In the present note the author, through use of the Grunsky inequalities, is able to prove that  $.49 < r_S[\Gamma(S)] \leq \frac{1}{2}$ .

**PROOF OF MAIN RESULT.** To find a lower bound for  $r_S[\Gamma(S)]$  we consider the nonvanishing of

$$\frac{f(z) + zf'(z) - f(\zeta) - \zeta f'(\zeta)}{z - \zeta}.$$

By use of the minimum principle we may assume  $|z| = |\zeta| < r$ . Since  $f$  is in  $S$  we may divide through by  $[f(z) - f(\zeta)]/(z - \zeta)$ . Thus it suffices to find the largest  $r$  such that

$$1 + \frac{zf'(z) - \zeta f'(\zeta)}{f(z) - f(\zeta)} \neq 0, \quad |z| = |\zeta| < r. \tag{1}$$

Consider for  $f$  in  $S$  the Grunsky coefficients defined by letting

$$\log \frac{f(z) - f(\zeta)}{z - \zeta} = \sum_{n,m=0}^{\infty} d_{nm} z^n \zeta^m. \tag{2}$$

Putting  $\zeta, z = 0$  respectively, in (2) we obtain

$$\log \frac{f(z)}{z} = \sum_{n=0}^{\infty} d_{n0} z^n, \quad \log \frac{f(\zeta)}{\zeta} = \sum_{m=0}^{\infty} d_{0m} \zeta^m.$$

Hence

$$\log \frac{f(z) - f(\zeta)}{z - \zeta} = \log \frac{f(z)}{z} + \log \frac{f(\zeta)}{\zeta} + \sum_{n,m=1}^{\infty} d_{nm} z^n \zeta^m. \tag{3}$$

Although Grunsky's inequalities are usually stated in terms of the function  $F$ , on  $|\xi| > 1$  defined by  $F(\xi) = 1/f(1/\xi)$ , it is more convenient for our purposes to express them directly in terms of  $f$  in  $S$ . To do this, we observe, by letting  $z' = 1/z, \zeta' = 1/\zeta$ , that

$$\begin{aligned} & \log \frac{f(z) - f(\zeta)}{z - \zeta} - \log \frac{f(z)}{z} - \log \frac{f(\zeta)}{\zeta} \\ &= \log \frac{f(1/z') - f(1/\zeta')}{z - \zeta} \\ &= \log \frac{f(1/z') - f(1/\zeta')}{f(1/z')f(1/\zeta')z'\zeta'[(1/z') - (1/\zeta')]} \\ &= \log \frac{1/f(1/z') - 1/f(1/\zeta')}{z' - \zeta'} = \log \frac{F(z') - F(\zeta')}{z' - \zeta'} \\ &= \sum_{n,m=1}^{\infty} d_{nm} (z')^{-n} (\zeta')^{-m} = \sum_{n,m=1}^{\infty} d_{nm} z^n \zeta^m. \end{aligned}$$

Thus we can use the following form of Grunsky's inequalities (see Pommerenke [9, p. 60]). For  $f$  in  $S$  and  $d_{nm}$  defined by (2) we have for arbitrary complex  $x_n$ ,

$$\sum_{n=1}^{\infty} n \left| \sum_{m=1}^{\infty} d_{nm} x_m \right|^2 < \sum_{n=1}^{\infty} \frac{|x_n|^2}{n} \tag{4}$$

provided the last series converges. Now, differentiating (3) with respect to  $z$  and  $\zeta$  we see from the uniform convergence of the series in (3) for  $|z| = |\zeta| < r < 1$  that

$$\frac{zf'(z)}{f(z) - f(\zeta)} - \frac{z}{z - \zeta} = \frac{zf'(z)}{f(z)} - 1 + \sum_{m,n=1}^{\infty} nd_{nm}z^n\zeta^m,$$

and

$$\frac{-\zeta f'(\zeta)}{f(z) - f(\zeta)} + \frac{\zeta}{z - \zeta} = \frac{\zeta f'(\zeta)}{f(\zeta)} - 1 + \sum_{n,m=1}^{\infty} md_{nm}z^n\zeta^m.$$

Adding these two expressions and rearranging we obtain

$$1 + \frac{zf'(z) - \zeta f'(\zeta)}{f(z) - f(\zeta)} = \frac{zf'(z)}{f(z)} + \frac{\zeta f'(\zeta)}{f(\zeta)} + \sum_{n,m=1}^{\infty} (n + m)d_{nm}z^n\zeta^m. \tag{5}$$

Thus from (1) we need to find the largest  $r$  for which the right-hand side of (5), which we denote by  $T(z, \zeta)$ , does not vanish for  $|z| = |\zeta| < r$ . We have by the use of Schwarz's inequality and (4) that

$$\begin{aligned} \Re e\{T(z, \zeta)\} &\geq \Re e\left\{\frac{zf'(z)}{f(z)} + \frac{\zeta f'(\zeta)}{f(\zeta)}\right\} - \left|\sum_{n,m=1}^{\infty} (n + m)d_{nm}z^n\zeta^m\right| \\ &\geq 2 \min_{|z|=r} \Re e\left\{\frac{zf'(z)}{f(z)}\right\} - \left(\left|\sum_{n=1}^{\infty} \sqrt{n} z^n \sum_{m=1}^{\infty} \sqrt{n} d_{nm}\zeta^m\right| \right. \\ &\qquad \qquad \qquad \left. + \left|\sum_{m=1}^{\infty} \sqrt{m} \zeta^m \sum_{n=1}^{\infty} \sqrt{m} d_{mn}z^n\right|\right) \\ &\geq 2 \min_{|z|=r} \Re e\left\{\frac{zf'(z)}{f(z)}\right\} - \left(\sum_{n=1}^{\infty} nr^{2n}\right)^{1/2} \left[\sum_{n=1}^{\infty} n \left|\sum_{m=1}^{\infty} d_{nm}\zeta^m\right|^2\right]^{1/2} \\ &\qquad - \left(\sum_{m=1}^{\infty} mr^{2m}\right)^{1/2} \left[\sum_{m=1}^{\infty} m \left|\sum_{n=1}^{\infty} d_{mn}z^n\right|^2\right]^{1/2} \\ &\geq 2 \min_{|z|=r} \Re e\left\{\frac{zf'(z)}{f(z)}\right\} - \left[\frac{r^2}{(1 - r^2)^2}\right]^{1/2} \left(\sum_{n=1}^{\infty} \frac{r^{2n}}{n}\right)^{1/2} \end{aligned}$$

$$\begin{aligned}
& - \left[ \frac{r^2}{(1-r^2)^2} \right]^{1/2} \left( \sum_{m=1}^{\infty} \frac{r^{2m}}{m} \right)^{1/2} \\
& \geq 2 \min_{|z|=r} \Re e \left\{ \frac{zf'(z)}{f(z)} \right\} - 2 \left( \frac{r}{1-r^2} \right) \left( \log \frac{1}{1-r^2} \right)^{1/2}. \quad (6)
\end{aligned}$$

Since  $f$  is in  $S$  we have the well-known inequality

$$\left| \log \frac{zf'(z)}{f(z)} \right| \leq \log \frac{1+r}{1-r}, \quad |z| \leq r,$$

where, in fact, for each real  $\alpha$ , and  $z$ ,  $|z| = r < 1$ , there exists an  $f$  in  $S$  such that

$$\log [zf'(z)/f(z)] = e^{i\alpha} \log [(1+r)/(1-r)]$$

(see Jenkins [4, p. 110]). Thus, if we let  $\log [zf'(z)/f(z)] = Re^{i\Phi}$ , we can assume  $R = \log[(1+r)/(1-r)]$ . In order to find the minimum of (6) for all  $f$  in  $S$  we consider

$$\begin{aligned}
\min_{f \in S} \min_{|z|=r} \Re e \left\{ \frac{zf'(z)}{f(z)} \right\} &= \min_{\Phi} \operatorname{Re} \{ \exp(Re^{i\Phi}) \} \\
&= \min_{\Phi} [ \exp(R \cos \Phi) ] [ \cos(R \sin \Phi) ].
\end{aligned}$$

Thus, from (6), we need to find the largest  $r$  for which

$$\min_{\Phi} [ \exp(R \cos \Phi) ] [ \cos(R \sin \Phi) ] \geq \frac{r}{1-r^2} [ -\log(1-r^2) ]^{1/2}. \quad (7)$$

It is easy to see that the left-hand side of (7), call it LS, is a decreasing function of  $r$  while the right-hand side of (7), call it RS, is an increasing function of  $r$ . A computer checked calculation shows that for  $r = .490$ ,  $RS < .3379$  while  $LS > .3393$  where the minimum value occurs when  $\Phi$  is approximately 2.5 radians. We note that for  $r = .491$ ,  $RS > .3398$ . Thus, inequality (1) holds for all  $f$  in  $S$  and  $r \leq .49$ . It follows that  $r_S[\Gamma(S)] > .49$ .

#### BIBLIOGRAPHY

1. R. W. Barnard, *On the radius of starlikeness of  $(zf')$  for  $f$  univalent*, Proc. Amer. Math. Soc. **53** (1975), 385–390.
2. R. W. Barnard and C. Kellogg, *Applications of convolution operator techniques to problems in univalent function theory* (to appear).
3. S. D. Bernardi, *The radius of univalence of certain analytic functions*, Proc. Amer. Math. Soc. **24** (1970), 312–318.
4. J. Jenkins, *Univalent functions and conformal mappings*, Springer-Verlag, Berlin and New York, 1965.
5. J. Krzyz, *The radius of close-to-convexity within the family of univalent functions*, Bull. Acad. Polon. Sci. Ser. Sci. Math. **10** (1962), 201–204.
6. R. J. Libera and A. E. Livingston, *On the univalence of some classes of regular functions*, Proc. Amer. Math. Soc. **30** (1971), 327–336.

7. A. E. Livingston, *On the radius of univalence of certain analytic functions*, Proc. Amer. Math. Soc. **17** (1966), 352–357.
8. K. S. Padmanabhan, *On the radius of univalence of certain classes of analytic functions*, J. London Math. Soc. **1** (1969), 225–231.
9. C. Pommerenke, *Univalent functions*, Vandenhoeck and Ruprecht, Göttingen, 1975.
10. R. Robinson, *Univalent majorants*, Trans. Amer. Math. Soc. **61** (1947), 1–35.
11. S. Ruscheweyh and T. Sheil-Small, *Hadamard products of schlicht functions and the Polya-Schöenberg conjecture*, Comment. Math. Helv. **48** (1973).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KENTUCKY, LEXINGTON, KENTUCKY 40506

*Current address:* Department of Mathematics, Texas Tech University, Lubbock, Texas 79409