

A NOTE ON THE BOREL FORMULA

RONALD M. DOTZEL

ABSTRACT. A new proof of the Borel formula is obtained for $G = (Z_p)^r$ actions on spaces X having $H_i(X; Z_p) = 0$, $i \neq n$ (some n) and $H_n(X; Z_p) = Z_p \oplus \text{Free } Z_p G$ module. Each X^H must be a Z_p -homology $n(H)$ -sphere and then $n - n(G) = \sum(n(H) - n(G))$, sum running over corank 1 subgroups. A discussion of examples follows.

1. Introduction. If an elementary abelian p -group $G = (Z_p)^r$ acts on a Z_p -homology n -sphere, then Smith theory [3, Chapter 3] says that the fixed point set of a subgroup H of G is a Z_p -homology $n(H)$ -sphere. If p is odd $n - n(H)$ must be even for all H . The Borel formula [2, p. 175] states that, moreover, $n - n(G) = \sum(n(H) - n(G))$ where the sum runs over all subgroups H of G such that $G/H = Z_p$ (i.e. H has corank 1).

In [4] the following converse result was established.

THEOREM A. *Let X be a finite CW complex with a cellular action of $G = (Z_p)^r$, such that each X^H ($H \neq 0$) is a Z_p -homology $n(H)$ -sphere. Suppose, moreover, that there exists an n such that $\tilde{H}_i(X; Z_p) = 0$, $i \neq n$, and so that if p is odd, $n - n(G)$ is even. Also assume the Borel formula holds for this action (and n). Then $H_n(X; Z_p) = Z_p \oplus F$ where F is $Z_p G$ free.*

This result was established by constructing a linear action of G on S^n and an equivariant map $\phi: X \rightarrow S^n$ which induces Z_p -homology isomorphisms $\phi^H: X^H \rightarrow (S^n)^H$, $H \neq 0$, and a Z_p -homology epimorphism when $H = 0$.

In the present paper we establish the converse to Theorem A, which provides an extension of the Borel formula. I am indebted to Professor Glen Bredon, who first suggested the approach I used here, and to Professor Gary C. Hamrick for many helpful conversations.

2. We begin with

PROPOSITION 1. *Let $G = (Z_p)^r$ ($r > 1$) act cellularly on X , a finite CW complex such that each X^H ($H \neq 0$) is a Z_p -homology $n(H)$ -sphere. Suppose there is an integer n so that $\tilde{H}_i(X; Z_p) = 0$, $i \neq n$, and $H_n(X; Z_p) = Z_p \oplus F$, where F is free over $Z_p G$. Then the Borel formula holds, i.e.,*

$$n - n(G) = \sum(n(H) - n(G)),$$

sum extending over all corank 1 subgroups H in G .

Received by the editors August 14, 1978 and, in revised form, December 4, 1978.

AMS (MOS) subject classifications (1970). Primary 55C35; Secondary 55B25.

© 1980 American Mathematical Society
0002-9939/80/0000-0176/\$02.25

PROOF. Let k be the cellular dimension of X . The proof proceeds in roughly two stages. At first we construct inductively a finite CW complex, Y_{k-1} , with the property that Y_{k-1} is k -dimensional, $k - 1$ -connected. Y_{k-1} will also have the property that

$$\text{Ext}_{Z_p G}^{s+(k-n)}(H_n(X; Z_p), Z_p) = \text{Ext}_{Z_p G}^s(H_k(Y_{k-1}; Z_p), Z_p)$$

for all $s \geq 2$.

On the other hand we define a finite CW complex \hat{Y}_{k-1} which satisfies the hypothesis of Theorem A in the introduction, for the integer $m = \Sigma(n(H) - n(G)) + n(G)$ (not, at this point, necessarily equal to n). It will result that

$$\text{Ext}_{Z_p G}^{s+(k-n)}(H_n(X; Z_p), Z_p) = \text{Ext}_{Z_p G}^{s+(k-m)}(H_m(\hat{Y}_{k-1}; Z_p), Z_p)$$

for any $s \geq 2$. Since G has nonperiodic group cohomology for $r > 1$, it follows that $n = m$, which is what we wanted to show.

To begin with, we will use extensively the fact that whenever $f: M \rightarrow T \oplus P$ is a ZG epimorphism of ZG modules with M free, P projective and T consisting of torsion prime to p , $\ker f$ is ZG projective. This follows from [7, 4.12] once it is clear that T is cohomologically trivial. But multiplication by $|G|$ on T is an automorphism, which induces an automorphism on cohomology. Coupling this with the fact that $H^*(G, \cdot)$ has exponent $|G|$ (see Cartan-Eilenberg, Chapter XII, e.g.), it follows that T is cohomologically trivial (this argument was pointed out to me by R. Oliver).

We may assume X is 1-connected (by suspending). Since $\tilde{H}_i(X; Z)$ is torsion prime to p , we may attach free orbits of cells in dimensions $\leq n$ to produce the complex Y such that Y is $n - 1$ -connected and $H_n(Y; Z) = H_n(X; Z) \oplus N$, N a ZG -projective (for a nice exposition of the procedure of "equivariantly attaching cells", see [5, §1]). Now add free orbits of $n + 1$ -cells to kill off N . This produces a complex X_0 with $H_n(X_0; Z) = H_n(X; Z)$ and $H_{n+1}(X_0; Z) = H_{n+1}(X; Z) \oplus N_0$, N_0 a ZG -projective. $H_i(X_0; Z) = H_i(X; Z)$, $i > n + 1$.

Add free orbits of $n + 1$ -cells to X_0 to kill off $H_n(X; Z)$. This creates an n -connected, k -dimensional complex Y_n , so that $H_i(Y_n) = H_i(X)$, $i > n + 1$, and

$$0 \rightarrow H_{n+1}(X_0; Z) \rightarrow H_{n+1}(Y_n; Z) \rightarrow \text{Ker } \partial_{n+1} \rightarrow 0 \tag{*}$$

where $0 \rightarrow \text{Ker } \partial_{n+1} \rightarrow H_{n+1}(Y_n, X_0; Z) \rightarrow H_n(X_0; Z) \rightarrow 0$. Now if $k = n + 1$, it follows that $H_{n+1}(X_0; Z) = N_0$, since $H_{n+1}(X; Z_p) = 0$. By [7, 3.5], $N_0 \otimes Z_p$ is $Z_p G$ free. After tensoring the above sequences with Z_p , it follows that

$$\text{Ext}_{Z_p G}^s(H_n(X; Z_p), Z_p) = \text{Ext}_{Z_p G}^{s-1}(H_{n+1}(Y_n; Z_p), Z_p) \text{ for any } s \geq 2.$$

If $k > n + 1$, we show below how to construct Y_{n+1} , which is $n + 1$ -connected and k -dimensional, from Y_n . If need be, the following argument may be repeated until one obtains Y_{k-1} .

Add free orbits of $n + 2$ -cells to Y_n to kill off $H_{n+1}(X_0; Z)$ (see sequence (*)). This produces a complex Y'_{n+1} with $H_{n+1}(Y'_{n+1}; Z) = \text{Ker } \partial_{n+1}$ and $H_{n+2}(Y'_{n+1}; Z) = H_{n+2}(X_0; Z) \oplus N_1$, where N_1 is ZG -projective.

Now add free orbits of $n + 2$ -cells to Y'_{n+1} to form Y_{n+1} and to kill off $\text{Ker } \partial_{n+1}$. Y_{n+1} is $n + 1$ -connected, k -dimensional. Further, we have:

$$H_{n+2}(Y_{n+1}, Y'_{n+1}; Z) \rightarrow H_{n+1}(Y'_{n+1}, X_0; Z) \xrightarrow{\partial_{n+1}} H_n(X_0; Z) \rightarrow 0$$

where the left-hand map is the composition

$$H_{n+2}(Y_{n+1}, Y'_{n+1}; Z) \rightarrow H_{n+1}(Y'_{n+1}; Z) \approx \text{Ker } \partial_{n+1} \rightarrow H_{n+1}(Y'_{n+1}, X_0; Z).$$

We also have the exact sequence,

$$\begin{aligned} 0 \rightarrow H_{n+2}(Y'_{n+1}; Z) \rightarrow H_{n+2}(Y_{n+1}; Z) \\ \rightarrow H_{n+2}(Y_{n+1}, Y'_{n+1}; Z) \xrightarrow{\partial_{n+2}} H_{n+1}(Y'_{n+1}; Z). \end{aligned}$$

Now if $k = n + 2$, it follows that $H_{n+2}(Y'_{n+1}; Z) = N_1$, a ZG -projective, since $H_{n+2}(X; Z_p) = 0$. Thus $H_{n+2}(Y'_{n+1}; Z_p)$ is $Z_p G$ free. As before we have the exact sequences

$$\begin{aligned} 0 \rightarrow H_{n+2}(Y'_{n+1}; Z_p) \rightarrow H_{n+2}(Y_{n+1}; Z_p) \rightarrow \text{Ker } \partial_{n+2} \otimes Z_p, \\ 0 \rightarrow \text{Ker } \partial_{n+2} \otimes Z_p \rightarrow H_{n+2}(Y_{n+1}, Y'_{n+1}; Z_p) \rightarrow H_{n+1}(Y'_{n+1}; Z_p). \end{aligned}$$

Then

$$\text{Ext}_{Z_p G}^s(H_{n+1}(Y'_{n+1}; Z_p), Z_p) = \text{Ext}_{Z_p G}^{s-1}(H_{n+2}(Y_{n+1}; Z_p), Z_p)$$

and

$$\text{Ext}_{Z_p G}^s(H_n(X; Z_p), Z_p) = \text{Ext}_{Z_p G}^{s-1}(H_{n+1}(Y'_{n+1}; Z_p), Z_p)$$

since $H_{n+1}(Y'_{n+1}; Z_p) = \text{Ker } \partial_{n+1} \otimes Z_p$ and where s is arbitrary, $s \geq 2$.

If $k > n + 2$, we have the exact sequence

$$0 \rightarrow H_{n+2}(Y'_{n+1}; Z) \rightarrow H_{n+2}(Y_{n+1}; Z) \rightarrow \text{Ker } \partial_{n+2}.$$

Since $H_{n+2}(Y'_{n+1}; Z) = H_{n+2}(X_0; Z) \oplus N_1$, we can repeat the procedure above to reach, eventually, Y_{k-1} . By adding additional free orbits of cells, if necessary, we may assume $k - m \geq 2$.

The same argument as above shows that for any $s \geq 2$,

$$\text{Ext}_{Z_p G}^{s+(k-n)}(H_n(X; Z_p), Z_p) = \text{Ext}_{Z_p G}^s(H_k(Y_{k-1}; Z_p), Z_p).$$

Now, on the other hand, consider $\hat{Y}_{k-1} = Y_{k-1}^{(m)} \cup Y_{k-1}^F$, where $Y_{k-1}^F = \bigcup_{H \neq 0} Y_{k-1}^H = \bigcup_{H \neq 0} X^H$. We claim that $\tilde{H}_i(\hat{Y}_{k-1}; Z_p) = 0$ for $i \neq m$. This amounts to showing that $H_*(\bigcup_{H \neq 0} X^H; Z_p) = 0$ for $* > m$. First of all, by induction on the rank of G, r , we may assume the Borel formula for any elementary abelian p -group of lesser rank. This induction implies that $n(H) \leq m$, for $H \neq 0$. Letting S be any nonempty collection of nonzero subgroups of G (without loss, not containing G itself) one actually establishes the stronger fact that $H_*(\bigcup_{H \in S} X^H; Z_p) = 0$ if $* > m$. To show this one uses induction on the cardinality of S and Mayer-Vietoris. Thus one is led to see it suffices to show that, for $M_0 \in S$,

$$H_*(X^{M_0}, \bigcup X^{MM_0}; Z_p) = 0, \text{ for } * > n(M_0),$$

where the union is over $S - \{M_0\}$. This latter fact is established by a double induction on rank M_0 and on the cardinality of $S - \{M_0\}$.

There are two naturally arising cases. If there is some M_1 in $S - \{M_0\}$ with $n(M_1M_0) < n(M_0)$ one considers the homology sequence of the triple $(X^{M_0}, \cup X^{MM_0}, \cup X^{MM_0})$, where the first union is over $S - \{M_0\}$ and the second is over $S - \{M_0, M_1\}$. An application of excision establishes the result in this case.

The second case is that every $M \in S$ has $n(MM_0) = n(M_0)$. Here, it suffices to show that if M_1 and M_2 both have $n(M_1M_0) = n(M_2M_0) = n(M_0)$ then $n(M_0) = n(M_1M_2M_0)$ (by a Mayer-Vietoris argument). To demonstrate this last property use the fact that the Borel formula holds whenever a group of rank $\leq r - 1$ is acting.

Now since $\tilde{H}_i(\hat{Y}_{k-1}; Z_p) = 0, i \neq m$, and the Borel formula holds on \hat{Y}_{k-1} , by Theorem A in the introduction (it is not hard to see that if $r > 1, m - n(G)$ is always even if p is odd)

$$H_m(\hat{Y}_{k-1}; Z_p) = Z_p \oplus F_1,$$

where F_1 is a free $Z_p G$ module.

Now consider $C_*(Y_{k-1}, \hat{Y}_{k-1}; Z_p)$. We have the exact sequence,

$$\begin{aligned} 0 \rightarrow H_k(Y_{k-1}; Z_p) \rightarrow C_k(Y_{k-1}, \hat{Y}_{k-1}; Z_p) \rightarrow \dots \\ \rightarrow C_{m+1}(Y_{k-1}, \hat{Y}_{k-1}; Z_p) \twoheadrightarrow H_m(\hat{Y}_{k-1}; Z_p). \end{aligned}$$

Since each $C_i(Y_{k-1}, \hat{Y}_{k-1}; Z_p)$ is $Z_p G$ free, it follows that for $s \geq 2$,

$$\text{Ext}_{Z_p G}^{s+(k-m)}(H_m(\hat{Y}_{k-1}; Z_p), Z_p) = \text{Ext}_{Z_p G}^s(H_k(Y_{k-1}; Z_p), Z_p).$$

Thus,

$$\text{Ext}_{Z_p G}^{s+(k-n)}(H_n(X; Z_p), Z_p) = \text{Ext}_{Z_p G}^{s+(k-m)}(H_m(\hat{Y}_{k-1}; Z_p), Z_p).$$

But since $H_n(X; Z_p)$ and $H_m(\hat{Y}_{k-1}; Z_p)$ are both stably Z_p , the result is that

$$\text{Ext}_{Z_p G}^{s+(k-n)}(Z_p, Z_p) = \text{Ext}_{Z_p G}^{s+(k-m)}(Z_p, Z_p).$$

This implies that $n = m$, since G has nonperiodic cohomology. \square

3. We now consider examples of Proposition 1 which are n -dimensional, $n - 1$ connected.

PROPOSITION 2. *Suppose that X satisfies the hypotheses of Proposition 1 and that X is n -dimensional, $n - 1$ connected. Then $H_n(X; Z) = Z \oplus P$, where P is ZG -projective and Z has trivial G -action if p is odd.*

PROOF. It was shown in [4] that there is a linear G -action on S^n and an equivariant map $\phi: X \rightarrow S^n$ having the properties described in the introduction. From the mapping cone C_ϕ , we have the exact sequence

$$0 \rightarrow H_{n+1}(C_\phi; Z) \rightarrow H_n(X; Z) \xrightarrow{\phi_*} H_n(S^n; Z) \rightarrow H_n(C_\phi; Z) \rightarrow 0.$$

Because ϕ_* is a Z_p -homology epimorphism we have

$$0 \rightarrow H_{n+1}(C_\phi; Z) \rightarrow H_n(X; Z) \twoheadrightarrow \text{Im } \phi_* \approx Z \quad (\text{with some action}). \quad (*)$$

Also, $H_n(C_\phi; Z) = Z_k$, where $(k, p) = 1$ and $\text{Im } \phi_* = (k)$. One may add free orbits of $n + 1$ cells to C_ϕ to kill $H_n(C_\phi; Z)$. This results in an $(n + 1)$ -dimensional,

n -connected G -complex Y such that $H_{n+1}(Y; Z) = H_{n+1}(C_\phi; Z) \oplus P$, P a projective ZG module. By [6, Lemma 3], $H_{n+1}(Y; Z)$ is ZG -projective and then so is $H_{n+1}(C_\phi; Z)$.

If $\text{Im } \phi_* \approx Z$ has trivial G -action then one has

$$\text{Ext}_{ZG}(Z; H_{n+1}(C_\phi; Z)) = H^1(G; H_{n+1}(C_\phi; Z)) = 0$$

by [6, 4.11]. Thus in this case, (*) splits.

If $\text{Im } \phi_* \approx Z$ has nontrivial G -action (denoted Z^-), it follows that $Z^- \otimes_Z Z^- \approx Z$ has trivial (diagonal) G -action and that whenever P is ZG -projective, so is $Z^- \otimes_Z P$ (diagonal action). This latter statement follows from the fact that $Z^- \otimes_Z ZG$ (diagonal action) is isomorphic, as ZG modules, to ZG . Now tensoring (*) over Z with Z^- one has the exact sequence of ZG modules,

$$0 \rightarrow H_{n+1}(C_\phi; Z) \otimes_Z Z^- \rightarrow H_n(X; Z) \otimes_Z Z^- \rightarrow Z \rightarrow 0. \quad (*)'$$

By the argument above, (*)' splits. So,

$$H_n(X; Z) \otimes_Z Z^- \approx Z \oplus H_{n+1}(C_\phi; Z) \otimes_Z Z^-.$$

Now, tensoring again with Z^- over Z , one has $H_n(X; Z) \approx Z^- \oplus H_{n+1}(C_\phi; Z)$.

□

As an example, one may take the wedge of a standard linear action on S^n with a bouquet of n -spheres permuted freely by G . On the other hand, suppose Y is an n -dimensional $n - 1$ -connected CW complex with G -action such that Y^H is Z_p -acyclic for each $H \triangleleft G, H \neq 0$. By [6, Lemma 3], $H_n(Y)$ is ZG -projective. Wedging Y with a linear S^n would yield other examples. However, it is a result of Swan (see [1, p. 178]) that $H_n(Y)$ must be stably free, when $G = Z_p$. Consequently, one can construct projective modules P over $ZG, G = (Z_p)^r, r > 1$, which could never be realised as an $H_n(Y)$, as above. Mainly, this is so because a necessary condition on P is that the image of P in $G_0(ZG)/S$ be zero, where $G_0(ZG)$ is the Grothendieck group on finitely generated ZG modules and S is the subgroup generated by signed permutation modules (see [1]). As a result it is not clear precisely when a projective ZG module can be an $H_n(Y)$, as above. According to [1] the necessary condition above is sufficient for $G = Z_p$.

BIBLIOGRAPHY

1. J. E. Arnold, *A solution of a problem of Steenrod for cyclic groups of prime order*, Proc. Amer. Math. Soc. **62** (1977), 177-182.
2. A. Borel, *Seminar on transformation groups*, Ann. of Math. Studies, no. 46, Princeton Univ. Press, Princeton, N. J., 1960.
3. G. Bredon, *Introduction to compact transformation groups*, Academic Press, New York, 1972.
4. R. Dotzel, *A converse of the Borel formula*, Trans. Amer. Math. Soc. **250** (1979), 275-287.
5. L. Jones, *The converse to the fixed point theorem of P. A. Smith. I*, Ann. of Math. (2) **94** (1971), 52-68.
6. R. Oliver, *Fixed point sets of group actions on finite acyclic complexes*, Comment. Math. Helv. **50** (1975), 155-177.
7. D. S. Rim, *Modules over finite groups*, Ann. of Math. (2) **69** (1959), 700-712.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS, AUSTIN, TEXAS 78712