

HOW TO RECOGNIZE A LOCALIZED SPHERE

J. AGUADÉ

ABSTRACT. Two cohomological characterizations of the sphere localized at a prime are given.

The purpose of this note is to obtain necessary and sufficient conditions on the cohomology of a nilpotent space in order to insure that it has the homotopy type of S_p^m , the sphere localized at the prime p (cf. [2]). p will denote a fixed prime integer. $\hat{\mathbf{Z}}_p$, $\mathbf{Z}_{(p)}$, \mathbf{Z}_n , \mathbf{Z}_{p^∞} denote the p -adic integers, the integers localized at p , the integers modulo n and the p^m th roots of unity with m running over all integers ≥ 0 , respectively. Group will always mean abelian group. A group is called p -local if it is q -divisible for every $q \neq p$ and has no q -torsion for every $q \neq p$. A nilpotent space is called p -local if its homology groups with integral coefficients are p -local [2, p. 53].

Recall that a subgroup B of a group A is called p -basic if the following holds:

- (i) B is a direct sum of cyclic p -groups and infinite cyclic groups,
- (ii) A/B is p -divisible,
- (iii) $B/p^k B$ is a direct summand of $A/p^k B$ for all k .

Every group contains p -basic subgroups [1, I, p. 137].

We use freely the most elementary properties of the functors Hom and Ext. A standard reference is [1, Chapters VIII and IX]. In particular we shall use the following:

$$\text{Ext}(\mathbf{Z}_n, A) = A/nA, \text{Ext}(\mathbf{Z}_{p^\infty}, \mathbf{Z}_p) = \mathbf{Z}_p, \text{Ext}(\mathbf{Q}, \mathbf{Z}_{(p)}) = \mathbf{Q}^{*0}, \text{Ext}(\mathbf{Z}_{p^\infty}, \mathbf{Z}_{(p)}) = \hat{\mathbf{Z}}_p.$$

All spaces are assumed to be of the homotopy type of CW-complexes.

LEMMA 1. *Let B be a p -basic subgroup of A . Then $\text{Hom}(A, \mathbf{Z}_p) = \text{Hom}(B, \mathbf{Z}_p)$.*

PROOF. Since A/B is p -divisible and \mathbf{Z}_p contains no p -divisible subgroups other than 0, the Hom-Ext exact sequence associated to $B \rightarrow A \rightarrow A/B$ yields that $\text{Hom}(A, \mathbf{Z}_p)$ is a subgroup of $\text{Hom}(B, \mathbf{Z}_p)$. Let $f: B \rightarrow \mathbf{Z}_p$. Clearly, f factorizes through $f': B/pB \rightarrow \mathbf{Z}_p$. Since B is a p -basic subgroup of A , B/pB is a direct summand of A/pB . Hence f' extends to A/pB and yields a homomorphism $g \in \text{Hom}(A, \mathbf{Z}_p)$. \square

LEMMA 2. *If A is a p -local group such that $\text{Hom}(A, \mathbf{Q}) = \text{Hom}(A, \mathbf{Z}_p) = 0$, then A is a divisible p -group.*

Received by the editors January 18, 1979.

AMS (MOS) subject classifications (1970). Primary 55J20, 20K30, 20K35.

Key words and phrases. Localization, nilpotent spaces, universal coefficient theorem.

© 1980 American Mathematical Society
0002-9939/80/0000-0179/\$02.00

PROOF. Since $\text{Hom}(A, \mathbf{Q}) = 0$, A has no elements of infinite order. Since A is p -local, A is a p -group. Let B be a p -basic subgroup. By Lemma 1, $\text{Hom}(B, \mathbf{Z}_p) = \text{Hom}(A, \mathbf{Z}_p) = 0$ and since B is a direct sum of cyclic p -groups then $B = 0$. That is A is p -divisible and so, since A is p -local, A is divisible. \square

LEMMA 3. Let E be a nilpotent space such that $H_*(E; \mathbf{Z}) = H_*(S_p^m; \mathbf{Z})$. Then E has the homotopy type of S_p^m .

PROOF. If $m > 1$ then $H_1 E = 0$. In other words, the abelianization of $\pi_1 E$ is trivial. Since $\pi_1 E$ is a nilpotent group we conclude that E is simply connected and so E is a Moore space $M(\mathbf{Z}_{(p)}, m)$, that is a localized sphere S_p^m .

If $m = 1$ then we have

$$[E, S_p^1] = [E, K(\mathbf{Z}_{(p)}, 1)] \cong H^1(E; \mathbf{Z}_{(p)}) = \mathbf{Z}_{(p)}$$

where $K(\mathbf{Z}_{(p)}, 1)$ denotes an Eilenberg-Mac Lane space of type $(\mathbf{Z}_{(p)}, 1)$. Then there exists a map $f: E \rightarrow S_p^1$ which induces isomorphisms in homology. Because E and S_p^1 are nilpotent spaces, f must be a homotopy equivalence. The result follows. \square

THEOREM 4. Let E be a nilpotent space such that

- (1) E is p -local,
- (2) $H^*(E; R) = H^*(S^m; R)$ where $R = \mathbf{Z}_p$ or $R = \hat{\mathbf{Z}}_p$,
- (3) $H^*(E; \mathbf{Q}) = H^*(S^m; \mathbf{Q})$,
- (4) $H^{m+1}(E; \mathbf{Z}_{(p)}) = 0$.

Then $E = S_p^m$. If any of the above conditions is omitted then the conclusion is false.

PROOF. It suffices to prove that $H_*(E; \mathbf{Z}) = H_*(S_p^m; \mathbf{Z})$. Let us consider the exact sequence $\hat{\mathbf{Z}}_p \xrightarrow{p} \hat{\mathbf{Z}}_p \rightarrow \mathbf{Z}_p$. It induces a long exact sequence in cohomology:

$$\dots \rightarrow H^r(E; \hat{\mathbf{Z}}_p) \xrightarrow{p} H^r(E; \hat{\mathbf{Z}}_p) \rightarrow H^r(E; \mathbf{Z}_p) \rightarrow H^{r+1}(E; \hat{\mathbf{Z}}_p) \rightarrow \dots$$

From this it follows easily that we may assume, without loss of generality that $R = \mathbf{Z}_p$ in condition (2). On the other hand, condition (3) implies that $H_r E$ has torsion-free rank zero if $r \neq m$ and has torsion-free rank one if $r = m$.

The universal coefficient theorem:

$$\text{Ext}(H_{r-1} E, \mathbf{Z}_p) \rightarrow H^r(E; \mathbf{Z}_p) \rightarrow \text{Hom}(H_r E, \mathbf{Z}_p)$$

yields: $\text{Hom}(H_r E, \mathbf{Z}_p) = 0$ if $r \neq m$ and $\text{Ext}(H_r E, \mathbf{Z}_p) = 0$ if $r \neq m - 1$. It follows from Lemma 2 that $H_r E$ is a divisible p -group if $r \neq m$. The structure theorem for divisible groups [1, I, p. 104] shows that $H_r E = \bigoplus \mathbf{Z}_{p^\infty}$. But $\text{Ext}(H_r E, \mathbf{Z}_p) = 0$ if $r \neq m - 1$, hence $H_r E = 0$ if $r \neq m, m - 1$ because $\text{Ext}(\mathbf{Z}_{p^\infty}, \mathbf{Z}_p) = \mathbf{Z}_p$.

Since $H^m(E; \mathbf{Z}_p) = \mathbf{Z}_p$ the universal coefficient theorem shows that there are two cases:

- (A) $\text{Ext}(H_{m-1} E, \mathbf{Z}_p) = \mathbf{Z}_p, \text{Hom}(H_m E, \mathbf{Z}_p) = 0$.

$H_{m-1} E$ is a divisible p -group and so $H_{m-1} E = \bigoplus \mathbf{Z}_{p^\infty}$. Since $\text{Ext}(H_{m-1} E, \mathbf{Z}_p) = \mathbf{Z}_p$ it follows that $H_{m-1} E = \mathbf{Z}_{p^\infty}$. On the other hand, let B be a p -basic subgroup of $H_m E$. By Lemma 1, $\text{Hom}(B, \mathbf{Z}_p) = \text{Hom}(H_m E, \mathbf{Z}_p) = 0$ and so $B = 0$ and $H_m E$ is divisible because it is p -local and p -divisible. Since $H_m E$ has torsion-free rank one and $\text{Ext}(H_m E, \mathbf{Z}_p) = 0$ we conclude that $H_m E = \mathbf{Q}$. Hence, we see that in this case

the space E has the homology of $M(\mathbf{Q}, m) \vee M(\mathbf{Z}_{p^\infty}, m - 1)$, the wedge of the Moore spaces $M(\mathbf{Q}, m)$ and $M(\mathbf{Z}_{p^\infty}, m - 1)$.

(B) $\text{Ext}(H_{m-1}E, \mathbf{Z}_p) = 0, \text{Hom}(H_mE, \mathbf{Z}_p) = \mathbf{Z}_p$.

$H_{m-1}E$ is a divisible p -group such that $\text{Ext}(H_{m-1}E, \mathbf{Z}_p) = 0$. Since $\text{Ext}(\mathbf{Z}_{p^\infty}, \mathbf{Z}_p) = \mathbf{Z}_p$ then $H_{m-1}E = 0$ and so E has homology $\neq 0$ only in dimension m . Let B be a p -basic subgroup of H_mE . We have $\text{Hom}(B, \mathbf{Z}_p) = \text{Hom}(H_mE, \mathbf{Z}_p) = \mathbf{Z}_p$. Since B is a direct sum of infinite cyclic groups and cyclic p -groups, either $B = \mathbf{Z}$ or $B = \mathbf{Z}_{p^k}$. If $B = \mathbf{Z}_{p^k}$ then condition (iii) in the definition of p -basic subgroup yields that \mathbf{Z}_{p^k} is a direct summand of H_mE , but this is impossible because $\text{Ext}(H_mE, \mathbf{Z}_p) = 0$. Hence \mathbf{Z} is a p -basic subgroup of H_mE . Since $D = H_mE/\mathbf{Z}$ is divisible and $\text{Ext}(H_mE, \mathbf{Z}_p) = 0$, the Hom-Ext exact sequence associated to $\mathbf{Z} \rightarrow H_mE \rightarrow D$ gives $\text{Ext}(D, \mathbf{Z}_p) = 0$. If we apply now the structure theorem for divisible groups we obtain that D has no p -torsion and so H_mE is torsion-free because it is p -local and so it can only have p -torsion. Then H_mE is a p -local torsion-free group of rank one. We apply now the classification theorem for these groups [1, II, p. 110]. Since H_mE is p -local, in order to prove that $H_mE = \mathbf{Z}_{(p)}$ it suffices to show that H_mE contains elements of p -height zero. Let us consider $1 \in \mathbf{Z} \subset H_mE$. If $1 = pa, a \in H_mE$ then in D we have $0 = p\bar{a}$ and since D has no p -torsion we get $a \in \mathbf{Z}$, a contradiction. Hence, the type of H_mE is $t_q = \infty$ if $q \neq p$ and $t_p = 0$. Then the space E has the same integral homology as S_p^m and then, by Lemma 3, $E = S_p^m$.

We have proved that if a nilpotent space E satisfies conditions (1), (2), (3) of the theorem then either $E = S_p^m$ or E has the same homology as $M(\mathbf{Q}, m) \vee M(\mathbf{Z}_{p^\infty}, m - 1)$. It suffices to show that if $H_mE = \mathbf{Q}$ then E does not satisfy condition (4). This follows easily from the fact $\text{Ext}(\mathbf{Q}, \mathbf{Z}_{(p)}) = \mathbf{Q}^{*0}$.

Finally, note that none of the conditions (1), (2), (3), (4) can be omitted. Let us consider the spaces $E_1 = S_p^m \vee M(\mathbf{Z}_q, 2)$ (q a prime $\neq p$); $E_2 = S_p^m \vee M(\mathbf{Z}_p, 2m)$; $E_3 = M(\mathbf{Z}_{p^\infty}, m - 1)$; $E_4 = M(\mathbf{Q}, m) \vee M(\mathbf{Z}_{p^\infty}, m - 1)$. It is easily seen that the space E_i verifies conditions (1), (2), (3), (4) except the i th, but $E_i \neq S_p^m$. \square

The above theorem shows that the cohomology with coefficients $\mathbf{Z}_p, \mathbf{Z}_p, \mathbf{Q}$ is not enough to characterize the localized spheres. The following theorem shows that the cohomology with coefficients $\mathbf{Z}_{(p)}$ is suitable for this purpose.

THEOREM 5. *Let E be a p -local nilpotent space such that $H^*(E; \mathbf{Z}_{(p)}) = H^*(S^m; \mathbf{Z}_{(p)})$. Then $E = S_p^m$.*

PROOF. From the universal coefficient theorem we obtain $\text{Hom}(H_rE, \mathbf{Z}_{(p)}) = 0$ if $r \neq m$ and $\text{Ext}(H_rE, \mathbf{Z}_{(p)}) = 0$ if $r \neq m - 1$. Further, $\text{Ext}(H_{m-1}E, \mathbf{Z}_{(p)})$ is a p -local subgroup of $\mathbf{Z}_{(p)}$. Then either $\text{Ext}(H_{m-1}E, \mathbf{Z}_{(p)}) = \mathbf{Z}_{(p)}$ or $\text{Ext}(H_{m-1}E, \mathbf{Z}_{(p)}) = 0$. But in the first case $\mathbf{Z}_{(p)}$ will be a cotorsion group (cf. [1, I, p. 232] and Theorem 54.6 of [1, I, p. 235]) and this is impossible because $\text{Ext}(\mathbf{Q}, \mathbf{Z}_{(p)}) = \mathbf{Q}^{*0} \neq 0$. Hence, $\text{Ext}(H_rE, \mathbf{Z}_{(p)}) = 0$ for all r .

Let B_r be a p -basic subgroup of H_rE . If B_r contains a direct summand of the form \mathbf{Z}_{p^k} then it follows easily from the definition of p -basic subgroup that \mathbf{Z}_{p^k} is a direct summand of H_rE . But $\text{Ext}(\mathbf{Z}_{p^k}, \mathbf{Z}_{(p)}) = \mathbf{Z}_{(p)}/p^k\mathbf{Z}_{(p)} \neq 0$. Hence, B_r is free. Let us consider the Hom-Ext exact sequence associated to $\mathbf{Z}_{(p)} \xrightarrow{p} \mathbf{Z}_{(p)} \rightarrow \mathbf{Z}_p$:

$$0 \rightarrow \text{Hom}(H_r E, \mathbf{Z}_{(p)}) \xrightarrow{p} \text{Hom}(H_r E, \mathbf{Z}_{(p)}) \rightarrow \text{Hom}(H_r E, \mathbf{Z}_p) \rightarrow 0.$$

We get that $\text{Hom}(H_r E, \mathbf{Z}_p) = 0$ if $r \neq m$. By Lemma 1, $\text{Hom}(B_r, \mathbf{Z}_p) = \text{Hom}(H_r E, \mathbf{Z}_p) = 0$, hence $B_r = 0$ if $r \neq m$. This shows that $H_r E$ is a divisible group for $r \neq m$. But the structure theorem for divisible groups and the equalities $\text{Ext}(\mathbf{Q}, \mathbf{Z}_{(p)}) = \mathbf{Q}^{\kappa_0}$, $\text{Ext}(\mathbf{Z}_{p^\infty}, \mathbf{Z}_{(p)}) = \hat{\mathbf{Z}}_p$, $\text{Ext}(H_r E, \mathbf{Z}_{(p)}) = 0$ yield $H_r E = 0$ if $r \neq m$.

In the case $r = m$ the above exact sequence shows that $\text{Hom}(H_m E, \mathbf{Z}_p) = \mathbf{Z}_p$. By Lemma 1, $B_m = \mathbf{Z}$. The Hom-Ext exact sequence associated to $\mathbf{Z} \rightarrow H_m E \rightarrow H_m E/\mathbf{Z}$ shows that $\text{Ext}(H_m E/\mathbf{Z}, \mathbf{Z}_{(p)})$ is the image of the countable group $\text{Hom}(\mathbf{Z}, \mathbf{Z}_{(p)})$. Since $\text{Ext}(\mathbf{Q}, \mathbf{Z}_{(p)})$ and $\text{Ext}(\mathbf{Z}_{p^\infty}, \mathbf{Z}_{(p)})$ are both uncountable and since $H_m E/\mathbf{Z}$ is a divisible group, we get that $H_m E/\mathbf{Z}$ is a torsion group without p -torsion. This leads to $\text{Hom}(H_m E/\mathbf{Z}, \mathbf{Q}) = 0$ and so $\text{Hom}(H_m E, \mathbf{Q}) = \text{Hom}(\mathbf{Z}, \mathbf{Q}) = \mathbf{Q}$. Hence $H_m E$ is a group of torsion-free rank one. Further, $H_m E$ is torsion-free because $H_m E/\mathbf{Z}$ has no p -torsion and $H_m E$ is p -local. Now we can show, as in the proof of Theorem 4 that $H_m E = \mathbf{Z}_{(p)}$ and so the space E has the same integral homology as S_p^m . Then, by Lemma 3, E is a localized sphere S_p^m . \square

REFERENCES

1. L. Fuchs, *Infinite abelian groups*, Academic Press, New York, 1970.
2. P. Hilton, G. Mislin, and J. Roitberg, *Localization of nilpotent groups and spaces*, North-Holland Mathematics Studies 15, North-Holland, Amsterdam, 1975.

SECCIÓ DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, BELLATERRA (BARCELONA), SPAIN

Current address: Forschungs Institut für Mathematik, ETH-Zentrum, 8092-Zürich, Switzerland