

## AN APPLICATION OF YAU'S MAXIMUM PRINCIPLE TO CONFORMALLY FLAT SPACES

S. I. GOLDBERG

**ABSTRACT.** Results of M. Tani on compact conformally flat manifolds and of M. Okumura on compact hypersurfaces of Euclidean space are extended to complete spaces by an application of S.-T. Yau's "maximum principle".

**1. Introduction.** M. Tani [3] proved that a compact and orientable Riemannian manifold admitting a conformally flat metric of positive Ricci curvature and constant scalar curvature is a space form, that is, it is a constant curvature space. It is our purpose to extend this result to complete Riemannian manifolds with Ricci curvature bounded from below. This will be accomplished by employing a "maximum principle" due to S.-T. Yau. In fact, the following statement is obtained.

**THEOREM 1.** *Let  $M$  be a  $d$ -dimensional,  $d > 3$ , complete, conformally flat Riemannian manifold whose Ricci curvature is bounded from below. If its scalar curvature  $r$  is a positive constant and  $\text{tr } Q^2 < r^2/(d - 1)$ , then  $M$  is a space form.*

**2. Definitions and notation.** Let  $(M, g)$  be a Riemannian manifold with metric  $g$ . The curvature transformation  $R(X, Y)$ ,  $X, Y \in M_m$ , where  $M_m$  is the tangent space at  $m \in M$ , and  $g$  are related by

$$R(X, Y) = \nabla_{[X, Y]} - [\nabla_X, \nabla_Y],$$

where  $\nabla$  is the Riemannian connection. In terms of a basis  $X_1, \dots, X_d$  of  $M_m$ , we set

$$R_{ijkh} = g(R(X_i, X_j)X_k, X_h), \quad R_{ij} = \text{tr}(X_k \rightarrow R(X_i, X_k)X_j),$$

$$t_{i_1 \dots i_p} = t(X_{i_1}, \dots, X_{i_p}), \quad \nabla_i t_{i_1 \dots i_p} = (\nabla_{X_i} t)(X_{i_1}, \dots, X_{i_p}).$$

We denote the scalar curvature by  $r$ , that is,  $r = \text{tr } Q$ , where  $Q = (R_j^i)$  and  $R_j^i = g^{ik}R_{jk}$ . The manifold  $(M, g)$  is *conformally flat* if  $g$  is conformally related to a locally flat metric.

**3. The Laplacian of  $\text{tr } Q^2$ .** The following formula may be found in [1]:

$$\frac{1}{2} \Delta \text{tr } Q^2 = g^{ab} \nabla_a R^{ij} \nabla_b R_{ij} + R^{ij} g^{ab} \nabla_a (\nabla_b R_{ij} - \nabla_i R_{bj}) + \frac{1}{2} R^{ij} \nabla_j \nabla_i r + K, \tag{3.1}$$

where  $\text{tr } Q^2$  is the square length of the Ricci tensor, and

$$K = R^{ik} (R_i^j R_{jk} + R^{hj} R_{ijhk}).$$

---

Received by the editors July 9, 1979.

AMS (MOS) subject classifications (1970). Primary 53C20; Secondary 53A07, 53C40.

Key words and phrases. Conformally flat, Ricci curvature, scalar curvature.

© 1980 American Mathematical Society  
 0002-9939/80/0000-0273/\$01.75

If  $r$  is a constant, the third term on the r. h. s. of (3.1) vanishes. If, moreover,  $M$  is conformally flat and  $d > 3$ , the second term on the right also vanishes (see [1]) and (3.1) reduces to

$$\frac{1}{2} \Delta \text{tr } Q^2 = K + g(\nabla Q, \nabla Q).$$

**4. Proof of Theorem 1.** Since  $M$  is conformally flat it can be shown that

$$(d-1)(d-2)K = d(d-1)\text{tr } Q^3 - r(2d-1)\text{tr } Q^2 + r^3.$$

Put  $S = Q - (r/d)I$ ,  $I = \text{identity}$ . Then, from  $\text{tr } S^2 > 0$ , we see that  $\text{tr } Q^2 > r^2/d$  with equality holding if and only if,  $M$  is an Einstein space. Since  $r$  is a constant, the Laplacian  $\Delta f^2$  of the function  $f^2 = \text{tr } S^2$ ,  $f > 0$ , satisfies  $\Delta f^2 = \Delta \text{tr } Q^2$ . Thus,

$$\frac{1}{2} \Delta f^2 = K + g(\nabla Q, \nabla Q). \quad (4.1)$$

Moreover,

$$(d-1)(d-2)K = d(d-1)\left(\text{tr } S^3 + \frac{3r}{d} f^2 + \frac{r^3}{d^2}\right) - r(2d-1)\left(f^2 + \frac{r^2}{d}\right) + r^3. \quad (4.2)$$

The following lemma may be found in [2].

**LEMMA 1.** Let  $a_i$ ,  $i = 1, \dots, d$ , be real numbers with

$$\sum_{i=1}^d a_i = 0, \quad \sum_{i=1}^d a_i^2 = k^2, \quad k = \text{const} > 0.$$

Then,

$$-\frac{d-2}{\sqrt{d(d-1)}} k^3 < \sum_{i=1}^d a_i^3 < \frac{d-2}{\sqrt{d(d-1)}} k^3.$$

Applying Lemma 1 to the eigenvalues of  $S$ , (4.2) yields the inequality

$$(d-1)K > f^2(r - \sqrt{d(d-1)})f.$$

We conclude from (4.1) that

$$\frac{d-1}{2} \Delta f^2 > f^2(r - \sqrt{d(d-1)})f. \quad (4.3)$$

**LEMMA 2** (S.-T. YAU [4]). Let  $M$  be a complete Riemannian manifold with Ricci curvature bounded below. Let  $u$  be a  $C^2$  function with  $\sup u < \infty$ . Then, there exists a sequence  $\{p_\nu\}$  in  $M$  such that

$$\lim_{\nu \rightarrow \infty} \|du(p_\nu)\| = 0, \quad \lim_{\nu \rightarrow \infty} (\Delta u)(p_\nu) < 0, \quad \lim_{\nu \rightarrow \infty} u(p_\nu) = \sup u.$$

Applying Lemma 2, the inequality (4.3) gives rise to the inequality

$$\lim_{\nu \rightarrow \infty} f^2(p_\nu)\{r - \sqrt{d(d-1)}f(p_\nu)\} < 0.$$

Hence, either  $f^2 \equiv 0$  or  $\sup f > r/\sqrt{d(d-1)}$ , the latter implying  $\sup \text{tr } Q^2 > r^2/(d-1)$ . The former says that  $\text{tr } Q^2 = r^2/d$ , so  $g$  is an Einstein metric. However, since  $g$  is conformally flat, it is a constant curvature metric.

The condition  $\text{tr } Q^2 < r^2/(d - 1)$  is essential. For, if  $M = M_1 \times N$ , where  $M_1$  has constant curvature and  $N$  is 1-dimensional, then  $M$  is conformally flat, its Ricci curvature is bounded below,  $r$  is constant and  $\text{tr } Q^2 = r^2/(d - 1)$ .

In a similar manner, we obtain the following extension of a theorem of Okumura [2].

**THEOREM 2.** *Let  $M$  be a  $d$ -dimensional complete connected hypersurface of  $R^{d+1}$  with Ricci curvature bounded from below. If its mean curvature  $\text{tr } H$  is constant and  $\text{tr } H^2 < (\text{tr } H)^2/(d - 1)$ , then  $M$  is a totally umbilical hypersurface.*

The inequality in Theorem 2 is the best possible as one sees by considering  $M = S^{d-1} \times R$ .

#### REFERENCES

1. S. I. Goldberg, *On conformally flat spaces with definite Ricci curvature*, Kōdai Math. Sem. Rep. **21** (1969), 226–232.
2. M. Okumura, *Hypersurfaces and a pinching problem on the second fundamental tensor*, Amer. J. Math. **86** (1974), 207–213.
3. M. Tani, *On a conformally flat Riemannian space with positive Ricci curvature*, Tôhoku Math. J. **19** (1967), 227–231.
4. S.-T. Yau, *Harmonic functions on complete Riemannian manifolds*, Comm. Pure Appl. Math. **28** (1975), 201–228.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA, ILLINOIS 61801