

NONEXTENDED QUADRATIC FORMS OVER POLYNOMIAL RINGS OVER POWER SERIES RINGS

RAMAN PARIMALA

ABSTRACT. If R is a complete discrete valuation ring, then every quadratic space over $R[T]$ is extended from R . We here show by an example that a corresponding result for higher-dimensional complete regular local rings is not valid.

It was proved in [3] that if R is any complete discrete valuation ring, every quadratic space over $R[T]$ is extended from R . We show in this note that if $R = \mathbf{R}[[X, Y]]$, there exist anisotropic quadratic spaces over $R[T]$ which are not extended from R . This is in contrast to a result of Mohan Kumar and Lindel in the linear case [2, Theorem 5.1, p. 150].

Let $R = \mathbf{R}[[X, Y]]$, \mathbf{R} denoting the field of real numbers. Let \mathbf{H} be the quaternions over \mathbf{R} and $A = \mathbf{H}[[X, Y]]$ the ring of formal power series in the variables X and Y over \mathbf{H} . We have an $A[T]$ -linear map $A[T]^2 \xrightarrow{\eta} A[T]$ defined by $(1, 0) \rightarrow XT + i$, $(0, 1) \rightarrow YT + j$ which is clearly a surjection. Let P be the kernel of η .

PROPOSITION. *The module P is nonfree projective over $A[T]$.*

PROOF. The first projection of $A[T]^2$ onto $A[T]$ maps P isomorphically onto the left ideal \mathfrak{A} of $A[T]$ generated by $1 + Y^2T^2$ and $1 + (iX + jY)T - kXYT^2$ [4, p. 143]. We prove that P is not free by showing that \mathfrak{A} is not principal. We note first that \mathfrak{A} is not the unit ideal since it is generated modulo X by $1 + jYT$ which is not a unit in $\mathbf{H}[[Y]][T]$. Suppose \mathfrak{A} is principal, generated by f . Then $\deg_T f < 2$. If $\deg_T f = 0$, then it follows that \mathfrak{A} is the unit ideal, which is not the case. If $\deg_T f = 2$, then $1 + Y^2T^2$ and $1 + (iX + jY)T - kXYT^2$ are unit left multiples of each other, which is again not possible. Let $\deg_T f = 1$ and $f = a + bT$, $a, b \in \mathbf{H}[[X, Y]]$. Then a is a unit and we assume $a = 1$ so that $f = 1 + bT$, $b \in \mathbf{H}[[X, Y]]$. We then have

$$1 + Y^2T^2 = (1 + cT)(1 + bT),$$

$$1 + (iX + jY)T - kXYT^2 = (1 + dT)(1 + bT).$$

From the first equation we get $c = -b$, $-b^2 = Y^2$. This implies that $b = \lambda Y$, $\lambda \in \mathbf{H}[[X, Y]]$. From the second equation we get $d + b = iX + jY$, $db = -kXY$ so that we have

$$(iX + jY)\lambda = -(kX + Y).$$

Received by the editors September 4, 1980.

AMS (MOS) subject classifications (1970). Primary 15A63, 18F25.

Key words and phrases. Projective modules, quadratic spaces, extendibility.

© 1981 American Mathematical Society
0002-9939/81/0000-0502/\$01.50

If $\lambda = \lambda_0 + \lambda_1 X + \lambda_2 Y + \dots$, $\lambda_i \in \mathbf{H}$, we have $i\lambda_0 = -k$, $j\lambda_0 = -1$, a contradiction, which proves the proposition.

The reduced norm on P [1, Theorem 2.1] gives rise to a quadratic space of rank 4 and discriminant 1 over $R[T]$. This space is anisotropic and not extended from R (in fact indecomposable) in view of [1, Theorem 4.6].

REFERENCES

1. M. A. Knus, M. Ojanguren and R. Sridharan, *Quadratic forms and Azumaya algebras*, J. Reine Angew. Math. **303/304** (1978), 231–248.
2. T. Y. Lam, *Serre's conjecture*, Lecture Notes in Math., vol. 635, Springer-Verlag, Berlin and New York, 1978.
3. Raman Parimala, *Quadratic forms over polynomial rings over Dedekind domains*, Amer. J. Math. **100** (1978), 913–928.
4. S. Parimala and R. Sridharan, *Projective modules over polynomial rings over division rings*, J. Math. Kyoto Univ. **15** (1975), 129–148.

TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400 005, INDIA