

ON A THEOREM OF FLANDERS

ROBERT E. HARTWIG

ABSTRACT. It is shown that if R is a regular strongly- π -regular ring, then R is unit-regular precisely when $(ab)^d \approx (ba)^d$ for all $a, b \in R$. This generalizes a result by Flanders, which states that the matrices AB and BA over a field \mathbf{F} have the same elementary divisors except possibly those divisible by λ .

1. Introduction. A classic theorem of Flanders [3] states that if A and B are $n \times n$ matrices over a field \mathbf{F} , then AB and BA have the same elementary divisors, except possibly for those that are powers of λ .

The purpose of this note is to point out that the real reason why this result is true, is because the matrix ring $\mathbf{F}_{n \times n}$ is both strongly- π -regular as well as unit-regular. We shall use the concept of pseudo-similarity, introduced in [5] to provide the necessary link between strong- π -regularity and unit-regularity.

We recall that a ring R is called (unit) regular, if for every $a \in R$, there exists a (unit) solution $x \in R$, to the equation $axa = a$. Such solutions will be denoted at a^- . A ring R is strongly- π -regular, $s\pi r$ for short, if for every $a \in R$, there is a solution to the equations

$$a^k xa = a^k, \quad xax = x, \quad ax = xa,$$

for some $k \geq 0$. The solution is unique and is called the Drazin inverse a^d of a [2]. In the special cases where $k = 0$ or $k = 1$, a^d is called the group inverse of a , and is denoted by $a^\#$. It is well known that the ring $\mathbf{F}_{n \times n}$ of $n \times n$ matrices over a field \mathbf{F} is both $s\pi r$ and unit-regular. Two ring elements are called pseudo-similar if

$$x^-ax = b, \quad xbx^- = a, \quad xx^-x = x$$

for some $x, x^- \in R$. It was shown in [6], that for a unit-regular ring, similarity (\approx) and pseudo-similarity (\approx), coincide. Two idempotents e and f in R , are said to be equivalent, $e \sim f$, if eR and fR are isomorphic (\cong) as R -modules. This may be rewritten as $e \sim f$ if $e = p^+p$, $f = pp^+$, for some $p, p^+ \in R$, that satisfy $pp^+p = p$ and $p^+pp^+ = p^+$. It is easily seen that $e \sim f \Leftrightarrow e \approx f$ [4] and that \sim actually coincides with the classical \mathcal{D} -relation on semigroups.

2. Main results. Our generalization of the theorem of Flanders is based on the following two main results.

THEOREM 1. *Let R be a strongly- π -regular ring with unity and let $x, y \in R$. Then the following are equivalent:*

- (i) $x^d \approx y^d$,
- (ii) $x^2x^d \approx y^2y^d$.

In which case

Received by the editors July 10, 1980 and, in revised form, December 9, 1981.

1980 *Mathematics Subject Classification.* Primary 15A21, 15A09.

© 1982 American Mathematical Society
 0002-9939/81/0000-0789/\$01.75

(iii) $xx^d \approx yy^d$ and $(x^d + 1 - xx^d) \approx (y^d + 1 - yy^d)$.

If in addition $R = \mathbb{F}_{n \times n}$, then each of the above is equivalent to

(iv) x and y have the same elementary divisors, except possibly those that are powers of λ .

PROOF. (i) \Leftrightarrow (ii) This is based on a result of Drazin [2], which states that if $pq = qp$, and $p^\#$ exists, then $p^\#q = qp^\#$. Now if $x^d q = qy^d$, with q invertible, then because $(x^d)^\# = x^2 x^d$, it follows that $x^2 x^d q = qy^2 y^d$. The converse follows also since $(p^\#)^\# = p$.

(i) \Rightarrow (iii) Let $x^d q = qy^d$, and hence by part (ii) $x^2 x^d q = qy^2 y^d$. Now $xx^d = (x^2 x^d)x^d$ and so $xx^d q = qyy^d$. The remaining result is now also clear.

(iii) \Rightarrow (iv) Let $R = \mathbb{F}_{n \times n}$, and suppose that

$$(2.1) \quad x = P \begin{bmatrix} U_1 & 0 \\ 0 & \eta_1 \end{bmatrix} P^{-1}, \quad y = Q \begin{bmatrix} U_2 & 0 \\ 0 & \eta_2 \end{bmatrix} Q^{-1},$$

are the core-nilpotent decompositions of x and y respectively, with U_i invertible and η_i nilpotent, $i = 1, 2$. Then

$$x^d = P \begin{bmatrix} U_1^{-1} & 0 \\ 0 & 0 \end{bmatrix} P^{-1},$$

and $xx^d \approx yy^d$ shows that U_1 and U_2 have the same size. Next,

$$\begin{bmatrix} U_1^{-1} & 0 \\ 0 & I \end{bmatrix} \approx x^d + 1 - xx^d \approx y^d + 1 - yy^d \approx \begin{bmatrix} U_2^{-1} & 0 \\ 0 & I \end{bmatrix}.$$

Hence, on using elementary divisors, it follows that $U_1^{-1} \approx U_2^{-1}$ and so $U_1 \approx U_2$. This is equivalent to (iv).

(iv) \Rightarrow (i) This is clear from the fact that $x^d \approx y^d \Leftrightarrow U_1^{-1} \approx U_2^{-1} \Leftrightarrow U_1 \approx U_2$. Our second result deals with the relation between ab and ba .

THEOREM 2. *Let R be a strongly- π -regular ring with unity, and let $a, b \in R$. Then the following are equivalent:*

- (i) R is unit-regular, and
- (ii) R is regular and $(ab)^d \approx (ba)^d$ for all $a, b \in R$.

PROOF. (i) \Rightarrow (ii) Cline's formula [1], states that $(xy)^d = x(yx)^{d^2}y$. If we apply this to ab and ba we may write

$$(ab)^d = a \cdot (ba)^d \cdot (ba)^d b$$

and

$$(ba)^d = (ba)^d b \cdot (ab)^d \cdot a.$$

Now since $(ba)^d b \cdot a \cdot (ba)^d b = (ba)^d b$, it follows that $(ab)^d \approx (ba)^d$. Moreover, $ab(ab)^d \approx ba(ba)^d$ as well. Next, since R is assumed to be unit-regular, we may conclude [6, p. 453] that semisimilarity implies similarity. Consequently $(ab)^d \approx (ba)^d$ and $ab(ab)^d \approx ba(ba)^d$.

(ii) \Rightarrow (i) Based on a result of Vidav [7], it suffices to show that two equivalent idempotents are similar. Suppose therefore that e and f are equivalent idempotents in R , and that $e = p^+ p f = p p^+$, with $p p^+ p = p$ and $p^+ p p^+ = p^+$. Now since $e^d = e$ and $f^d = f$, we have $e = e^d = (p^+ p)^d \approx (p p^+)^d = f^d = f$. Because R is regular, the proof is now complete.

COROLLARY 1. If $A, B \in \mathbf{F}_{n \times n}$, then

- (i) $(AB)^d \approx (BA)^d$,
- (ii) $(AB)^2(AB)^d \approx (BA)^2(BA)^d$, and
- (iii) the elementary divisors of AB and BA coincide except possibly for those that are powers of λ .

PROOF. (i) Since $\mathbf{F}_{n \times n}$ is unit-regular and $s\pi r$, this follows from Theorem 2. The remaining two results follow at once from Theorem 1, thereby completing the proof of Flander's Theorem.

COROLLARY 2. If $A, B \in \mathbf{F}_{n \times n}$, then $AB \approx BA$ if and only if $\text{rank}(AB)^k = \text{rank}(BA)^k$, $k = 1, 2, \dots$.

PROOF. The necessity is clear. For the sufficiency, suppose that $AB = x$ and $BA = y$ are given as in (2.1). It then suffices to show that $U_1 \approx U_2$ and $\eta_1 \approx \eta_2$. From Corollary 1, we recall that $U_1 \approx U_2$ always holds. Lastly, $\text{rank}(AB)^k = \text{rank}(BA)^k$ implies that $\text{rank}(\eta_1^k) = \text{rank}(\eta_2^k)$, for all $k = 1, 2, \dots$, which is well known to suffice for $\eta_1 \approx \eta_2$.

REMARKS. 1. The results of Corollary 1 can easily be modified to include the case where A and B are rectangular, by means of adding zeros suitably.

2. The equivalence of the pseudo-similarity and similarity of $(ab)^d$ and $(ba)^d$, hinges on the existence of a unit solution to the equation $(ba)^d b \cdot x \cdot (ba)^d b = (ba)^d b$. Even though several obvious solutions exist, such as $x = a$ or $x = a + 1 - (ba)(ba)^d$, it is not known whether there exists a unit solution u that can be expressed in terms of a, b and $(\cdot)^d$ exclusively.

3. It is not known whether the converse of Theorem 1 holds in general. That is whether $xx^d \approx yy^d$ and $x^d + 1 - xx^d \approx y^d + 1 - yy^d$ ensure that $x^d \approx y^d$.

REFERENCES

1. R. E. Cline, *An application of representations for the generalized inverse of a matrix*, MRC Technical report, #592, 1965.
2. M. P. Drazin, *Pseudo-inverses in associative rings and semigroups*, Amer. Math. Monthly **65** (1958), 506-514.
3. H. Flanders, *Elementary divisors of AB and BA* , Proc. Amer. Math. Soc. **2** (1951), 871-874.
4. R. E. Hartwig, *Unit and unitary solutions to $AX = B$* (submitted).
5. R. E. Hartwig and F. Hall, *Pseudo-similarity for matrices over a field*, Proc. Amer. Math. Soc. **71** (1978), 6-10.
6. R. E. Hartwig and J. Luh, *A note on the group structure of unit regular ring elements*, Pacific J. Math. **71** (1977), 449-461.
7. I. Vidav, *Modules over regular rings*, Math. Balkanica **1** (1971), 287-292.

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NORTH CAROLINA 27650