

ANOTHER CHARACTERIZATION OF BLO

COLIN BENNETT¹

ABSTRACT. It is shown that a locally integrable function f on \mathbb{R}^n has bounded lower oscillation ($f \in \text{BLO}$) if and only if $f = MF + h$, where F has bounded mean oscillation ($F \in \text{BMO}$) and $MF < \infty$ a.e., and h is bounded. Here, MF is a variant of the familiar Hardy-Littlewood maximal function: $MF = \sup_{Q \ni x} Q(F)$ (no absolute values), where $Q(F)$ is the mean value of F over the cube Q .

We consider real-valued locally integrable functions f on \mathbb{R}^n . When Q is a cube in \mathbb{R}^n with sides parallel to the coordinate axes, we denote by $Q(f)$ the mean value of f over Q . Define a maximal function Mf of f by

$$(Mf)(x) = \sup_{Q \ni x} Q(f) = \sup_{Q \ni x} \frac{1}{|Q|} \int_Q f(y) dy \quad (x \in \mathbb{R}^n),$$

where the supremum extends over all cubes Q that contain x . Note that f is not assumed to be nonnegative so that Mf may take on negative values. However, it follows from the differentiation theorem that $Mf \geq f$ a.e. Moreover, it is clear $|Mf|$ is dominated by $M|f|$, the latter being the familiar Hardy-Littlewood maximal function of f , and so, in particular, the maximal operator M is bounded on L^2 . We shall make use of the obvious fact that if Q and Q' are two cubes with $Q \subset Q'$, then $Q'(f)$ is dominated by $(Mf)(x)$ for every x in Q , and so

$$(1) \quad Q'(f) \leq \inf_Q Mf \quad (Q \subset Q').$$

A locally integrable function f on \mathbb{R}^n is said to be of *bounded mean oscillation* ($f \in \text{BMO}$) if

$$(2) \quad \|f\|_{\text{BMO}} = \sup_Q Q(|f - Q(f)|)$$

is finite. The space BMO is a linear space and, when the constant functions are factored out, the functional in (2) is a norm under which BMO is a Banach space. It is well known (cf. [3, p. 227]) that replacing (2) by the corresponding quadratic functional results in an equivalent norm on BMO. In particular, there is a constant c , depending only on the dimension n , such that

$$(3) \quad \left(\frac{1}{|Q|} \int_Q |f(x) - Q(f)|^2 dx \right)^{1/2} \leq c \|f\|_{\text{BMO}},$$

for every cube Q in \mathbb{R}^n .

Received by the editors July 30, 1981.

AMS (MOS) subject classifications (1970). Primary 44A25.

¹Research supported in part by NSF Grant MCS-8102194.

©1982 American Mathematical Society
 0002-9939/81/0000-0799/\$02.25

Recently, R. R. Coifman and R. Rochberg [2] have introduced the space BLO of functions of *bounded lower oscillation*. The space is defined analogously to BMO except that in (2) one subtracts from f the essential infimum

$$f_Q = \operatorname{ess\,inf}_{x \in Q} f(x)$$

instead of the mean value $Q(f)$. Thus, a locally integrable function f on \mathbb{R}^n is in BLO if

$$(4) \quad \|f\|_{\text{BLO}} = \sup_Q (Q(f) - f_Q)$$

is finite. The BLO property need not be preserved under multiplication by negative constants so, in particular, BLO is not a linear space, and, despite the notation, the functional in (4) is not a norm. It is easy to check, however, that $\|\cdot\|_{\text{BLO}}$ is subadditive and positive-homogeneous.

Note that every L^∞ -function is in BLO, with

$$(5) \quad \|f\|_{\text{BLO}} \leq 2\|f\|_{L^\infty} \quad (f \in L^\infty),$$

and every BLO-function is in BMO, with

$$(6) \quad \|f\|_{\text{BMO}} \leq 2\|f\|_{\text{BLO}} \quad (f \in \text{BLO}).$$

Using a representation theorem of Carleson for BMO-functions, Coifman and Rochberg [2] have shown that every BMO-function is representable as the difference of two BLO-functions. Moreover, they have shown that BLO-functions arise, modulo bounded functions, as logarithms of maximal functions. In other words, a locally integrable function f belongs to BLO if and only if

$$(7) \quad f = \alpha \log MF + h,$$

where α is a nonnegative constant, F is a nonnegative locally integrable function whose maximal function MF is finite a.e., and h is an L^∞ -function.

Let us note also the result of [1] that if $f \in \text{BMO}$ (and has finite maximal function), then the decreasing rearrangement f^* (of $|f|$) is in BLO; in fact, this condition characterizes the rearrangements of BMO-functions.

In this note, we shall obtain a different description of BLO-functions. We shall show that they arise, modulo bounded functions, as the maximal functions MF of BMO-functions F . Of course, we must exclude those F for which MF is identically infinite.

We shall need two lemmas, the first of which may be regarded as a refinement of Theorem 4.2 in [1].

LEMMA 1. *If $F \in \text{BMO}$ and if Q is any cube in \mathbb{R}^n , then*

$$(8) \quad Q(MF) \leq c\|F\|_{\text{BMO}} + \inf_Q MF,$$

where c is a constant depending on the dimension n . In particular, if MF is not identically infinite, then $MF \in \text{BLO}$ and

$$(9) \quad \|MF\|_{\text{BLO}} \leq c\|F\|_{\text{BMO}}.$$

PROOF. Fix Q and let \bar{Q} be the cube concentric to Q with dimensions three times as large. Write

$$F = (F - \bar{Q}(F))\chi_{\bar{Q}} + [\bar{Q}(F)\chi_{\bar{Q}} + F\chi_{\bar{Q}^c}] = G + H,$$

say.

By the Cauchy-Schwarz inequality,

$$Q(MG) \leq \left(\frac{1}{|Q|} \int_Q (MG)^2 dx \right)^{1/2} \leq |Q|^{-1/2} \|MG\|_{L^2(\mathbb{R}^n)}.$$

But M is bounded on L^2 , so using the definition of G , the fact that $|\bar{Q}| = 3^n |Q|$, and (3), we obtain

$$(10) \quad Q(MG) \leq c\|F\|_{\text{BMO}}.$$

Next we shall show that

$$(11) \quad Q(MH) \leq c\|F\|_{\text{BMO}} + \inf_Q MF,$$

which, together with (10) and the fact that $F = G + H$, will establish the desired result (8). In order to establish (11), it will suffice to show that $MH(x)$ is dominated by the right-side of (11) for every $x \in Q$, and for this it will be enough to show that

$$(12) \quad P(H) \leq c\|F\|_{\text{BMO}} + \inf_Q MF$$

for every cube P in \mathbb{R}^n containing x .

The result follows directly from (1) if P does not meet \bar{Q}^c , for then $H = \bar{Q}(F)$ on P and so $P(H) = \bar{Q}(F) \leq \inf_Q MF$. So suppose $P \cap \bar{Q}^c \neq \emptyset$, and let P' be the smallest cube containing both P and \bar{Q} . Since P contains the point x of Q , it is clear that

$$(13) \quad |P'| \leq c|P|$$

for some constant c depending only on the dimension n . Furthermore, from the definition of H ,

$$\begin{aligned} \int_P (H - P'(F)) &\leq \int_{P'} |H - P'(F)| = \int_{\bar{Q}} |\bar{Q}(F) - P'(F)| + \int_{P' \cap \bar{Q}^c} |F - P'(F)| \\ &\leq \left(\int_{\bar{Q}} + \int_{P' \cap \bar{Q}^c} \right) |F - P'(F)| = \int_{P'} |F - P'(F)|, \end{aligned}$$

so, using (2) and (13), we obtain $P(H - P'(F)) \leq c\|F\|_{\text{BMO}}$. Since $P' \supset Q$, an appeal to (1) now gives

$$P(H) = P(H - P'(F)) + P'(F) \leq c\|F\|_{\text{BMO}} + \inf_Q MF.$$

This establishes (12) and hence, as we remarked above, completes the proof of (8).

It is now clear from (8) that if $(MF)(x)$ is finite at any one point x in \mathbb{R}^n , then $(MF)(y)$ is finite a.e. on every cube Q containing x , hence on all of \mathbb{R}^n . In that case, we may subtract $\inf_Q MF$ from each side of (8) and take the supremum over all Q to obtain (9).

We require one further lemma.

LEMMA 2. *A locally integrable function f on \mathbb{R}^n belongs to BLO if and only if $Mf - f$ belongs to L^∞ . Furthermore,*

$$(14) \quad \|Mf - f\|_{L^\infty} = \|f\|_{\text{BLO}}.$$

PROOF. Suppose first that f belongs to BLO. Let x be any Lebesgue point of f and let Q be any cube containing x . Then $f(x) \geq f_Q$ and so

$$Q(f) - f(x) \leq Q(f) - f_Q \leq \|f\|_{\text{BLO}}.$$

Taking the supremum over all cubes Q containing x , and then the supremum over all Lebesgue points x of f , we find that $Mf - f$ belongs to L^∞ and

$$(15) \quad \|Mf - f\|_{L^\infty} \leq \|f\|_{\text{BLO}}.$$

Conversely, suppose $Mf - f$ belongs to L^∞ and let Q be any cube in \mathbb{R}^n . Any point x in Q for which

$$(16) \quad f(x) < Q(f) - \|Mf - f\|_{L^\infty}$$

must satisfy

$$(Mf)(x) - f(x) \geq Q(f) - f(x) > \|Mf - f\|_{L^\infty},$$

and consequently such points x constitute a set of measure zero. Hence the essential infimum f_Q is at least as large as the value on the right-hand side of (16), and so

$$Q(f) - f_Q \leq \|Mf - f\|_{L^\infty}.$$

Taking the supremum over all Q , we obtain the reverse inequality to (15), and hence (14) is established.

Our main result is as follows.

THEOREM. *A locally integrable function f on \mathbb{R}^n belongs to BLO if and only if there are functions h in L^∞ and F in BMO with MF finite a.e. such that*

$$(17) \quad f = MF + h.$$

Furthermore,

$$(18) \quad \|f\|_{\text{BLO}} \sim \inf(\|F\|_{\text{BMO}} + \|h\|_{L^\infty}),$$

where the infimum extends over all representations of the form (17).

PROOF. If f has a representation as in (17), then MF belongs to BLO by virtue of Lemma 1. Since h is bounded, hence in BLO, we see that f is in BLO. Furthermore, from (5) and (9),

$$\|f\|_{\text{BLO}} \leq \|MF\|_{\text{BLO}} + \|h\|_{\text{BLO}} \leq c\|F\|_{\text{BMO}} + 2\|h\|_{L^\infty},$$

so taking the infimum over all representations (17) of f , we obtain

$$(19) \quad \|f\|_{\text{BLO}} \leq c \inf(\|F\|_{\text{BMO}} + \|h\|_{L^\infty}).$$

Conversely, if $f \in \text{BLO}$, then Mf is finite a.e., by Lemma 2. Furthermore, by the same lemma, the function $f - Mf$ is bounded, and so $f = Mf + (f - Mf)$ is a

representation of the form (17) with $F = f$ and $h = f - Mf$. Moreover, from (6) and (14),

$$\|F\|_{\text{BMO}} + \|h\|_{L^\infty} = \|f\|_{\text{BMO}} + \|f - Mf\|_{L^\infty} \leq 3\|f\|_{\text{BLO}}$$

so, combining this observation with (19), we establish (18).

REMARKS. The above analysis can also be carried through for a cube Q_0 instead of all of \mathbb{R}^n . In this case, BLO-functions on Q_0 are bounded below and so, by addition of suitable constants, can be rendered nonnegative. In this case, one obtains a result of the form (17) with M the usual Hardy-Littlewood maximal function $\sup_Q Q(|f|)$.

I wish to thank my colleagues, Professors R. A. DeVore, W. Nestlerode, and R. Sharpley, for some interesting discussions concerning this material.

REFERENCES

1. C. Bennett, R. A. DeVore and R. Sharpley, *Weak- L^∞ and BMO*, Ann. of Math. (2) **113** (1981), 601–611.
2. R. R. Coifman and R. Rochberg, *Another characterization of BMO*, Proc. Amer. Math. Soc. **79** (1980), 249–254.
3. C. Sadosky, *Interpolation of operators and singular integrals*, Dekker, New York, 1979.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SOUTH CAROLINA 29208