

## SYMMETRIC CONTINUITY OF REAL FUNCTIONS

C. L. BELNA

**ABSTRACT.** It is shown that the set of points where a real function is both symmetrically continuous and not continuous has inner measure zero but may have full outer measure.

We shall consider an arbitrary function  $f$  that maps the real line  $\mathbf{R}$  into itself. Such a function is said to be *symmetrically continuous at*  $x \in \mathbf{R}$  if

$$f(x+h) - f(x-h) \rightarrow 0 \quad \text{as } h \rightarrow 0.$$

The set of all points where  $f$  is symmetrically continuous is denoted by  $\mathcal{S}\mathcal{C}$ , and the set of all points where  $f$  is discontinuous is denoted by  $\mathcal{D}$ . (Recall that  $\mathcal{D}$  is of type  $F_\sigma$ ; on the other hand,  $\mathcal{S}\mathcal{C}$  need not be measurable and need not possess the Baire property—examples are given below.)

The object of this investigation is the set  $\mathcal{S}\mathcal{C} \cap \mathcal{D}$ . Interest in this set apparently dates back to 1935 when F. Hausdorff [3] asked whether  $\mathcal{D}$  could be uncountable when  $\mathcal{S}\mathcal{C} = \mathbf{R}$ ; an affirmative answer to this question was given in 1971 by D. Preiss [4]. Hausdorff also asked whether an arbitrarily prescribed  $F_\sigma$ -set could equal  $\mathcal{D}$  when  $\mathcal{S}\mathcal{C} = \mathbf{R}$ . In response to this question, H. Fried [2] proved the following result.

**THEOREM F.** *If  $\mathcal{S}\mathcal{C}$  is residual, then  $\mathcal{D}$  is of first category.*

The analogue for  $f$  restricted to an interval is clearly valid; and because  $\mathcal{D}$  has the Baire property and any residual subset of an open interval has the Baire property, this theorem has the following equivalent formulations.

**THEOREM F\*.**  *$\mathcal{S}\mathcal{C} \cap \mathcal{D}$  is residual on no open interval; or, equivalently,  $\mathcal{S}\mathcal{C} \cap \mathcal{D}$  contains no set of second category that has the Baire property.*

In the following sense, Theorem F\* is the “best” statement that can be made concerning the category of  $\mathcal{S}\mathcal{C} \cap \mathcal{D}$ . P. Erdős [1] has shown that the continuum hypothesis implies the existence of an additive subgroup  $G$  of  $\mathbf{R}$  that is of measure 0 and of second category on every open interval. Thus, if  $f$  is the characteristic function of  $G$ , then  $G \subset \mathcal{S}\mathcal{C}$  and  $\mathcal{D} = \mathbf{R}$ ; that is,  $\mathcal{S}\mathcal{C} \cap \mathcal{D}$  is of second category on every open interval.

Erdős also proved that the continuum hypothesis implies the existence of a subgroup  $G$  of  $\mathbf{R}$  that is of first category and has full exterior measure on each open

---

Received by the editors March 12, 1982.  
1980 *Mathematics Subject Classification.* Primary 26A15.

interval. Thus, if  $f$  is the characteristic function of  $G$ , then

$$|(\mathfrak{S}\mathcal{C} \cap \mathfrak{O}) \cap I|_e = |I|$$

for each open interval  $I$  (where  $|\cdot|_e$  and  $|\cdot|$  denote exterior Lebesgue measure and Lebesgue measure). Consequently, the following metric analogue of Theorem F\* is the “best” statement that can be made concerning the measure of  $\mathfrak{S}\mathcal{C} \cap \mathfrak{O}$ . (Here  $|\cdot|_i$  denotes inner Lebesgue measure.)

**THEOREM.**  $|\mathfrak{S}\mathcal{C} \cap \mathfrak{O}|_i = 0$ .

It is our goal to furnish a proof of this theorem. First we note an immediate corollary: *if  $\mathfrak{S}\mathcal{C}$  is measurable, then  $|\mathfrak{S}\mathcal{C} \cap \mathfrak{O}| = 0$ ; in particular, if  $\mathfrak{S}\mathcal{C}$  is of full measure, then  $|\mathfrak{O}| = 0$ . This contains a previous result of Preiss [4], who proved that  $\mathfrak{S}\mathcal{C} = \mathbf{R}$  implies  $|\mathfrak{O}| = 0$ . We also note that E. M. Stein and A. Zygmund [5, Lemma 9] presented our theorem with the added hypothesis that  $f$  be measurable. However, their proof contains an oversight; for example, if  $f(x) = x$ , then in their notation  $E = \mathbf{R}$  and  $E_n = \emptyset$  for each  $n$ . (In a footnote [5, p. 266], they do allude to a second proof which may be correct.)*

To begin the proof, we introduce an auxiliary function of two variables and prove two key lemmas concerning it. For each  $x \in \mathbf{R}$  and each  $\epsilon > 0$  we define  $\delta(x; \epsilon)$  to be the supremum of all numbers  $\delta \geq 0$  for which

$$|f(x + h) - f(x - h)| < \epsilon \quad \text{whenever } |h| \leq \delta.$$

(Note that  $\delta(x; \epsilon) \geq 0$ , and that  $\delta(x; \epsilon) > 0$  for each  $\epsilon > 0$  if and only if  $x \in \mathfrak{S}\mathcal{C}$ .)

**LEMMA 1.** *Let  $\epsilon$  and  $\delta$  be positive numbers, and let  $I$  be an open interval with  $|I| < \delta/2$ . If  $|\{x: \delta(x; \epsilon) > \delta\} \cap I|_e > 0$ , then there exists a positive number  $\tau$  such that  $\delta(x; 5\epsilon) > \tau$  for each  $x \in \mathfrak{S}\mathcal{C} \cap I$ .*

**PROOF.** Set  $E = \{x: \delta(x; \epsilon) > \delta\} \cap I$ . Since  $|E|_e > 0$ , the distance set

$$d(E) \equiv \{|x - y| : x, y \in E\}$$

is dense on some interval  $(0, \bar{\tau})$ . (This follows from the identity  $\overline{d(E)} = d(\bar{E})$  and the well-known fact that the distance set of a set of positive measure contains some interval  $(0, \bar{\tau})$ .) Choose any point  $x \in \mathfrak{S}\mathcal{C} \cap I$ , and without loss of generality assume  $x = 0$ . Let  $h \in (0, \bar{\tau})$  be arbitrary. To establish the lemma for any  $\tau \in (0, \bar{\tau})$ , we shall show that

$$(1) \quad |f(h) - f(-h)| < 5\epsilon.$$

Inequality (1) is certainly satisfied if  $\delta(0; \epsilon) > h$ . Suppose  $\delta(0; \epsilon) \leq h$ . Because  $d(E)$  is dense on  $(0, \bar{\tau})$ , there exist two points  $x, y \in E$  such that  $x < y$  that

$$(2) \quad 0 < h - 2(y - x) < \delta(0; \epsilon).$$

Let  $h_y$  be the reflection of  $h$  in  $y$ , and let  $h_{yx}$  be the reflection of  $h_y$  in  $x$ . Since  $\tau \leq |I| < \delta/2$ , it readily follows that

$$|h - y| < |I| + \tau < \delta \quad \text{and} \quad |h_y - x| < |I| + \tau < \delta.$$

Thus, since  $x, y \in E$  we have

$$|f(h) - f(h_y)| < \epsilon \quad \text{and} \quad |f(h_y) - f(h_{yx})| < \epsilon;$$

that is,

$$(3) \quad |f(h) - f(h_{yx})| < 2\epsilon.$$

A similar argument shows that

$$(4) \quad |f(-h) - f((-h)_{xy})| < 2\epsilon,$$

where  $(-h)_{xy}$  is the reflection in  $y$  of the reflection of  $-h$  in  $x$ . Furthermore,

$$(-h)_{xy} = -h + 2(y - x) = -h_{yx},$$

and by (2) we have  $0 < h_{yx} < \delta(0; \epsilon)$ ; therefore,

$$|f(h_{yx}) - f((-h)_{xy})| < \epsilon.$$

This together with (3) and (4) and the triangle inequality gives (1), and the proof is complete.

**LEMMA 2.** *Let  $\epsilon$  and  $\delta$  be positive numbers, and let  $I$  be an open interval with  $|I| < \delta$ . If  $|\{x: \delta(x; \epsilon) > \delta\} \cap I|_i > 3|I|/4$ , then  $|f(a) - f(b)| < 2\epsilon$  for any two points  $a, b \in I$ .*

**PROOF.** If  $E = \{x: \delta(x; \epsilon) > \delta\} \cap I$ , then it readily follows from the hypothesis on  $E$  that  $(0, |I|/2) \subset d(E)$ . Thus if  $a, b \in I$  with  $a < b$ , there exist points  $x, y \in E$  such that  $x < y$  and  $b - a = 2(y - x)$ . Let  $a_x$  denote the reflection of  $a$  in  $x$ , and let  $a_{xy}$  denote the reflection of  $a_x$  in  $y$ . Because  $|I| < \delta$  and  $a_{xy} = b$  we have  $|a - x| < \delta$  and  $|a_x - y| < \delta$ ; and because  $x, y \in E$  we have

$$|f(a) - f(a_x)| < \epsilon \quad \text{and} \quad |f(a_x) - f(a_{xy})| < \epsilon.$$

As  $a_{xy} = b$ , this implies that  $|f(a) - f(b)| < 2\epsilon$ , and the lemma is established.

**PROOF OF THEOREM.** Assume  $|\mathfrak{S}\mathcal{C} \cap \mathfrak{O}|_i > 0$ . If  $\mathfrak{O}_t$  denotes the set of points where the oscillation of  $f$  exceeds  $t$ , then  $|\mathfrak{S}\mathcal{C} \cap \mathfrak{O}_t|_i > 0$  for some  $t > 0$  (since each  $\mathfrak{O}_t$  is measurable). Let  $M$  be a measurable subset of  $\mathfrak{S}\mathcal{C} \cap \mathfrak{O}_t$  with  $|M| > 0$ . Choose any  $\epsilon \in (0, t/11)$ . Since  $M \subset \mathfrak{S}\mathcal{C}$ , there exists a  $\delta > 0$  such that

$$|\{x: \delta(x, \epsilon) > \delta\} \cap M|_e > 0.$$

Therefore, there exists a point  $p \in \{x: \delta(x; \epsilon) > \delta\} \cap M$  that is a point of density of  $M$  and a point of exterior density of the set  $\{x: \delta(x; \epsilon) \geq \delta\}$ . Let  $I$  be an open interval containing  $p$  with  $|I| < \delta/2$ . By Lemma 1 there exists a  $\tau > 0$  such that  $\delta(x; 5\epsilon) > \tau$  for each  $x \in M$ . Let  $J$  be an open interval containing  $p$  such that  $|J| < \tau$  and  $|M \cap J| > 3|J|/4$ . Then

$$|\{x: \delta(x; 5\epsilon) > \tau\} \cap J|_i > 3|J|/4,$$

and by Lemma 2 the oscillation of  $f$  on  $J$  does not exceed  $10\epsilon$  which is less than  $10t/11$ . This contradicts  $p \in \mathfrak{O}_t$ , and the theorem is proved.

The author wishes to thank M. J. Evans for bringing reference [5] to his attention.

#### REFERENCES

1. P. Erdős, *Some remarks on subgroups of real numbers*, Colloq. Math. **42** (1979), 119–120.
2. H. Fried, *Über die symmetrische Stetigkeit von Funktionen*, Fund. Math. **29** (1937), 134–137.
3. F. Hausdorff, *Problème #62*, Fund. Math. **25** (1935), 578.
4. D. Preiss, *A note on symmetrically continuous functions*, Časopis Pěst. Mat. **96** (1971), 262–264.
5. E. M. Stein and A. Zygmund, *On the differentiability of functions*, Studia Math. **23** (1964), 247–283.

SYRACUSE RESEARCH CORPORATION, MERRILL LANE, SYRACUSE, NEW YORK 13210