

A THEOREM OF CRAMÉR AND WOLD REVISITED

ALLADI SITARAM

ABSTRACT. Let $H = \{(x, y) : x > 0\} \subseteq \mathbf{R}^2$ and let E be a Borel subset of H of positive Lebesgue measure. We prove that if μ and ν are two probability measures on \mathbf{R}^2 such that $\mu(\sigma(E)) = \nu(\sigma(E))$ for all rigid motions σ of \mathbf{R}^2 , then $\mu = \nu$. This generalizes a well-known theorem of Cramér and Wold.

1. Introduction. A celebrated theorem of H. Cramér and H. Wold particularly well known to probabilists (see [3]) asserts: If μ and ν are probability measures on \mathbf{R}^2 such that they agree on all half planes, then $\mu = \nu$. This can be reformulated in the following way: Let $H = \{(x, y) \in \mathbf{R}^2; x \geq 0\}$. If μ and ν are probability measures such that $\mu(\sigma(H)) = \nu(\sigma(H))$ for all rigid motions σ of \mathbf{R}^2 , then $\mu = \nu$. The aim of this note is to generalize this result to an arbitrary Borel set E of positive Lebesgue measure contained in H . The results in this paper are valid for \mathbf{R}^n , $n \geq 2$, but for notational simplicity we consider \mathbf{R}^2 —the same proofs go through for any $n \geq 2$. A special case of our result—for a restricted class of Borel sets E , i.e. those that “pave” the half space H —appears in [4, §I.2.4]. However, the methods in [4] are different, where the Radon transform is used.

2. Notation and terminology. For any unexplained notation or terminology please see [5].

By a rigid motion of \mathbf{R}^2 we mean a homeomorphism of \mathbf{R}^2 of the form $(x, y) \rightarrow T(x, y) + (x_0, y_0)$, where (x_0, y_0) is a fixed vector in \mathbf{R}^2 and T is a special orthogonal linear transformation of \mathbf{R}^2 (i.e. T is a matrix of the form

$$\begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}.$$

Throughout this paper λ denotes the Lebesgue measure on \mathbf{R}^2 . Let C denote the class of all (finite) complex measures on \mathbf{R}^2 . If T is a tempered distribution (in the sense of Schwartz), then \hat{T} denotes the Fourier transform of T (which is again a distribution) and $\text{Supp } T$ denotes the (closed) support of T . For standard facts regarding distributions, Fourier transforms etc., see [5]. If g is a bounded Borel function on \mathbf{R}^2 , then g defines a tempered distribution and \hat{g} will denote the (distributional) Fourier transform of g . If μ is a finite complex measure, then $\mu * g$ is the bounded Borel function defined by

$$(\mu * g)(x) = \int_{\mathbf{R}^2} g(x - y) d\mu(y).$$

Received by the editors September 11, 1981 and, in revised form, July 21, 1982.
1980 *Mathematics Subject Classification*. Primary 60B15, 60E10.

© 1983 American Mathematical Society
0002-9939/82/0000-00787/\$01.50

Finally we note that for a complex measure or an L^1 -function the usual notion of Fourier transform coincides with the notion of distributional Fourier transform.

If $E \subset \mathbf{R}^2$, let 1_E denote the indicator function of E , i.e. $1_E(x) = 1$ if $x \in E$ and $1_E(x) = 0$ if $x \notin E$.

H will always stand for the subset of \mathbf{R}^2 defined by $H = \{(x, y) \in \mathbf{R}^2; x \geq 0\}$.

We end this section by quoting a result that will be needed in the next section.

PROPOSITION. *Let f be a bounded measurable function on \mathbf{R}^2 and $\mu \in C$. If $\mu * f$ vanishes identically, then $\hat{\mu}$ vanishes on $\text{Supp } \hat{f}$.*

(*Note.* For a proof of this theorem, we refer to p. 232 of [2]. In [2] μ is taken to be an L^1 -function but by convolving μ with an L^1 -function whose Fourier transform is nowhere vanishing (e.g. the Gaussian) we can get the theorem quoted above. Note also that $\text{Supp } \hat{f}$ is called “spectrum of f ” in [2].)

3. The main result. We start with a proposition which combined with the proposition quoted in §2 yields the main result.

PROPOSITION 3.1. *Let h be a nonnegative bounded Borel measurable function on \mathbf{R}^2 such that h is positive on a set of positive Lebesgue measure and such that $\text{Supp } h \subset H$. Then $\mathbf{R} \times \{0\} \subset \text{Supp } \hat{h}$.*

PROOF. First assume $h \in L^1(\mathbf{R}^2)$. Let $f(x) = \int_{\mathbf{R}} h(x, y) dy$. The hypotheses on h easily imply that f is a nontrivial, nonnegative L^1 -function on \mathbf{R} which is supported in \mathbf{R}^+ . Now if \hat{f} is the one-dimensional Fourier transform of f , then \hat{f} can be extended to a bounded function g in the region $S = \{z \in \mathbf{C}; \text{Im } z \leq 0\}$. g will be analytic in $S_0 = \{z \in \mathbf{C}; \text{Im } z < 0\}$ and continuous in S . Thus \hat{f} is the “boundary-value” of a bounded analytic function in S_0 and consequently \hat{f} cannot vanish identically on any nonempty open subset of \mathbf{R} , i.e. $\text{Supp } \hat{f} = \mathbf{R}$. Now observe that if \hat{h} is the (two dimensional) Fourier transform of h , then

$$\hat{h}(\lambda, 0) = \int_{\mathbf{R}^2} h(x, y) e^{-i\lambda x} dx dy = \hat{f}(\lambda).$$

Thus $\text{Supp } \hat{h} \supset \mathbf{R} \times \{0\}$ because $\text{Supp } \hat{f} = \mathbf{R}$.

Now we drop the assumption that $h \in L^1(\mathbf{R}^2)$. To prove the proposition let us assume \hat{h} vanishes in a neighborhood U (in \mathbf{R}^2) of a point $(\lambda_0, 0) \in \mathbf{R} \times \{0\}$. Choose ϵ sufficiently small such that the open ball of radius 2ϵ with centre at $(\lambda_0, 0)$ is contained in U . Let $0 \neq h_1$ be a nonnegative function in $L^1(\mathbf{R}^2)$ such that \hat{h}_1 is a C^∞ -function and $\text{Supp } \hat{h}_1$ is contained in the ball of radius ϵ around 0. (It is always possible to do this.) Then $\text{Supp}(\hat{h}h_1) = \text{Supp}(\hat{h} * \hat{h}_1) \subset \text{Supp } \hat{h} + \text{Supp } \hat{h}_1$. So if $U' = \{(x, y) \in \mathbf{R}^2; \sqrt{(x - \lambda_0)^2 + y^2} < \epsilon\}$, then $\text{Supp}(\hat{h}h_1) \cap U' = \emptyset$. However hh_1 is a nonnegative L^1 -function with $\text{Supp } hh_1 \subseteq H$ and by the first part of our proof hh_1 must be zero almost everywhere on \mathbf{R}^2 . Since \hat{h}_1 is a C^∞ -function of compact support, h_1 is the restriction of an entire function to \mathbf{R}^2 and hence $h_1(x) \neq 0$ a.e. on \mathbf{R}^2 . Thus h is zero a.e. which gives us a contradiction and the proof of our proposition is complete.

Proposition 3.1 and the Proposition in §2 easily imply the following generalization of the Cramér-Wold theorem.

THEOREM 3.2. *Let E be a Borel subset of H such that $\lambda(E) > 0$. Let $\mu, \nu \in C$ such that $\mu(\sigma(E)) = \nu(\sigma(E))$ for all rigid motions σ of \mathbf{R}^2 . Then $\mu = \nu$.*

PROOF. Let l be any line through $(0,0)$ in \mathbf{R}^2 . We will prove $\hat{\mu} = \hat{\nu}$ on l . By Proposition 3.1, $\text{Supp } \hat{1}_E \supseteq \mathbf{R} \times \{0\}$. This implies that there exists a rotation T of \mathbf{R}^2 such that $\text{Supp } \hat{1}_{TE} \supseteq l$. Now $\mu(\sigma(E)) = \nu(\sigma(E))$ for all rigid motions σ , implies that $\check{\mu} * 1_{TE} = \check{\nu} * 1_{TE}$ (where $\check{\mu}(A) = \mu(-A)$), for every rotation T of \mathbf{R}^2 . Thus by the Proposition in §2 it follows that $(\check{\mu})^\wedge = (\check{\nu})^\wedge$ on l . Since l is arbitrary this implies $(\check{\mu})^\wedge = (\check{\nu})^\wedge$, i.e. $\check{\mu} = \check{\nu}$, i.e. $\mu = \nu$, and the proof of our theorem is complete.

REMARK. The technique used in this paper is essentially that of [1]—this paper could be considered a continuation of [1].

ACKNOWLEDGEMENT. The author thanks B. V. Rao, R. L. Karandikar and S. C. Bagchi for useful conversations.

REFERENCES

1. S. C. Bagchi and A. Sitaram, *Determining sets for measures on \mathbf{R}^2* , Illinois J. Math. (to appear).
2. W. F. Donoghue, Jr., *Distributions and Fourier transforms*, Academic Press, New York, 1969.
3. W. Feller, *An introduction to probability theory and its applications*, Vol. II, Wiley, New York, 1966.
4. A. Hertle, *Zur Radon-Transformation von Funktionen und Massen*, Thesis, Erlangen, 1979.
5. W. Rudin, *Functional analysis*, McGraw-Hill, New York, 1973.

STATISTICS-MATHEMATICS DIVISION, INDIAN STATISTICAL INSTITUTE, CALCUTTA 700035, INDIA

Current address: Department of Mathematics, University of Washington, Seattle, Washington 98195