

ON SUMMABILITY OF FOURIER SERIES AT A POINT

G. D. DIKSHIT

ABSTRACT. In this paper summability of Fourier series by a regular linear method of summation determined by a triangular matrix, has been studied and various results—some known and some new—on Cesàro and Nörlund summability have been deduced. A convergence criterion has also been obtained.

1. Let $C = (c_{n,k})$, $k = 0, 1, 2, \dots, n$, be a triangular matrix and let

$$t_n = \sum_{k=0}^n c_{n,k} s_k,$$

where $\{s_k\}$ is a given sequence of numbers. If $t_n \rightarrow s$ as $n \rightarrow \infty$, $\{s_n\}$ is called summable (C) to s . In this paper we assume $c_{n,k} \geq 0$ for $k = 0, 1, 2, \dots, n$, and $\sum_{k=0}^n c_{n,k} = 1$. Then a necessary and sufficient condition for regularity of the method (C) is

$$\lim_{n \rightarrow \infty} c_{n,k} = 0 \quad \text{for each } k.$$

In the case

$$c_{n,k} = A_{n-k}^{\alpha-1} / A_n^\alpha, \quad \alpha \geq 0,$$

where $\{A_n^{\alpha-1}\}$ is determined by the identity

$$(1-x)^{-\alpha} = \sum_0^\infty A_n^{\alpha-1} x^n \quad (|x| < 1),$$

the method (C) reduces to the well-known Cesàro method (C, α) . For

$$c_{n,k} = p_{n-k} / P_n, \quad P_n = p_0 = p_1 + \dots + p_n > 0,$$

the method (C) reduces to the Nörlund method (N, p) . In the case $p_n = 1/(n+1)$, the Nörlund method $(N, 1/(n+1))$ is also known as the harmonic method.

Let f be a Lebesgue integrable periodic function with period 2π and let

$$f(x) \sim \frac{1}{2}a_0 + \sum_1^\infty (a_n \cos nx + b_n \sin nx) \equiv \sum_0^\infty A_n(x).$$

We write

$$\begin{aligned} \phi(t) &= \frac{1}{2} \{f(x+t) + f(x-t) - 2f(x)\}, \\ \Phi(t) &= \int_0^t |\phi(u)| du \quad \text{and} \quad s_n(x) = \sum_0^n A_k(x). \end{aligned}$$

Received by the editors October 13, 1983.

1980 *Mathematics Subject Classification*. Primary 42A24; Secondary 42A20, 40C05.

Key words and phrases. Triangular matrices, Cesàro method, Nörlund method, convergence criteria for Fourier series.

Let

$$C_n(k) = \sum_{m=0}^k c_{n,n-m}$$

and, for $u \geq 0$, define $C_n(u) = C_n([u])$, where $[u]$ is the greatest integer function.

Throughout the paper K is used to denote an absolute constant, not necessarily the same at each occurrence.

2. We establish the following

THEOREM. *Let $\{c_{n,k}\}$ be nondecreasing with respect to k . Let χ be a positive function defined over $(0, \infty)$ such that as $n \rightarrow \infty$, (i) $n\chi(n) = O(1)$ and (ii) $\int_1^n \chi(u)C_n(u) du = O(1)$. Then if $\Phi(t) = o(\chi(\pi/t))$, as $t \rightarrow 0+$, the series $\sum A_n(x)$ is summable (C) to $f(x)$.*

3. Proof. We have that $\{c_{n,k}\}$ is nonnegative and nondecreasing in k . Hence,

$$(n - k)c_{n,k} \leq \sum_{m=k+1}^n c_{n,m} \leq 1.$$

Thus for each fixed k , $c_{n,k} \rightarrow 0$ as $n \rightarrow \infty$, that is, (C) is a regular method.

In view of the fact that the convergence of Fourier series at a point is a local property of the generating function, we may take $\phi(t) = 0$ over $[\delta, \pi]$, where $0 < \delta < \pi$. We choose δ such that $\Phi(t) = o(\chi(\pi/t))$ for $t \in (0, \delta)$. Let

$$t_n(x) = \sum_0^n c_{n,k} s_k(x).$$

Then we need to show that $t_n(x) - f(x) = o(1)$ as $n \rightarrow \infty$. After the Dirichlet integral, for $n > \pi/\delta$,

$$\begin{aligned} t_n(x) - f(x) &= \sum_0^n c_{n,k} s_k(x) - f(x) = \frac{1}{\pi} \int_0^\delta \phi(t)L(n, t) dt \\ &= \frac{1}{\pi} \left\{ \int_0^{\pi/n} + \int_{\pi/n}^\delta \right\} = I_1 + I_2, \quad \text{say,} \end{aligned}$$

where

$$L(n, t) = \sum_0^n \frac{c_{n,k} \sin(k + \frac{1}{2})t}{\sin(\frac{1}{2}t)}.$$

As

$$|L(n, t)| \leq \pi \sum_0^n (k + \frac{1}{2}) c_{n,k} \leq \pi (n + \frac{1}{2}),$$

we get

$$|I_1| \leq \left(n + \frac{1}{2}\right) \int_0^{\pi/n} |\phi(t)| dt = o(n\chi(n)) = o(1),$$

as $n \rightarrow \infty$.

Next, in view of the order estimates of McFadden [4, Lemma 5.11],

$$\left| \sum_{k=a}^b c_{n,n-k} e^{i(n-k)t} \right| \leq KC_n(\pi/t),$$

where $0 \leq a \leq b \leq \infty$, $0 < t \leq \pi$, and n a positive integer, we obtain

$$\begin{aligned} |I_2| &\leq K \int_{\pi/n}^{\delta} \frac{|\phi(t)|C_n(\pi/t)}{t} dt \\ &= K \sum_{k=r}^{n-1} \int_{\pi/(k+1)}^{\pi/k} \frac{|\phi(t)|C_n(\pi/t)}{t} dt + K \int_{\pi/r}^{\delta} \frac{|\phi(t)|C_n(\pi/t)}{t} dt, \end{aligned}$$

where r is a positive integer such that $\pi/r \leq \delta < \pi/(r+1)$. As

$$\int_{\pi/(k+1)}^{\pi/k} \frac{|\phi(t)|C_n(\pi/t)}{t} dt = \left[\frac{C_n(\pi/t)}{t} \Phi(t) \right]_{\pi/(k+1)}^{\pi/k} + \int_{\pi/(k+1)}^{\pi/k} \frac{\Phi(t)C_n(\pi/t)}{t^2} dt,$$

$$\begin{aligned} |I_2| &\leq o(C_n(r)) + o(n\chi(n)C_n(n)) + K \int_{\pi/n}^{\delta} \frac{\Phi(t)C_n(\pi/t)}{t^2} dt \\ &= o(1) + o\left(\int_1^n \chi(u)C_n(u) du\right) = o(1). \end{aligned}$$

This completes the proof of the Theorem.

4. The four corollaries in this section follow as a result of our Theorem.

COROLLARY 1 (HARDY [2]). *Let $\alpha > 0$. If $\Phi(t) = o(t)$, as $t \rightarrow 0+$, then $\sum A_n(x)$ is summable (C, α) to $f(x)$.*

The case $\alpha = 1$ is the classical result of Lebesgue (see [10, Theorem III 3.9]).

PROOF. Let $\chi(u) = \pi/u$ and $c_{n,k} = A_{n-k}^{\alpha-1}/A_n^\alpha$. Then $\chi(\pi/t) = t$ and

$$C_n(u) = \sum_{m=0}^{[u]} c_{n,n-m} = \sum_{m=0}^{[u]} \frac{A_m^{\alpha-1}}{A_n^\alpha} = \frac{A_{[u]}^\alpha}{A_n^\alpha}.$$

Thus $n\chi(n) = \pi$ and

$$\int_1^n \chi(u)C_n(u) du = O(n^{-\alpha}) \int_1^n u^{\alpha-1} du = O(1) \quad \text{as } n \rightarrow \infty.$$

Hence all the hypotheses of the Theorem are satisfied and the result follows.

COROLLARY 2. (i) (SIDDIQI [6]). *If $\Phi(t) = o(t/\log(2\pi/t))$, as $t \rightarrow 0+$, then $\sum A_n(x)$ is summable $(N, 1/(n+1))$ to $f(x)$.*

(ii) *If $\Phi(t) = o(t/\{\log(3\pi/t) \log \log(3\pi/t)\})$, as $t \rightarrow 0+$, then $\sum A_n(x)$ is summable $(N, 1/\{(n+2) \log(n+2)\})$.*

(iii) *If $\Phi(t) = o(t/\{\log(k\pi/t) \log_2(k\pi/t) \cdots \log_q(k\pi/t)\})$, as $t \rightarrow 0+$, then $\sum A_n(x)$ is summable $(N, 1/\{(n+k) \log(n+k) \cdots \log_{q-1}(n+k)\})$, to $f(x)$, where $\log_r x = \log(\log_{r-1} x)$, for $r \geq 2$, and k is such that $\log_q k > 0$.*

PROOF. To deduce this corollary, note that, in case (i) taking

$$\chi(u) = \frac{\pi}{u \log 2u} \quad \text{and} \quad c_{n,k} = \frac{1/(n+1-k)}{\sum_0^n 1/(k+1)},$$

we obtain

$$\begin{aligned} \chi(\pi/t) &= t/\log(2\pi/t), \\ n\chi(n) &= \pi/\log 2n = o(1) \quad \text{as } n \rightarrow \infty, \\ C_n(u) &= \sum_0^{[u]} 1/(m+1) / \sum_0^n 1/(k+1), \end{aligned}$$

and thus

$$\int_1^n \chi(u)C_n(u) du = O\left(\frac{1}{\log n}\right) \int_1^n \frac{1}{u} du = O(1).$$

Thus the hypotheses of the Theorem are satisfied and the result follows.

The choice of χ , $c_{n,k}$, $C_n(u)$, etc., is similarly suggested in each of the cases (ii) and (iii), and the proof of the corollary is completed.

COROLLARY 3. *Let $\{p_n\}$ be a nonnegative, nonincreasing sequence and let $p(1/t) = p([1/t])$ and $P(1/t) = P([1/t])$.*

(i) (SINGH [7]). *If (a) $\Phi(t) = o(t/\log(\pi/t))$ as $t \rightarrow 0+$, and (b) $\sum_1^n (P_k/k \log(k+1)) = O(P_n)$, then $\sum A_n(x)$ is summable (N, p) to $f(x)$.*

(ii) (PATI [5]). *If (c) $\Phi(t) = o(t/P(1/t))$ as $t \rightarrow 0+$, and (d) $\log n = O(P_n)$, then $\sum A_n(x)$ is summable (N, p) to $f(x)$.*

(iii) (SINGH [8]). *If (e) $\Phi(t) = o(p(1/t)/P(1/t))$, as $t \rightarrow 0+$, then $\sum A_n(x)$ is summable (N, p) to $f(x)$.*

REMARKS. In their theorems both Pati and Singh have assumed an extra hypothesis on $\{P_n\}$: " $P_n \rightarrow \infty$, as $n \rightarrow \infty$ ".

PROOF. Since $\{p_n\}$ is nonnegative and nonincreasing,

$$(n+1)p_n \leq p_0 + p_1 + \dots + p_n = P_n.$$

Therefore $np_n/P_n = O(1)$, as $n \rightarrow \infty$. Taking $c_{n,k} = p_{n-k}/P_n$ we obtain

$$C_n(u) = P(u)/P_n.$$

Case (i). Take $\chi(u) = 1$, for $u \in (0, 2)$ and $\chi(u) = \pi/(u \log u)$ for $u \in [2, \infty)$. Then for $t \in (0, 1/2)$,

$$\chi(\pi/t) = t/\log(\pi/t),$$

and, for $n \geq 2$,

$$n\chi(n) = \pi/\log n.$$

Thus

$$n\chi(n) = o(1) \quad \text{as } n \rightarrow \infty.$$

Also

$$\begin{aligned} \int_1^n \chi(u)C_n(u) du &= \frac{P_1}{P_n} + \frac{\pi}{P_n} \int_2^n \frac{P(u)}{u \log u} du \\ &= \frac{P_1}{P_n} + \frac{\pi}{P_n} \sum_2^{n-1} \int_k^{k+1} \frac{P(u)}{u \log u} du \\ &\leq \frac{1}{P_n} \left\{ P_1 + \pi \sum_2^{n-1} \frac{P_k}{k \log k} \right\} \\ &\leq K \left(\frac{1}{P_n} \right) \sum_1^n \frac{P_k}{k \log(k+1)} \\ &= O(1) \quad \text{as } n \rightarrow \infty, \end{aligned}$$

and the hypotheses of the Theorem are satisfied.

Case (ii). Take $\chi(u) = 1/uP(u)$. Then

$$n\chi(n) = 1/P(n) = O(1), \quad \text{as } n \rightarrow \infty,$$

and

$$\int_1^n \chi(u)C_n(u) du = \frac{1}{P_n} \int_1^n \frac{1}{u} du = \frac{\log n}{P_n} = O(1).$$

Case (iii). Let $\chi(u) = p(u)/P(u)$. Then

$$n\chi(n) = np_n/P_n = O(1),$$

as shown earlier, and also

$$\int_1^n \chi(u)C_n(u) du = \frac{1}{P_n} \int_1^n p(u) du = O(1).$$

Thus in each of these cases, the hypotheses of the Theorem are satisfied and the corollary follows.

COROLLARY 4 (A CONVERGENCE CRITERION). *Let χ be a decreasing function such that $\int_1^n \chi(u) du = O(1)$. If $\Phi(t) = o(\chi(\pi/t))$, as $t \rightarrow 0+$, then $\sum A_n(x)$ converges to $f(x)$.*

In particular, if $\chi(\pi/t)$ denotes any of the following:

- (i) $t/(\log(2\pi/t))^{1+\varepsilon}$,
- (ii) $t/\{\log(k\pi/t)(\log \log(k\pi/t))^{1+\varepsilon}\}, \dots$ where $\varepsilon > 0$ and k is appropriately chosen, then $\Phi(t) = O(\chi(\pi/t))$ implies that $\sum A_n(x)$ converges to $f(x)$.

REMARKS. This result may be compared with the corresponding classical results on nonconvergence of a Fourier series at a point of continuity, e.g. see [10, Theorem VIII 2.4, p. 303]. Thus, in the suggested particular cases, $\varepsilon > 0$ may not be replaced by $\varepsilon = 0$. For other alternate convergence criteria involving the case $\varepsilon = 0$, see [3, Theorems 3, 10; 9, Theorems 2, 3].

We shall need the following result for a proof of Corollary 4.

LEMMA [1]. *Let $\{p_n\}$ satisfy the Kaluza conditions:*

$$\text{for } n \geq 0, \quad p_n > 0 \quad \text{and} \quad p_{n+1}/p_n \leq p_{n+2}/p_{n+1} \leq 1.$$

Then if $\{P_n\}$ is bounded, the method (N, p) is ineffective, i.e. only convergent sequences are summable by the method.

PROOF OF COROLLARY 4. We first note that as χ is decreasing,

$$n\chi(n) \leq \int_1^n \chi(u) du = O(1).$$

Now choosing $c_{n,k} = p_{n-k}/P_n$ such that $\{p_n\}$ satisfies the requirements of the Lemma (e.g. $\{p_n\}$ may be taken to be one of the sequences

$$\left\{ \frac{1}{(n+1)(n+2)} \right\}, \quad \left\{ \frac{1}{2^n} \right\}, \quad \left\{ \frac{1}{(n+2)(\log(n+2))^{1+\varepsilon}}, \varepsilon > 0 \right\},$$

etc.), we see that the hypotheses of the Theorem are satisfied, and thus we complete the proof.

In the case of the particular instances cited, we note that

$$\Phi(t) = O(t/(\log(2\pi/t))^{1+\varepsilon}), \quad \text{as } t \rightarrow 0+$$

implies that

$$\Phi(t) = o(t/(\log(2/t))^{1+2/\varepsilon}), \quad \text{as } t \rightarrow 0+,$$

and similarly in the other cases, and then the results as claimed follow.

REFERENCES

1. G. D. Dikshit, *A note on Riesz and Nörlund means*, Rend. Circ. Mat. Palermo (2) **18** (1969), 49–61.
2. G. H. Hardy, *On the summability of Fourier series*, Proc. London Math. Soc. (2) **12** (1913), 365–372; also in *Collected papers of G. H. Hardy*, Vol. 3, Oxford Univ. Press, 1969, pp. 118–125.
3. G. H. Hardy and J. E. Littlewood, *Some new convergence criteria for Fourier series*, Ann. Scuola Norm. Sup. Pisa (2) **3** (1934), 43–62; also in *Collected papers of G. H. Hardy*, op. cit., pp. 68–88.
4. L. McFadden, *Absolute Nörlund summability*, Duke Math. J. **9** (1942), 168–207.
5. T. Pati, *A generalisation of a theorem of Iyengar on the harmonic summability of Fourier series*, Indian J. Math. **3** (1961), 85–90.
6. J. A. Siddiqi, *On the harmonic summability of Fourier series*, Proc. Nat. Acad. Sci. India Sect. A **28** (1948), 527–531.
7. T. Singh, *On Nörlund summability of Fourier series and its conjugate series*, Proc. Nat. Inst. Sci. India Part A **29** (1963), 65–73.
8. —, *Nörlund summability of Fourier series and its conjugate series*, Ann. Mat. Pura Appl. (4) **64** (1964), 123–132.
9. G. Sunouchi, *Convergence criteria for Fourier series*, Tôhoku Math. J. (2) **4** (1952), 187–193.
10. A. Zygmund, *Trigonometric series*, Vol. 1, Cambridge Univ. Press, 1968.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF AUCKLAND, AUCKLAND, NEW ZEALAND