

**SOME ALGEBRAIC SETS OF HIGH LOCAL COHOMOLOGICAL
 DIMENSION IN PROJECTIVE SPACE**

GENNADY LYUBEZNIK

ABSTRACT. Let $V_0, \dots, V_{[n/t]}$ be algebraic sets of pure codimension t in P^n , and suppose $\cap V_i$ is empty. Then $P^n - \cup V_i$ has cohomological dimension $n - [n/t]$.

If U is a scheme, then $cd(U)$, the cohomological dimension of U , is the largest integer i such that there exists a quasi-coherent sheaf F on U such that $H^i(F) \neq 0$.

In [1], G. Faltings proved that if V is an algebraic set of pure codimension t in P^n , then

$$(1) \quad cd(P^n - V) \leq n - [n/t].$$

This note gives some algebraic sets for which equality holds in (1).

THEOREM. Put $s = [n/t]$ and let $V = V_0 \cup V_1 \cup \dots \cup V_s$ be the union of $s + 1$ algebraic sets of pure codimension t in general position in P^n (i.e. such that the intersection of all of them is empty). Then

$$cd(P^n - V) = n - [n/t].$$

This theorem (from the author's thesis [4]) answers the conjecture from [3] in the affirmative and covers all three examples from [3], but not the statement of the main theorem.

For a proof it is convenient to translate the problem into an algebraic language. Put $R_n = k[x_0, \dots, x_n]_{(x_0, \dots, x_n)}$ and let \mathfrak{A} be the defining ideal of V in R_n . Then the cohomological dimension of $P^n - V$ is the largest integer i such that $H_{\mathfrak{A}}^{i+1}(R_n) \neq 0$ (cf. [2]).

LEMMA. Put $s = [n/t]$ and let $\mathfrak{A}_0, \mathfrak{A}_1, \dots, \mathfrak{A}_j$ be $j + 1$ homogeneous ideals of pure height t in R_n . Put $\beta_j = \sum_{r=0}^j \mathfrak{A}_r$. Then $H_{\beta_j}^i(R_n) = 0$ if $i \geq n - s + j + 2$.

PROOF. If $j = 0$, the result follows from (1). Put $\beta_{j-1} = \sum_{r=0}^{j-1} \mathfrak{A}_r$. Then $\beta_{j-1} \cap \mathfrak{A}_j$ has the same radical as $\gamma_{j-1} = \sum_{r=0}^{j-1} \mathfrak{A}_r \cap \mathfrak{A}_j$. Since β_{j-1} and γ_{j-1} are sums of $j - 1$ ideals of pure heights t in R_n , we may assume that $H_{\beta_{j-1}}^i(R_n) = H_{\gamma_{j-1}}^i(R_n) = 0$ for all $i \geq n - s + j + 1$. We also know that $H_{\mathfrak{A}_j}^i(R_n) = 0$ if $i \geq n - s + 2$. The Mayer-Vietoris long exact sequence gives

$$H_{\gamma_{j-1}}^i(R_n) \rightarrow H_{\beta_{j-1}}^{i+1}(R_n) \rightarrow H_{\beta_j}^{i+1}(R_n) \oplus H_{\mathfrak{A}_j}^{i+1}(R_n)$$

and this proves the Lemma.

Received by the editors April 5, 1984.

1980 *Mathematics Subject Classification*. Primary 14B15, 13D03.

PROOF OF THE THEOREM. Let $\mathfrak{A}_0, \dots, \mathfrak{A}_s$ be the defining ideals of V_0, \dots, V_s in R_n . Put $\mathfrak{S}_j = \mathfrak{A}_0 \cap \mathfrak{A}_1 \cap \dots \cap \mathfrak{A}_j + \mathfrak{A}_{j+1} + \dots + \mathfrak{A}_s$. Then the biggest integer i for which $H_{\mathfrak{S}_j}^i(R_n) \neq 0$ is $i = n - j + 1$. We are going to prove this by induction on j and the theorem will follow by putting $j = s$.

For $j = 0$, \mathfrak{S}_j is m -primary, where m is the maximal ideal of R_n and the above claim is well known in this case. Assume $j > 0$ and assume the Theorem proven for $j - 1$. Put $\mathfrak{A}' = \mathfrak{A}_j + \mathfrak{A}_{j+1} + \dots + \mathfrak{A}_s$ and $\mathfrak{A}'' = \mathfrak{A}_0 \cap \mathfrak{A}_1 \cap \dots \cap \mathfrak{A}_{j-1} + \mathfrak{A}_{j+1} + \mathfrak{A}_{j+2} + \dots + \mathfrak{A}_s$. Then $\mathfrak{S}_j = \mathfrak{A}' \cap \mathfrak{A}''$ and $\mathfrak{S}_{j-1} = \mathfrak{A}' + \mathfrak{A}''$. By the Lemma $H_{\mathfrak{A}'}^i(R_n) = H_{\mathfrak{A}''}^i(R_n) = 0$ for all $i \geq n - j + 2$. The claim now follows from the Mayer-Vietoris sequence considering the induction hypothesis. Q.E.D.

REMARK. The above Lemma gives a lower bound on the number of algebraic sets of given codimension which are needed to cut out a given algebraic subset of \mathbf{P}^n set-theoretically. Namely, if $V \subset \mathbf{P}^n$ and $\text{cd}(\mathbf{P}^n - V) = v$, then we need at least $v + 1 - (n - [n/t])$ algebraic subsets of pure codimension t to cut out V set-theoretically.

Faltings' inequality (1) and the fact that it is exact for all n and t (our Theorem) suggest the following.

CONJECTURE. Every algebraic subset of \mathbf{P}^n of pure codimension t is a set-theoretic intersection of $n + 1 - [n/t]$ hypersurfaces [4, p. 8].

For additional supporting evidence for this conjecture see [4, Theorem 6].

REFERENCES

1. G. Faltings, *Über locale Kohomologiegruppen hoher Ordnung*, J. Reine Angew. Math. **313** (1980), 43–51.
2. R. Hartshorne, *Cohomological dimension of algebraic varieties*, Ann. of Math. **88** (1968), 403–450.
3. G. Lyubeznik, *On set-theoretic intersections*, J. Algebra **87** (1984), 105–112.
4. ———, *Set-theoretic intersections and monomial ideals*, Thesis, Columbia University, 1984.

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NEW YORK 10027

Current address: Department of Mathematics, Purdue University, West Lafayette, Indiana 47907