

EXPONENTIAL SUMS OF LERCH'S ZETA FUNCTIONS

KAI WANG

ABSTRACT. For x not an integer and $\operatorname{Re}(s) > 0$, let

$$F(x, s) = \sum_{k=1}^{\infty} \frac{e^{2\pi i k x}}{k^s}$$

be the Lerch's zeta function. In this note, we will show that

$$\sum_{\gamma=1}^{m-1} e^{-2\pi i \gamma \alpha / m} F\left(\frac{\gamma}{m}, 1-n\right) = \frac{1}{n} \left(B_n - m^n B_n \left(\frac{\alpha}{m} - \left[\frac{\alpha}{m} \right] \right) \right)$$

where α is an integer and $\alpha \not\equiv 0 \pmod{m}$ and $n \geq 1$. For $n = 1$, this formula is equivalent to the classical Eisenstein formula

$$\frac{\alpha}{m} - \left[\frac{\alpha}{m} \right] - \frac{1}{2} = -\frac{1}{2m} \sum_{\gamma=1}^{m-1} \sin \frac{2\pi \gamma \alpha}{m} \cot \frac{\pi \gamma}{n}.$$

1. Introduction. About fifteen years ago, S. Chowla [3] proved the following interesting result:

THEOREM A. *Let p be a prime. Then the $\frac{1}{2}(p-1)$ real numbers $\cot \gamma\pi/p$, $\gamma = 1, \dots, \frac{1}{2}(p-1)$, are linear independent over Q , the field of rational numbers.*

Recently, we have extended this result to nonprime numbers and the derivatives of all orders of $\cot x$ [5].

THEOREM B. *For an arbitrary positive integer m , let $\Lambda = \{\gamma | 1 \leq \gamma < \frac{1}{2}m, (\gamma, m) = 1\}$. Then for each $s = 0, 1, \dots$, the $\frac{1}{2}\phi(m)$ real numbers*

$$\left\{ \left(\frac{d^s}{dx^s} \cot \frac{(x+\gamma)\pi}{m} \right)_{x=0} \mid \gamma \in \Lambda \right\}$$

are linear independent over Q where $\phi(m)$ is the Euler function.

On the other hand, there is a classical Eisenstein formula

$$\frac{\alpha}{m} - \left[\frac{\alpha}{m} \right] - \frac{1}{2} = -\frac{1}{2m} \sum_{\gamma=1}^{m-1} \sin \frac{2\pi \gamma \alpha}{m} \cot \frac{\pi \gamma}{m}.$$

Our recent result suggests that it may be of interest to extend the Eisenstein formula to the derivatives of all orders of $\cot x$. Our main result is the following theorem.

Received by the editors June 29, 1984 and, in revised form, November 16, 1984.
 1980 *Mathematics Subject Classification*. Primary 10G05, 10G15; Secondary 10H10.

THEOREM C. *Let m be an arbitrary positive integer. For $s = 1$, let $E_s = \frac{1}{2}$ and for $s \neq 1$, let $E_s = 0$. If s is odd and $s \geq 1$, then*

$$\begin{aligned} \sum_{\alpha=0}^{m-1} \sin \frac{2\pi\alpha\gamma}{m} \left(\frac{d^{s-1}}{dx^{s-1}} \cot \left(x + \frac{\alpha\pi}{m} \right) \right)_{x=0} \\ = \frac{2^s i^{s-1}}{s} \left(m^s B_s \left(\frac{\gamma}{m} - \left[\frac{\gamma}{m} \right] \right) - B_s + E_s \right). \end{aligned}$$

If s is even and $s \geq 1$, then

$$\begin{aligned} \sum_{\alpha=0}^{m-1} \cos \frac{2\pi\alpha\gamma}{m} \left(\frac{d^{s-1}}{dx^{s-1}} \cot \left(x + \frac{\alpha\pi}{m} \right) \right)_{x=0} \\ = \frac{(2i)^s}{s} \left(m^s B_s \left(\frac{\gamma}{m} - \left[\frac{\gamma}{m} \right] \right) - B_s \right), \end{aligned}$$

where $\gamma \not\equiv 0 \pmod{m}$, B_s and $B_s(x)$ are Bernoulli numbers and Bernoulli polynomials, respectively.

We refer to §2 for the definitions of Bernoulli polynomials and Bernoulli numbers.

Theorem C will be proved as a corollary to Theorem D in §2 which involves Lerch's zeta functions.

2. Lerch's zeta functions. Let $\phi(z, s)$ be the power Dirichlet series defined by

$$\phi(z, s) = \sum_{k=1}^{\infty} \frac{z^k}{k^s}$$

which converges for all s if $|z| < 1$, for $\operatorname{Re}(s) > 0$ if $|z| = 1$, $z \neq 1$, and for $\operatorname{Re}(s) > 1$ if $z = 1$. It is known that $\phi(z, s)$ can be extended to the whole s -plane by means of the contour integral. Let $z = e^{2\pi ix}$, where x is real, and the periodic function $F(x, s)$ is defined by the equation $F(x, s) = \phi(e^{2\pi ix}, s)$. $F(x, s)$ is called Lerch's zeta function in the literature. For x not an integer and $\operatorname{Re}(s) > 0$, we have the series representation

$$F(x, s) = \sum_{k=1}^{\infty} \frac{e^{2\pi ikx}}{k^s}.$$

On the other hand, let $B_n(x)$ be the n th Bernoulli polynomials, defined by the power series

$$\frac{te^{xt}}{e^t - 1} = \sum_{k=0}^{\infty} B_k(x) \frac{t^k}{k!}$$

and $B_n = B_n(0)$ is the n th Bernoulli number.

In this note we will prove the following result:

THEOREM D. *Let $m > 2$ be a fixed positive integer and let α be an integer such that $\alpha \not\equiv 0 \pmod{m}$. Then, for $n \geq 1$,*

$$\sum_{\gamma=1}^{m-1} e^{-2\pi i \gamma \alpha / m} F \left(\frac{\gamma}{m}, 1 - n \right) = \frac{1}{n} \left(B_n - m^n B_n \left(\frac{\alpha}{m} - \left[\frac{\alpha}{m} \right] \right) \right).$$

Our proof involves the Fourier series for the Bernoulli polynomials. This was suggested by a referee of a previous version of this note.

3. A proof of Theorem D. Let $B_n(x)$ be the n th Bernoulli polynomial as defined in the Introduction. Then $B_n(x - [x])$ is a periodic function of period 1. It is known that the Fourier series of $B_n(x - [x])$ are given as follows [4, p. 16]:

$$B_{2k-1}(x - [x]) = 2(-1)^k(2k-1)! \sum_{h=1}^{\infty} \frac{\sin 2\pi hx}{(2\pi h)^{2k-1}}$$

and

$$B_{2k}(x - [x]) = 2(-1)^{k-1}(2k)! \sum_{h=1}^{\infty} \frac{\cos 2\pi hx}{(2\pi h)^{2k}}$$

where $B_1(x - [x])$ has to be exempted for integral x because the discontinuity of $B_1(x - [x])$ at x is equal to 0. For our purpose, we may combine the above two formulas into one;

$$B_n(x - [x]) = -(-i)^n n! \sum_{h=1}^{\infty} \frac{e^{2\pi i h x} + (-1)^n e^{-2\pi i h x}}{(2\pi h)^n}$$

for $n > 1$.

Recall that $F(x, s)$ satisfies the following functional equation;

$$F(x, s) = \frac{\Gamma(1-s)}{(2\pi)^{1-s}} (e^{\pi i(1-s)/2} \xi(1-s, x) + e^{\pi i(s-1)/2} \xi(1-s, 1-x)),$$

where $\Gamma(s)$ is the gamma function, $\xi(s, x)$ is the Hurwitz zeta function.

Recall that for $0 < x \leq 1$ and $\operatorname{Re}(s) > 1$,

$$\xi(s, x) = \sum_{h=0}^{\infty} \frac{1}{(h+x)^s}.$$

Since for $n = 1$, Theorem D is equivalent to Eisenstein's formula, we will only consider the case $n > 1$:

$$\begin{aligned} & \sum_{\gamma=1}^{m-1} e^{-2\pi i \gamma \alpha / m} F\left(\frac{\gamma}{m}, 1-n\right) \\ &= \frac{(n-1)! i^n}{(2\pi)^n} \left(\sum_{\gamma=1}^{m-1} e^{-2\pi i \gamma \alpha / m} \xi\left(n, \frac{\gamma}{m}\right) + (-1)^n \sum_{\gamma=1}^{m-1} e^{-2\pi i \gamma \alpha / m} \xi\left(n, \frac{m-\gamma}{m}\right) \right) \\ &= \frac{(n-1)! i^n}{(2\pi)^n} \left(\sum_{\gamma=1}^{m-1} e^{-2\pi i \gamma \alpha / m} \xi\left(n, \frac{\gamma}{m}\right) + (-1)^n \sum_{\beta=1}^{m-1} e^{2\pi i \beta \alpha / m} \xi\left(n, \frac{\beta}{m}\right) \right) \\ &= \frac{(n-1)! i^n}{(2\pi)^n} \left(\sum_{\gamma=1}^{m-1} \sum_{k=0}^{\infty} \frac{e^{2\pi i \gamma \alpha / m} + (-1)^n e^{-2\pi i \gamma \alpha / m}}{(k + \gamma/m)^n} \right) \end{aligned}$$

(and letting $h = mk + \gamma$)

$$\begin{aligned}
&= m^n (n-1)! i^n \sum_{h=1, h \neq O(m)}^{\infty} \frac{e^{2\pi i h \alpha / m} + (-1)^n e^{-2\pi i h \alpha / m}}{(2\pi h)^n} \\
&= m^n (n-1)! i^n \left(\sum_{h=1}^{\infty} \frac{e^{2\pi i h \alpha / m} + (-1)^n e^{-2\pi i h \alpha / m}}{(2\pi h)^n} \right. \\
&\quad \left. - \sum_{k=1}^{\infty} \frac{e^{2\pi i \alpha k} + (-1)^n e^{-2\pi i \alpha k}}{(2\pi m k)^n} \right) \\
&= m^n \left(-\frac{1}{n} B_n \left(\frac{\alpha}{m} - \left[\frac{\alpha}{m} \right] \right) + \frac{1}{nm^n} B_n(\alpha - [\alpha]) \right) \\
&= \frac{1}{n} \left(B_n - m^n B_n \left(\frac{\alpha}{m} - \left[\frac{\alpha}{m} \right] \right) \right).
\end{aligned}$$

This completes the proof.

4. Examples. The first few values of $F(x, -n)$ are given in [2] as follows:

$$\begin{aligned}
F(x, 0) &= -\frac{1}{2}(1 - i \cot \pi x), \quad F(x, -1) = -\frac{1}{4}(1 + \cot^2 \pi x), \\
F(x, -2) &= -\frac{i}{8}(2 \cot \pi x + 2 \cot^3 \pi x).
\end{aligned}$$

It follows that

$$\begin{aligned}
\sum_{\alpha=1}^{m-1} \cos \frac{2\pi \alpha \gamma}{m} \cot^2 \frac{\alpha \pi}{m} &= \frac{2}{3} + 2m^2 B_2 \left(\frac{\gamma}{m} - \left[\frac{\gamma}{m} \right] \right), \\
\sum_{\alpha=1}^{m-1} \sin \frac{2\pi \alpha \gamma}{m} \cot^3 \frac{\alpha \pi}{m} &= -2m B_1 \left(\frac{\gamma}{m} - \left[\frac{\gamma}{m} \right] \right) + \frac{4m^3}{3} B_3 \left(\frac{\gamma}{m} - \left[\frac{\gamma}{m} \right] \right).
\end{aligned}$$

5. A proof of Theorem C. Now Theorem C follows easily from Theorem D because

$$F(x, 0) = \frac{1}{2}(1 - i \cot \pi x).$$

6. Comments. (1) Theorem C is equivalent to the following:

THEOREM E. *Let $m > 2$ be a fixed positive integer and let α be an integer such that $\alpha \not\equiv 0 \pmod{m}$. Then,*

$$\begin{aligned}
\frac{1}{2i} \sum_{\gamma=1}^{m-1} e^{-2\pi i \alpha \gamma / m} \cot \left(x + \frac{\gamma \pi}{m} \right) \\
= \frac{1}{2} - \frac{e^{2xi}}{e^{2xi} - 1} + \frac{me^{2mxi}(\gamma/m - [\gamma/m])}{e^{2mxi} - 1}.
\end{aligned}$$

This formula was originally proved by different arguments without involving Lerch's zeta function. The computation is considerably longer.

(2) Theorem D can also be used to derive the formula for Dirichlet L -function using the following representation of Dirichlet L -function for primitive mod m Dirichlet character χ [2]:

$$\tau(\bar{\chi})L(s, \chi) = \sum_{\alpha=1}^{m-1} \bar{\chi}(\alpha) F \left(\frac{\alpha}{m}, s \right),$$

where $\tau(\bar{\chi})$ is the Gauss sum.

REFERENCES

1. T. M. Apostol, *Introduction to analytic number theory*, Springer-Verlag, Berlin and New York, 1976.
2. ———, *Dirichlet L-functions and character power sums*, J. Number Theory **2** (1970), 223–234.
3. S. Chowla, *The nonexistence of nontrivial linear relations between roots of a certain irreducible equation*, J. Number Theory **2** (1970), 120–123.
4. H. Rademacher, *Topics in analytic number theory*, Springer-Verlag, Berlin and New York, 1973.
5. K. Wang, *On a theorem of S. Chowla*, J. Number Theory **15** (1982), 1–4.

DEPARTMENT OF MATHEMATICS, WAYNE STATE UNIVERSITY, DETROIT, MICHIGAN
48202