

SHORTER NOTES

The purpose of this department is to publish very short papers of unusually polished character, for which there is no other outlet.

A FORMALLY NORMAL OPERATOR HAVING NO NORMAL EXTENSION

KONRAD SCHMÜDGEN

ABSTRACT. We give an example of a formally normal operator N satisfying $\dim \mathcal{D}(N^*)/\mathcal{D}(N) = 1$ which has no normal extension in any larger Hilbert space.

For a linear operator T on a Hilbert space \mathcal{H} , we denote by $\mathcal{D}(T)$ its domain. A *formally normal operator* N on \mathcal{H} is a densely defined closed linear operator on \mathcal{H} such that $\mathcal{D}(N) \subseteq \mathcal{D}(N^*)$ and $\|Nx\| = \|N^*x\|$ for all $x \in \mathcal{D}(N)$. A *normal operator* is a formally normal operator N on \mathcal{H} satisfying $\mathcal{D}(N) = \mathcal{D}(N^*)$.

The first example of a formally normal operator N which has no normal extension in a larger Hilbert space is due to E. A. Coddington [1]. Coddington's example is minimal in the sense that $\dim \mathcal{D}(N^*)/\mathcal{D}(N) = 1$. The aim of this note is to present a very simple example of that kind. For let S be the unilateral shift on the Hardy space $\mathcal{H} = H^2(\mathbb{T})$, and let $A := i(S + I)(S - I)^{-1}$. Set $B := S + S^*$.

PROPOSITION. *The operator $N := A + iB$ is a formally normal operator on \mathcal{H} which has no normal extension in a possibly larger Hilbert space. Moreover, $\dim \mathcal{D}(N^*)/\mathcal{D}(N) = 1$.*

PROOF. First we show that $BA \subseteq A^*B$. Let $x \in \mathcal{D}(A)$. Then $x = (S - I)y$ for some $y \in \mathcal{H}$ and $B(A - i)x = 2iBy$. Let $P := z^0 \otimes z^0$. Since $(I - P)BS = SB$ and $z^0 \in \ker(A^* - i)$, $Bx = (S - I)By + PBSy \in \mathcal{D}(A^*)$ and $(A^* - i)Bx = (A - i)(S - I)By = 2iBy = B(A - i)x$. This proves $BA \subseteq A^*B$.

From $BA \subseteq A^*B$ and $B = B^*$, it follows that $\langle Ax, Bx \rangle = \langle Bx, Ax \rangle$ for $x \in \mathcal{D}(A)$. Since $N^* = A^* - iB$ and $\mathcal{D}(N) \subseteq \mathcal{D}(N^*)$, the latter implies $\|Nx\|^2 = \|Ax\|^2 + \|Bx\|^2 = \|N^*x\|^2$ for $x \in \mathcal{D}(N) = \mathcal{D}(A)$. That is, N is formally normal. Since A has deficiency indices $(0, 1)$, $\dim \mathcal{D}(N^*)/\mathcal{D}(N) = 1$.

Received by the editors January 18, 1985.

1980 *Mathematics Subject Classification.* Primary 47A20, 47B15.

Key words and phrases. Formally normal operator, unbounded normal operator.

©1985 American Mathematical Society
0002-9939/85 \$1.00 + \$.25 per page

Assume that there exists a normal extension N_1 of N in a Hilbert space $\mathcal{H}_1 \supseteq \mathcal{H}$. Let U_1 denote the Cayley transform of the selfadjoint operator $A_1 := \operatorname{Re} N_1$ on \mathcal{H}_1 . From $A_1 \supseteq A$ and $(A - i)\mathcal{D}(A) = \mathcal{H}$ it follows that $U_1 \upharpoonright \mathcal{H} = S$. Clearly, $B \subseteq B_1 := \operatorname{Im} N_1$. Because N is normal, $U_1 B_1 \subseteq B_1 U_1$. Since $B = B_1 \upharpoonright \mathcal{H}$ and $S = U_1 \upharpoonright \mathcal{H}$, this implies $SB = BS$. Since $B = S + S^*$ by definition, this is a contradiction.

REMARKS. 1. A slight modification of the above proof shows the following: Suppose A is a maximal symmetric operator on a Hilbert space \mathcal{H} . Suppose B is a bounded selfadjoint operator on H such that $BA \subseteq A^*B$, but $BA \not\subseteq AB$. Then $N := A + iB$ is a formally normal operator which cannot be extended to a normal operator in any larger Hilbert space.

2. In the above example, B can be replaced by an arbitrary bounded selfadjoint Toeplitz operator B_φ with symbol $\varphi \neq 0$ in $L^\infty(\mathbb{T})$. (By Example 1.10 in [2], $B_\varphi A \subseteq A^*B_\varphi$ and $B_\varphi A \not\subseteq AB_\varphi$.)

3. Suppose A and B are selfadjoint operators which commute on a common core \mathcal{D} and for which the spectral projections do not commute. Then $N := (A + iB) \upharpoonright \mathcal{D}$ is formally normal and has no normal extension in a possibly larger Hilbert space [3, Lemma 1.5]. But it seems to be more difficult in general to calculate $\dim \mathcal{D}(N^*)/\mathcal{D}(N)$ in that case.

REFERENCES

1. E. A. Coddington, *Formally normal operators having no normal extensions*, *Canad. J. Math.* **17** (1965), 1030–1040.
2. K. Schmüdgen, *Unbounded commutants and intertwining spaces of unbounded symmetric operators and *-representations* (to appear).
3. _____, *On commuting unbounded self-adjoint operators*. III, Preprint, 1984; *Manuscripta Math.* (to appear).

SEKTION MATHEMATIK, KARL-MARX-UNIVERSITÄT LEIPZIG, 7010 LEIPZIG, GERMAN DEMOCRATIC REPUBLIC