

WHITNEY LEVELS IN $C_p(X)$ ARE ARS

MARK LYNCH¹

ABSTRACT. For X a metric continuum, and $p \in X$, we show that the Whitney levels in the relative hyperspace $C_p(X) = \{K \in C(X) | p \in K\}$ are absolute retracts.

1. Introduction. For (X, d) a metric continuum, let $C(X)$ denote the hyperspace of subcontinua with the Hausdorff metric H . A *Whitney map* $\mu: C(X) \rightarrow [0, 1]$ is a map such that $\mu(\{x\}) = 0$ for each $x \in X$, $\mu(X) = 1$, and $\mu(K) < \mu(M)$ whenever $K \subsetneq M$. Such maps may always be constructed [8]. The point-inverses $\mu^{-1}(t)$, $0 \leq t \leq 1$, are subcontinua in $C(X)$ [4], and are called *Whitney levels*. For $p \in X$, let $C_p(X) = \{K \in C(X) | p \in K\}$ be the relative hyperspace, and let μ_p denote the restriction of μ to $C_p(X)$. Eberhart [3] showed that $C_p(X)$ is always an AR. In this paper we consider the relative Whitney levels $\mu_p^{-1}(t)$ in $C_p(X)$. We will use a construction in a space of order arcs to show that each Whitney level $\mu_p^{-1}(t)$ is an AR.

Krasinkiewicz and Nadler [7] and Rogers [11] have shown that $\mu_p^{-1}(t)$ is arcwise connected, and Rogers [10] showed that $\mu_p^{-1}(t)$ is acyclic. We point out that the Whitney levels $\mu^{-1}(t)$ need not be arcwise connected [6], and even if X is a 2-cell, they need not be ARs [9].

2. The space of orders arcs $\Lambda_p(X)$. An arc $\alpha \subset C(X)$ is an *order arc* if, for all $K, M \in \alpha$, either $K \subset M$ or $M \subset K$. Then $\cap \alpha = \cap \{M | M \in \alpha\}$ and $\cup \alpha = \cup \{M | M \in \alpha\}$ are the endpoints of α . For every pair $A, B \in C(X)$ with $A \subset B$, there exists an order arc α with $\cap \alpha = A$ and $\cup \alpha = B$ [6]. An order arc α may be parametrized by defining $\alpha(t)$ to be the unique $K \in \alpha$ such that $\mu(K) = (1-t) \cdot \mu(\cap \alpha) + t \cdot \mu(\cup \alpha)$. Let Λ_p denote the space of *maximal order arcs* in $C_p(X)$, i.e., $\Lambda_p = \{\text{order arcs } \alpha \subset C_p(X) | \alpha(0) = \{p\} \text{ and } \alpha(1) = X\} \subset C(C(X))$. For each $t \in [0, 1]$, the *evaluation map* $e_t: \Lambda_p \rightarrow \mu_p^{-1}(t)$, defined by $e_t(\alpha) = \alpha(t)$, is onto. We will show that $\mu_p^{-1}(t)$ is an absolute extensor for metric spaces by imitating Dugundji's proof of the extension property for maps into locally convex linear metric spaces [2]. To do this, we utilize the evaluation map e_t and a type of convex structure on Λ_p .

For $\alpha_1, \dots, \alpha_n \in \Lambda_p$ and $t_1, \dots, t_{n-1} \in [0, 1]$, let $\beta = \langle \alpha_1, t_1; \dots; \alpha_{n-1}, t_{n-1}; \alpha_n \rangle$ be the element of Λ_p defined by

$$\beta = \{\alpha_1(s) | 0 \leq s \leq t_1\} \cup \{\alpha_1(t_1) \cup \alpha_2(s) | 0 \leq s \leq t_2\} \\ \cup \dots \cup \{\alpha_1(t_1) \cup \dots \cup \alpha_{n-1}(t_{n-1}) \cup \alpha_n(s) | 0 \leq s \leq 1\}.$$

Received by the editors July 24, 1985.

1980 *Mathematics Subject Classification*. Primary 54B20, 54C55; Secondary 54F20.

Key words and phrases. Absolute retract, hyperspace of subcontinua, Whitney level.

¹This work represents part of the author's dissertation, written at Louisiana State University under the supervision of Professor Doug Curtis.

This construction is continuous, in the sense that if $\alpha_i^m \rightarrow \alpha_i$ and $t_i^m \rightarrow t_i$ as $m \rightarrow \infty$, for each i , then $(\alpha_1^m, t_1^m; \dots; \alpha_{n-1}^m, t_{n-1}^m; \alpha_n^m) \rightarrow (\alpha_1, t_1; \dots; \alpha_{n-1}, t_{n-1}; \alpha_n)$.

3. Extending maps into $\mu_p^{-1}(t)$. Let (Z, ρ) be a metric space, and $A \subset Z$ a closed subset. Given a map $g: A \rightarrow \mu_p^{-1}(t)$, we define an extension $\hat{g}: Z \rightarrow \mu_p^{-1}(t)$ of g by a Dugundji-type construction. For every $x \in Z - A$, let $B_x = \{z \in Z \mid \rho(x, z) < \frac{1}{2} \cdot \rho(x, A)\}$. Let $\mathbf{U} = \{U_\alpha \mid \alpha \in \mathbf{A}\}$ be a neighborhood finite open refinement of $\{B_x \mid x \in Z - A\}$, indexed by a well-ordered set \mathbf{A} . Let $\{\phi_\alpha \mid \alpha \in \mathbf{A}\}$ be a partition of unity of $Z - A$ subordinated to \mathbf{U} . With each $\alpha \in \mathbf{A}$, associate $a_\alpha \in A$ as follows: Choose $x_\alpha \in U_\alpha$, and take $a_\alpha \in A$ with $\rho(x_\alpha, a_\alpha) < 2 \cdot \rho(x_\alpha, A)$. For each $\alpha \in \mathbf{A}$, choose $\beta_\alpha \in \Lambda_p$ such that $e_t(\beta_\alpha) = \beta_\alpha(t) = g(a_\alpha)$. Then with each $x \in Z - A$ there is associated a finite set $\{\beta_\alpha \mid \phi_\alpha(x) > 0\}$ of elements of Λ_p . The extension $\hat{g}: Z \rightarrow \mu_p^{-1}(t)$ is defined in the following steps:

(1) For $x \in Z - A$, let $\alpha_1 < \alpha_2 < \dots < \alpha_n$ be the ordering in \mathbf{A} of those elements α for which $\phi_\alpha(x) > 0$, and define

$$\tau(x, \alpha_i) = \frac{\phi_{\alpha_i}(x)}{\phi_{\alpha_1}(x) + \dots + \phi_{\alpha_n}(x)}, \quad i = 1, 2, \dots, n.$$

Note that $\tau(x, \alpha_n) = 1$.

(2) With $\beta_{\alpha_1}, \dots, \beta_{\alpha_n}$ the elements of Λ_p corresponding to x , define $\beta_x = \langle \beta_{\alpha_1}, \tau(x, \alpha_1); \dots; \beta_{\alpha_{n-1}}, \tau(x, \alpha_{n-1}); \beta_{\alpha_n} \rangle$.

(3) Define $\hat{g}(x) = e_t(\beta_x) = \beta_x(t)$.

CLAIM 1. \hat{g} is continuous on $Z - A$.

Consider $x \in Z - A$, with $\alpha_1 < \dots < \alpha_n$ as in step (1) above. For any $y \in Z$ sufficiently close to x , each $\phi_{\alpha_i}(y)$ will be near $\phi_{\alpha_i}(x)$, $i = 1, \dots, n$. Clearly, this implies that if $\gamma_1 < \dots < \gamma_{n+m}$ is the ordering in \mathbf{A} of $\{\alpha \mid \phi_\alpha(y) > 0\}$, then for each k such that $\gamma_k = \alpha_i$ for some i , $\tau(y, \gamma_k)$ is near $\tau(x, \alpha_i)$, and for all other γ_k , either $\tau(y, \gamma_k)$ is near 0 or $\alpha_n < \gamma_k$. It follows that β_y is near β_x , and $\hat{g}(y)$ is near $\hat{g}(x)$.

CLAIM 2. \hat{g} is continuous on $\text{bd } A$.

Consider $a \in \text{bd } A$, and let $y \in Z - A$ denote a point near a . Let $\{\alpha_1, \dots, \alpha_n\} = \{\alpha \in \mathbf{A} \mid \phi_\alpha(y) > 0\}$. Then for each $i \leq n$, the point $a_{\alpha_i} \in A$ associated with α_i is near a , thus $\beta_{\alpha_i}(t) = g(a_{\alpha_i})$ is near $g(a)$. Note that the construction of the order arc β_y forces $\beta_y(t) \subset \beta_{\alpha_1}(t) \cup \beta_{\alpha_2}(t) \cup \dots \cup \beta_{\alpha_n}(t)$. Since $M = \bigcup_1^n \beta_{\alpha_i}(t)$ is an element of $C_p(X)$ near $g(a) \in \mu_p^{-1}(t)$, $\mu(M)$ is near $t = \mu(\beta_y(t))$. Since $\beta_y(t) \subset M$, the nature of the Whitney map μ forces $\beta_y(t)$ and M to be close. Then $\hat{g}(y) = \beta_y(t)$ is near $\hat{g}(a) = g(a)$.

Thus, the extension $\hat{g}: Z \rightarrow \mu_p^{-1}(t)$ of g is continuous, and this concludes the proof of our main result:

THEOREM. *Each Whitney level $\mu_p^{-1}(t)$ in $C_p(X)$ is an AR.*

There are several easy corollaries.

COROLLARY 1. *$\{X\}$ is an unstable point in $C_p(X)$.*

PROOF. Given $\varepsilon > 0$, choose $t < 1$ sufficiently close to 1 so that

$$\text{diam}_H(\mu_p^{-1}([t, 1])) < \varepsilon.$$

Since $\mu_p^{-1}(t)$ is an AR, there is a retraction $r: \mu_p^{-1}([t, 1]) \rightarrow \mu_p^{-1}(t)$. Then r extends by the identity to a retraction $R: C_p(X) \rightarrow \mu_p^{-1}([0, t])$. We have $R(C_p(X)) \subset C_p(X) - \{X\}$ and $H(R(M), M) < \varepsilon$ for each $M \in C_p(X)$. Thus, $\{X\}$ is unstable in $C_p(X)$.

Note that the above retraction R shows that $\mu_p^{-1}([0, t])$ is an AR. In fact, a similar argument shows that $\mu_p^{-1}([s, t])$ is an AR for all $s < t$.

COROLLARY 2. $C_p(X) - \{X\}$ is an AR.

PROOF. Since $C_p(X) - \{X\}$ is an ANR, it suffices to show that it is n -connected for all n [5]. Let $f: S^n \rightarrow C_p(X) - \{X\}$ be a map of the n -sphere. Choose $t < 1$ such that $f(S^n) \subset \mu_p^{-1}([0, t])$. Then f is null-homotopic in the AR $\mu_p^{-1}([0, t])$, thus $C_p(X) - \{X\}$ is n -connected for all n .

If X has a cut point p , then clearly $\mu^{-1}(t) = \mu_p^{-1}(t)$ for all t in some neighborhood of 1. Thus we have the following

COROLLARY 3. If X has a cut point, then

- (i) for all t in some neighborhood of 1, $\mu^{-1}(t)$ is an AR; and
- (ii) $\{X\}$ is unstable in $C(X)$.

The general question suggested by part (ii) has been answered in [1]: For X a Peano continuum, $\{X\}$ is stable in $C(X)$ if and only if X is a finite graph with no cut points.

BIBLIOGRAPHY

1. D. W. Curtis, *Stable points in hyperspaces of Peano continua*, preprint.
2. J. Dugundji, *An extension of Tietze's theorem*, Pacific J. Math. **1** (1951), 353-367.
3. Carl Eberhart, *Intervals of continua which are Hilbert cubes*, Proc. Amer. Math. Soc. **68** (1978), 220-224.
4. Carl Eberhart and Sam B. Nadler, Jr., *The dimension of certain hyperspaces*, Bull. Polon. Acad. Sci. Math. **19** (1971), 1027-1034.
5. S. T. Hu, *Theory of retracts*, Wayne State Univ. Press, 1965.
6. J. L. Kelley, *Hyperspaces of a continuum*, Trans. Amer. Math. Soc. **52** (1942), 22-36.
7. J. Krasinkiewicz and Sam B. Nadler, Jr., *Whitney properties*, Fund. Math. **98** (1978), 165-180.
8. Sam B. Nadler, Jr., *Hyperspaces of sets*, Dekker, New York and Basel, 1978.
9. Ann Petrus, *Whitney maps and Whitney properties in $C(X)$* , Topology Proc., vol. 1 (Proc. 1976 Topology Conference, Auburn Univ.), Math. Dept., Auburn Univ., Auburn, Ala., 1977, pp. 147-172.
10. James T. Rogers, Jr., *Applications of a Vietoris-Begle theorem for multi-valued maps to the cohomology of hyperspaces*, Michigan J. Math. **22** (1976), 315-319.
11. —, *Whitney continua in the hyperspace $C(X)$* , Pacific J. Math. **58** (1975), 569-584.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA 70803

Current address: 5466 Harpers Farm Road, Apartment A4, Columbia, Maryland 21044