

A SHORT PROOF OF CAUCHY'S POLYGONAL NUMBER THEOREM

MELVYN B. NATHANSON

ABSTRACT. This paper presents a simple proof that every nonnegative integer is the sum of $m + 2$ polygonal numbers of order $m + 2$.

Let $m \geq 1$. The polygonal numbers of order $m + 2$ are the integers

$$p_m(k) = \frac{m}{2}(k^2 - k) + k$$

for $k = 0, 1, 2, \dots$. Fermat [3] asserted that every nonnegative integer is the sum of $m + 2$ polygonal numbers of order $m + 2$. For $m = 2$, Lagrange [5] proved that every nonnegative integer is the sum of four squares $p_2(k) = k^2$. For $m = 1$, Gauss [4] proved that every nonnegative integer is the sum of three triangular numbers $p_1(k) = (k^2 + k)/2$, or, equivalently, that every positive integer $n \equiv 3 \pmod{8}$ is the sum of three odd squares. Cauchy [1] proved Fermat's statement for all $m \geq 3$, and Legendre [6] refined and extended this result. For $m \geq 3$ and $n \leq 120m$, Pepin [8] published tables of explicit representations of n as a sum of $m + 2$ polygonal numbers of order $m + 2$, at most four of which are different from 0 or 1. Dickson [2] prepared similar tables. Pall [7] obtained important related results on sums of values of a quadratic polynomial.

Uspensky and Heaslet [9, p. 380] and Weil [10, p. 102] have written that there is no short and easy proof of Cauchy's polygonal number theorem. The object of this note is to present a short and easy proof.

Because of Pepin's and Dickson's tables, it suffices to consider only $n \geq 120m$. For completeness, I also include a proof of Cauchy's lemma.

CAUCHY'S LEMMA. *Let a and b be odd positive integers such that $b^2 < 4a$ and $3a < b^2 + 2b + 4$. Then there exist nonnegative integers s, t, u, v such that*

$$(1) \quad a = s^2 + t^2 + u^2 + v^2,$$

$$(2) \quad b = s + t + u + v.$$

PROOF. Since a and b are odd, it follows that $4a - b^2 \equiv 3 \pmod{8}$, and so, by Gauss's triangular number theorem, there exist odd integers $x \geq y \geq z > 0$ such that

$$(3) \quad 4a - b^2 = x^2 + y^2 + z^2.$$

Received by the editors August 26, 1985 and, in revised form, December 20, 1985.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 11P05, 11B83, 11B13.

Key words and phrases. Polygonal numbers, sums of squares, Cauchy-Fermat theorem.

©1987 American Mathematical Society
0002-9939/87 \$1.00 + \$.25 per page

Choose the sign of $\pm z$ so that $b + x + y \pm z \equiv 0 \pmod{4}$. Define integers s, t, u, v by

$$s = \frac{b + x + y \pm z}{4}, \quad t = \frac{b + x}{2} - s = \frac{b + x - y \mp z}{4},$$

$$u = \frac{b + y}{2} - s = \frac{b - x + y \mp z}{4}, \quad v = \frac{b \pm z}{2} - s = \frac{b - x - y \pm z}{4}.$$

Then equations (1) and (2) are satisfied, and $s \geq t \geq u \geq v$. To show these integers are nonnegative, it suffices to prove that $v \geq 0$, or $v > -1$. This is true if $b - x - y - z > -4$, or, equivalently, if $x + y + z < b + 4$. The maximum value of $x + y + z$ subject to the constraint (3) is $\sqrt{12a - 3b^2}$, and the inequality $3a < b^2 + 2b + 4$ implies that $x + y + z \leq \sqrt{12a - 3b^2} < b + 4$. This proves the lemma.

THEOREM 1. *Let $m \geq 3$ and $n \geq 120m$. Then n is the sum of $m + 1$ polygonal numbers of order $m + 2$, at most four of which are different from 0 or 1.*

PROOF. Let b_1 and b_2 be consecutive odd integers. The set of numbers of the form $b + r$, where $b \in \{b_1, b_2\}$ and $r \in \{0, 1, \dots, m - 3\}$, contains a complete set of residue classes modulo m , and so $n \equiv b + r \pmod{m}$ for some $b \in \{b_1, b_2\}$ and $r \in \{0, 1, \dots, m - 3\}$. Define

$$(4) \quad a = 2\left(\frac{n - b - r}{m}\right) + b = \left(1 - \frac{2}{m}\right)b + 2\left(\frac{n - r}{m}\right).$$

Then a is an odd integer, and

$$(5) \quad n = \frac{m}{2}(a - b) + b + r.$$

If $0 < b < \frac{2}{3} + \sqrt{8(n/m) - 8}$, then the quadratic formula implies that

$$b^2 - 4a = b^2 - 4\left(1 - \frac{2}{m}\right)b - 8\left(\frac{n - r}{m}\right) < 0$$

and so $b^2 < 4a$. Similarly, if $b > \frac{1}{2} + \sqrt{6(n/m) - 3}$, then $3a < b^2 + 2b + 4$. Since the length of the interval

$$(6) \quad I = \left(\frac{1}{2} + \sqrt{6\left(\frac{n}{m}\right) - 3}, \frac{2}{3} + \sqrt{8\left(\frac{n}{m}\right) - 8}\right)$$

is greater than 4, it follows that I contains two consecutive odd positive integers b_1 and b_2 . Thus, there exist odd positive integers a and b that satisfy (5) and the inequalities $b^2 < 4a$ and $3a < b^2 + 2b + 4$. Cauchy's Lemma implies that there exist s, t, u, v satisfying (1) and (2), and so

$$n = \frac{m}{2}(a - b) + b + r = \frac{m}{2}(s^2 - s) + s + \dots + \frac{m}{2}(v^2 - v) + v + r$$

$$= p_m(s) + p_m(t) + p_m(u) + p_m(v) + r.$$

This completes the proof.

Note that this result is slightly stronger than Cauchy's theorem. Legendre [6] proved that every sufficiently large integer is the sum of five polygonal numbers of order $m + 2$, one of which is either 0 or 1. This can also be easily proved.

THEOREM 2. *Let $m \geq 3$. If m is odd, then every sufficiently large integer is the sum of four polygonal numbers of order $m + 2$. If m is even, then every sufficiently large integer is the sum of five polygonal numbers of order $m + 2$, one of which is either 0 or 1.*

PROOF. There is an absolute constant c such that if $n > cm^3$, then the length of the interval I defined in (6) is greater than $2m$, and so I contains at least m consecutive odd integers.

If m is odd, these form a complete set of residues modulo m , and so $n \equiv b \pmod{m}$ for some odd number $b \in I$. Let $r = 0$. Define a by formula (4).

If m is even and $n > cm^3$, then $n \equiv b + r \pmod{m}$ for some odd integer $b \in I$ and $r \in \{0, 1\}$. Define a by (4).

In both cases, the theorem follows immediately from Cauchy's Lemma.

REFERENCES

1. A. Cauchy, *Démonstration du théorème général de Fermat sur les nombres polygones*, Mém. Sci. Math. Phys. Inst. France (1) **14** (1813–15), 177–220 = *Oeuvres* (2), vol. 6, 320–353.
2. L. E. Dickson, *All positive integers are sums of values of a quadratic function of x* , Bull. Amer. Math. Soc. **33** (1927), 713–720.
3. P. Fermat, quoted in T. L. Heath, *Diophantus of Alexandria*, Dover, New York, 1964, p. 188.
4. C. F. Gauss, *Disquisitiones arithmeticae*, Yale Univ. Press, New Haven, Conn., and London, 1966.
5. J. L. Lagrange, *Démonstration d'un théorème d'arithmétique*, Nouveaux Mémoires de l'Acad. Royale des Sci. et Belles-L. de Berlin, 1770, pp. 123–133 = *Oeuvres*, vol. 3, pp. 189–201.
6. A.-M. Legendre, *Théorie des nombres*, 3rd ed., vol. 2, 1830, pp. 331–356.
7. G. Pall, *Large positive integers are sums of four or five values of a quadratic function*, Amer. J. Math. **54** (1932), 66–78.
8. T. Pepin, *Démonstration du théorème de Fermat sur les nombres polygones*, Atti Accad. Pont. Nuovi Lincei **46** (1892–93), 119–131.
9. J. V. Uspensky and M. A. Heaslet, *Elementary number theory*, McGraw-Hill, New York and London, 1939.
10. A. Weil, *Number theory, an approach through history from Hammurabi to Legendre*, Birkhäuser, Boston, Mass., 1983.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEWARK, NEW JERSEY 07102

Current address: Office of the Provost and Vice President for Academic Affairs, Lehman College (CUNY), Bronx, New York 10468