

CESÀRO AND BOREL-TYPE SUMMABILITY

DAVID BORWEIN AND TOM MARKOVICH

(Communicated by R. Daniel Mauldin)

ABSTRACT. Though summability of a series by the Cesàro method C_p does not in general imply its summability by the Borel-type method (B, α, β) , it is shown that the implication holds under an additional condition.

1. Introduction. Suppose throughout that $\sum_{n=0}^{\infty} a_n$ is a series with partial sums $s_n := \sum_{k=0}^n a_k$, and that $\alpha > 0$ and $\alpha N + \beta > 0$ where N is a nonnegative integer. The series $\sum_{n=0}^{\infty} a_n$ is said to be summable (B, α, β) to s if

$$\alpha e^{-x} \sum_{n=N}^{\infty} s_n \frac{x^{\alpha n + \beta - 1}}{\Gamma(\alpha n + \beta)} \rightarrow s \quad \text{as } x \rightarrow \infty.$$

The Borel-type summability method (B, α, β) is regular, and $(B, 1, 1)$ with $N = 0$ is the standard Borel summability method B .

We shall also be concerned with the Cesàro summability method C_p ($p > -1$) and the Valiron method V_α defined as follows:

$$\sum_{n=0}^{\infty} a_n = s(C_p) \quad \text{if } c_n^p := \frac{s_n^p}{\binom{n+p}{n}} \rightarrow s \quad \text{as } n \rightarrow \infty$$

where

$$s_n^p := \sum_{k=0}^n \binom{n-k+p-1}{n-k} s_k;$$

$$\sum_{n=0}^{\infty} a_n = s(V_\alpha) \quad \text{if } \left(\frac{\alpha}{2\pi n}\right)^{1/2} \sum_{k=0}^{\infty} \exp\left(-\frac{\alpha(n-k)^2}{2n}\right) s_k \rightarrow s \quad \text{as } n \rightarrow \infty.$$

Consider the series $\sum_{n=1}^{\infty} a_n := \sum_{n=1}^{\infty} n^{a-1} \exp(Ain^a)$ where $A > 0$ and $0 < a < 1/2$. It is known [5, p. 213] that this series is summable C_p for every $p > 0$ but is not convergent. However, since $a_n = o(n^{-1/2})$, it follows by the Borwein Tauberian Theorem [1, Theorem 1] that the series is not summable (B, α, β) for any α and β . This example shows that, in general, summability C_p does not imply summability (B, α, β) . The following theorem indicates how to strengthen the C_p summability hypothesis in order to ensure summability (B, α, β) .

THEOREM 1. *Suppose that p is a nonnegative integer and that $c_n^p = s + o(n^{-p/2})$ as $n \rightarrow \infty$. Then $\sum_{n=0}^{\infty} a_n = s(B, \alpha, \beta)$.*

The special case $\alpha = \beta = 1$, $p = 1$ of Theorem 1 has been proved by Hardy [5, Theorem 149]. Hardy and Littlewood [4, §3] proved that the condition

Received by the editors July 14, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 40G05, 40G10.

This research was supported in part by the Natural Sciences and Engineering Council of Canada.

$c_n^p = s + o(n^{-1/2})$ is not sufficient for the summability of $\sum a_n$ by the Borel method. Hyslop [7, Theorem VIII] has obtained a more general result than Hardy, namely the case $\alpha = \beta = 1$ of Theorem 1. More recently, Swaminathan [10] has proved Theorem 1 with $p = 1$ and (B, α, β) summability replaced by the more general $F(a, q)$ summability introduced by Meir [9].

2. Preliminary results.

LEMMA 1 [8, LEMMA 7]. *Let $m < x_0 < n - 1$ where m, n are integers and let the nonnegative function $f(x)$ be increasing on $[m, x_0]$ and decreasing on $[x_0, n]$. Then*

$$\sum_{k=m}^n f(k) \leq \int_m^n f(x)dx + f(x_0).$$

LEMMA 2 [2, THEOREM 3]. *Suppose that $s_n = O(n^r)$ where $r \geq 0$. Then $\sum_{n=0}^\infty a_n = s(B, \alpha, \beta)$ if and only if $\sum_{n=0}^\infty a_n = s(V_\alpha)$.*

THEOREM 2 (CF. [6, THEOREM 2]). *Suppose that p is a nonnegative integer and that $c_n^p = s + o(n^{-p/2})$ as $n \rightarrow \infty$. Then $\sum_{n=0}^\infty a_n = s(V_\alpha)$.*

PROOF. Suppose, as we may without loss of generality, that $s = 0$.

Let $v_n(x) := \exp(-\alpha(n - x)^2/2n)$ and denote the p th difference of $v_n(k)$ by $\Delta^p v_n(k)$, so that

$$\Delta^p v_n(k) = \sum_{r=0}^p \binom{p}{r} (-1)^r v_n(k + r).$$

Applying Abel’s partial summation formula $p (< m)$ times, we have that

$$\sum_{k=0}^m s_k v_n(k) = \sum_{k=0}^{m-p} s_k^p \Delta^p v_n(k) + \sum_{r=0}^{p-1} s_{m-r}^{r+1} \Delta^r v_n(m - r).$$

Letting $m \rightarrow \infty$ and applying the limitation theorem for Cesàro summability [5, Theorem 46], we see that

$$F(n) := \sum_{k=0}^\infty s_k v_n(k) = \sum_{k=0}^\infty s_k^p \Delta^p v_n(k).$$

In order to prove the theorem we must show that $F(n) = o(n^{1/2})$. Since, by the hypothesis, $s_k^p = o(k^{p/2})$ as $k \rightarrow \infty$ and $k^{p/2} \Delta^p v_n(k) = o(n^{1/2})$ as $n \rightarrow \infty$, it suffices to show that

$$(1) \quad G(n) := n^{-1/2} \sum_{k=0}^\infty k^{p/2} |\Delta^p v_n(k)|$$

is bounded.

It is familiar that $\Delta^p v_n(k) = (-1)^p v_n^{(p)}(k + c)$ for some $c \in [0, p]$. Hence there is a $\theta = \theta(n, k) \in [0, p]$ such that

$$(2) \quad |\Delta^p v_n(k)| \leq |v_n^{(p)}(k + \theta)|.$$

Since $v_n^{(p)}(x) = v_n(x) \sum_{0 \leq r \leq p/2} b_r(n-x)^{p-2r} n^{r-p}$, where the b_r 's are constants, we get from (1) and (2) that

$$G(n) = O \left(\sum_{0 \leq r \leq p/2} |b_r| n^{r-p-1/2} \sum_{k=0}^{\infty} k^{p/2} |n-k-\theta|^{p-2r} v_n(k+\theta) \right).$$

Therefore to establish that $G(n)$ is bounded it is enough to show that, for $0 \leq r \leq p/2$ and $0 \leq \theta \leq p$,

$$H(n) := \sum_{k=0}^{\infty} k^{p/2} |n-k-\theta|^{p-2r} v_n(k+\theta) = O(n^{p-r+1/2}).$$

Write

$$(3) \quad H(n) = \left\{ \sum_{k=0}^{n-p-1} + \sum_{k=n-p}^n + \sum_{k=n+1}^{\infty} \right\} k^{p/2} |n-k-\theta|^{p-2r} v_n(k+\theta) \\ := \sum_1 + \sum_2 + \sum_3.$$

Since $|n-k-\theta| \leq 2p$ for $0 \leq \theta \leq p$ and $n-p \leq k \leq n$, and $0 < v_n(k+\theta) \leq 1$, it is immediate that

$$(4) \quad \sum_2 = O(n^{p/2}).$$

Next, setting $f(x) := x^{p-2r} \exp(-\alpha x^2/2n)$ and applying Lemma 1, we have that

$$\sum_1 \leq \sum_{k=0}^{n-p-1} k^{p/2} (n-k)^{p-2r} v_n(k+p) \leq \sum_{k=p}^{n-1} k^{p/2} (n-k+p)^{p-2r} v_n(k) \\ \leq M n^{p/2} \sum_{k=p}^{n-1} f(n-k) \leq M n^{p/2} \sum_{k=1}^n f(k) \\ \leq M n^{p/2} \int_1^n f(x) dx + M C n^{p/2} \left(\frac{(p-2r)n}{\alpha} \right)^{p/2-r},$$

where $M := (1+p)^{p-2r}$ and $C := \exp(r-p/2)$. Letting $u = \alpha x^2/2n$, we get that

$$(5) \quad \sum_1 = O \left(n^{p-r+1/2} \int_0^{\infty} u^{(p-1)/2-r} e^{-u} du \right) + O(n^{p-r}) = O(n^{p-r+1/2}).$$

Further, with M and $f(x)$ as above and $g(x) := x^{3p/2-2r} \exp(-\alpha x^2/2n)$, we see that

$$\sum_3 \leq \sum_{k=n+1}^{\infty} k^{p/2} (k-n+p)^{p-2r} v_n(k) \\ \leq M \left(\sum_{k=n+1}^{2n} + \sum_{k=2n+1}^{\infty} \right) k^{p/2} (k-n)^{p-2r} v_n(k) \\ \leq M (2n)^{p/2} \sum_{k=1}^n f(k) + M 2^{p/2} \sum_{k=n+1}^{\infty} g(k) := \sum_{3,1} + \sum_{3,2}.$$

As above $\sum_{3,1} = O(n^{p-r+1/2})$. And finally, as $n \rightarrow \infty$,

$$\begin{aligned} \sum_{3,2} &= O\left(\int_n^\infty g(x)dx\right) + o(1) \\ &= O\left(n^{3p/4-r+1/2} \int_{\alpha n/2}^\infty u^{3p/4-r-1/2} e^{-u} du\right) + o(1) \\ &= o(1). \end{aligned}$$

Thus,

$$(6) \quad \sum_3 = O(n^{p-r+1/2}) + o(1) \quad \text{as } n \rightarrow \infty.$$

It now follows from (3)–(6) that $H(n) = O(n^{p-r+1/2})$. This completes the proof. \square

3. Proof of Theorem 1. The limitation theorem for Cesàro summability [5, Theorem 46] implies that $s_n = o(n^p)$. Therefore, by Theorem 2 and Lemma 2, we have that $\sum_{n=0}^\infty a_n = s(B, \alpha, \beta)$. \square

4. Related results. The methods of Euler E_δ , Meyer-Konig S_δ , and Taylor T_δ ($0 < \delta < 1$) are defined as follows:

$$\begin{aligned} \sum_{n=0}^\infty a_n = s(E_\delta) & \quad \text{if } \sum_{k=0}^n \binom{n}{k} \delta^k (1-\delta)^{n-k} s_k \rightarrow s \quad \text{as } n \rightarrow \infty; \\ \sum_{n=0}^\infty a_n = s(S_\delta) & \quad \text{if } (1-\delta)^{n+1} \sum_{k=0}^\infty \binom{n+k}{k} \delta^k s_k \rightarrow s \quad \text{as } n \rightarrow \infty; \\ \sum_{n=0}^\infty a_n = s(T_\delta) & \quad \text{if } (1-\delta)^{n+1} \sum_{k=0}^\infty \binom{n+k}{k} \delta^k s_{n+k} \rightarrow s \quad \text{as } n \rightarrow \infty. \end{aligned}$$

These methods, as well as the Borel-type and Valiron methods, are contained in the $F(a, q)$ family of methods mentioned in the introduction. The following theorem generalizes Swaminathan’s result [10], via Theorem 2 and [3, Satz III], for the Euler, Meyer-Konig, and Taylor methods.

THEOREM 3. *Suppose that p is a nonnegative integer and that $c_n^p = s + o(n^{-p/2})$ as $n \rightarrow \infty$. Then for $0 < \delta < 1$, the series $\sum_{n=0}^\infty a_n$ is summable to s by the E_δ , S_δ , and T_δ methods.*

REFERENCES

1. D. Borwein, *A Tauberian theorem for Borel-type methods of summability*, *Canad. J. Math.* **21** (1969), 740–747.
2. D. Borwein and T. Markovich, *A Tauberian theorem concerning Borel-type and Cesàro methods of summability*, (submitted).
3. G. Faulhaber, *Äquivalenzsätze für die Kreisverfahren der Limitierungstheorie*, *Math. Z.* **66** (1956), 34–52.
4. G. H. Hardy and J. E. Littlewood, *Theorems concerning the summability of series by Borel’s exponential method*, *Rend. Circ. Mat. Palermo* **41** (1916), 36–53.
5. G. H. Hardy, *Divergent series*, Oxford, 1949.

6. J. M. Hyslop, *On the summability of series by a method of Valiron*, Proc. Edinburgh Math. Soc. (2) **4** (1936), 218–223.
7. ———, *The generalisation of a theorem on Borel summability*, Proc. London Math. Soc. (2) **41** (1936), 243–256.
8. B. Kwee, *An improvement on a theorem of Hardy and Littlewood*, J. London Math. Soc. (2) **28** (1983), 93–102.
9. A. Meir, *Tauberian constants for a family of transformations*, Ann. of Math. (2) **78** (1963), 594–599.
10. V. Swaminathan, *A note on the family $F(a, q)$ of summability methods*, Math. Z. **138** (1974), 119–122.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WESTERN ONTARIO, LONDON, ONTARIO, CANADA N6A 5B7