

INJECTIVE HULLS OF SIMPLE $\mathfrak{sl}(2, \mathbb{C})$ MODULES ARE LOCALLY ARTINIAN

RANDALL P. DAHLBERG

(Communicated by Donald S. Passman)

ABSTRACT. Let L denote the simple Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ over the complex numbers \mathbb{C} . For any simple L -modules S , considered as a left unital module over the universal enveloping algebra of L , $U(L)$, the injective hull of S , $E_L(S)$, is a locally Artinian $U(L)$ -module.

In this paper a module will mean unital left module over the universal enveloping algebra $U(L)$ of L and will be referred to simply as an L -module. A two-sided ideal will be referred to as an ideal. We recall that the standard ordered basis for L is

$$e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

with the relations $[e, f] = h$, $[h, e] = 2e$, and $[h, f] = -2f$. It is well-known that the Casimir element $Q = 4ef + h^2 - 2h$ lies in the center $Z(L)$ of $U(L)$ and that $Z(L) = \mathbb{C}[Q]$. Furthermore, if I is a minimal primitive ideal of $U(L)$ then $I = U(L)(Q - c)$ for some complex number c . [3, 4.9.22, p. 167].

We will need some facts about primitive factor rings of $U(L)$ which we record here as lemmas for our convenience. If R is a ring, the Krull dimension of R will be denoted by $K - \dim(R)$.

Lemma 1. *Let R be a non-Artinian primitive factor ring of $U(L)$. Then $K - \dim(R) = 1$.*

Proof. The case where R is simple follows from the work of Arnal-Pinczon [1] and Roos [6]. If R is not simple, the result was proven by S. P. Smith [7].

Lemma 2 [5, Proposition 5.5, p. 465]. *Let R be a prime ring with $K - \dim(R) = n$. Then either R is simple Artinian or $K - \dim(W) < n$ whenever W is a finitely generated essential extension of a simple R -module.*

Received by the editors October 31, 1988. Presented at the Phoenix Meeting of the AMS on January 13, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 17B10.

©1989 American Mathematical Society
0002-9939/89 \$1.00 + \$.25 per page

Lemma 3 [4, Lemma 2, p. 131]. *Let z be a centralizing element in a left Noetherian ring U and E be a left U -module such that $\text{ann}_E(Uz) = \{e \in E \mid (Uz)e = 0\}$ is Artinian and essential in E . Then E is an Artinian U -module.*

We are now prepared to prove our principal result.

Theorem. *Let S be a simple $U(L)$ -module. Then the injective hull of S is locally Artinian.*

Proof. Let E be a finitely generated essential extension of S . It suffices to show that E is Artinian. Let I be the minimal primitive ideal which annihilates S . Then $I = Uz$ for some central element z in $U = U(L)$. By Lemma 1, $K - \dim(U/I) = 1$. Let $M = \{m \in E \mid Im = 0\}$. Then M is a finitely generated (U/I) -module and S is an essential submodule of M . Hence by Lemma 2, M is Artinian. Thus M is an essential Artinian submodule of E , so E is Artinian by Lemma 3.

Corollary. *Let V be an Artinian L -module. Then $E_L(V)$ is locally Artinian.*

Proof. Since $U(L)$ is Noetherian we may express $E_L(V)$ as a direct sum of indecomposable injective modules. Each summand must intersect V non-trivially so contains an Artinian submodule and hence contains a simple submodule which must then be an essential submodule of the summand. Thus each summand is the injective hull of a simple module so is locally Artinian by the Theorem. But a direct sum of locally Artinian modules is again locally Artinian, hence $E_L(V)$ is locally Artinian.

Remark 1. If L is a solvable Lie algebra over a field k of characteristic zero, then $E_L(V)$ is locally finite dimensional when V is a locally finite dimensional L -module [2, Corollary 12, p. 467]. If L is a semisimple Lie algebra over k then $E_L(V)$ cannot be locally finite dimensional in view of Weyl's Theorem on complete reducibility since any finite dimensional submodule $\neq V$ would contain V as a direct summand by complete reducibility.

Remark 2. If $M(\lambda)$ is a Verma module with a finite dimensional quotient $L(\lambda)$, then its \mathfrak{h} -finite dual $\delta M(\lambda)$ is an essential extension of $L(\lambda)$. As one allows \mathfrak{h} to vary according to automorphisms of $U(\mathfrak{sl}(2, \mathbb{C}))$, and twists $\delta M(\lambda)$ according to the automorphism, one gets further essential extensions of $L(\lambda)$. Note that twisting $L(\lambda)$ by an automorphism leaves $L(\lambda)$ unchanged up to isomorphism since it is completely determined by its dimension.

ACKNOWLEDGMENT

The author would like to thank the reviewer for many helpful suggestions and improvements.

REFERENCES

1. D. Arnal and G. Pinczon, *Idéaux à gauche dans les quotients simples de l'algèbre enveloppante de $\mathfrak{sl}(2)$* , Bull. Soc. Math. France **101** (1973), 381–395.
2. R. P. Dahlberg, *Injective hulls of Lie modules*, J. Algebra **87**, No. 2 (1984), 458–471.
3. J. Dixmier, *Enveloping algebras*, North-Holland, Amsterdam, 1977.
4. A. V. Jategaonkar, *Certain injectives are Artinian*, Proc. Kent State Conf., Lect. Notes in Math. No. 545, Springer-Verlag, Berlin, 1975.
5. I. M. Musson, *Injective modules for group rings of polycyclic groups II*, Quart. J. Math. Oxford (2) **31** (1980), 449–466.
6. J. E. Roos, *Compléments à l'étude des quotients primitifs des algèbres enveloppantes des algèbres de Lie semi-simples*, C. R. Acad. Sci. Paris Sér. A **276** (1973), 447–450.
7. S. P. Smith, *The primitive factor rings of the enveloping algebra of $\mathfrak{sl}(2, \mathbb{C})$* , J. London Math. Soc. (2) **24** (1981), 97–108.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, SETON HALL UNIVERSITY, SOUTH ORANGE, NEW JERSEY 07079