

TWO EXAMPLES OF LOCAL ARTINIAN RINGS

WEIMIN XUE

(Communicated by Donald S. Passman)

ABSTRACT. We answer a question of D. A. Hill in the negative by providing two local artinian rings R and S such that R is right serial but the left indecomposable injective R -module is not uniserial, and that S is not right serial but the left indecomposable injective S -module is uniserial. Moreover R possesses a Morita duality but fails to have self-duality.

A module is *uniserial* in case its submodules are linearly ordered by inclusion. A ring is *left (right) serial* if it is a direct sum of uniserial left (right) modules. It was shown in [5] that an artinian ring R is right serial if and only if every left indecomposable injective R -module is uniserial, provided R is commutative modulo its radical. Hill [5] asked if this result can be extended to arbitrary artinian rings. The answer is “No”. In this note we construct two examples of local artinian rings R and S such that: (1) R is right serial but the left indecomposable injective R -module is not uniserial, and (2) S is not right serial but the left indecomposable injective S -module is uniserial. Moreover the local artinian ring R possesses a (Morita) duality but fails to have self-duality, which solves a question some people are concerned about (see Müller [9, p. 1149], for example). Our examples are based on the profound results of Schofield [10] and Dowbor, Ringel and Simson [4].

Theorem ([10, pp. 214–217] and [4]). *There are division ring extensions $G \leq F$ and $H \leq F$ with $\dim({}_G F) = \dim(F_H) = 3$, $\dim(F_G) = \dim({}_H F) = 2$, and ring isomorphisms $\alpha: F \cong G$ and $\beta: F \cong H$.*

Throughout this note the above notations will be fixed.

Example 1. Let $R = F \times F$ as abelian group and define the multiplication via

$$(a, b)(c, d) = (ab, \beta(a)d + bc).$$

Then R is a local artinian ring with $c({}_R R) = 3$ and $c(R_R) = 2$. Hence R is right serial with radical square zero. Let E_R be the right indecomposable injective R -module; then we have

$$E/\text{Soc}(E_R) \cong \text{Hom}_R({}_R J(R)_R, \text{Soc}(E_R))_R$$

Received by the editors August 22, 1988 and, in revised form, January 23, 1989.
1980 *Mathematics Subject Classification* (1985 Revision). Primary 16A35, 16A49.
Key words and phrases. Uniserial modules, serial rings, Morita duality.

©1989 American Mathematical Society
0002-9939/89 \$1.00 + \$.25 per page

whose length $= \dim(\text{Hom}_F({}_H F_F, F_F)_H) = \dim(F_H) = 3$. It follows that $c(E_R) = 4$. According to Dlab and Ringel [3, Proposition II.3.3], R is not an exceptional ring. It follows from [3, Proposition II.3.2] that the length of the left indecomposable injective R -module ${}_R U$ is greater than 2. Hence ${}_R U$ is not uniserial.

Azumaya [2] and Morita [7] proved that a right artinian ring has a (Morita) duality if and only if every right indecomposable injective module has finite length. A presentation of duality and self-duality can be found in Anderson and Fuller [1, §23, §24]. We note that the local artinian right serial ring R in Example 1 does possess a duality but does not have self-duality since $c(E_R) \neq c({}_R R)$.

Example 2. Let $T = F \times F$ as abelian group and define the multiplication via

$$(a, b)(c, d) = (ab, \alpha(a)d + bc).$$

Then T is a local artinian ring with radical square zero, $c({}_T T) = 4$, and $c(T_T) = 2$. Let V_T be the right indecomposable injective T -module, and $S = \text{End}(V_T)$. Using the same calculation as that of Example 1 we have $c(V_T) = 3$. It follows that ${}_S V_T$ defines a duality. Therefore S is a local artinian ring (the right artinian-ness of S follows from Müller [8, Theorem 10] and a result of [4]) with radical square zero, ${}_S V$ is the left indecomposable injective S -module, $c({}_S V) = c(T_T) = 2$, $c({}_S S) = c(V_T) = 3$, and $T = \text{End}({}_S V)$. So ${}_S V$ is uniserial. Now since $c({}_T T) = 4$, T is not an exceptional ring, neither is S by [3, Proposition III.6.1 and Theorem III.3.1]. According to [3, Proposition II.3.2], $c(S_S) > 2$, since $c({}_S S) = 3$ and $c({}_S V) = 2$. Hence S_S is not uniserial.

ACKNOWLEDGMENT

The author wishes to express his gratitude for the hospitality of the University of Iowa during the preparation of this paper.

REFERENCES

1. F. W. Anderson and K. R. Fuller, *Rings and categories of modules*, Springer-Verlag, Berlin/New York, 1974.
2. G. Azumaya, *A duality theory for injective modules*, Amer. J. Math. **81** (1959), 249–278.
3. V. Dlab and C. M. Ringel, *Balanced rings*, Lecture Notes in Math. 246, Springer-Verlag, Berlin/New York, 1972.
4. P. Dowbor, C. M. Ringel and D. Simson, *Hereditary artinian rings of finite representation type*, Lecture Notes in Math. 832, Springer-Verlag, Berlin/New York, 1980.
5. D. A. Hill, *Rings whose indecomposable injective modules are uniserial*, Canad. J. Math. **34** (1982), 797–805.
6. J. Kraemer, *Characterizations of the existence of (Quasi-) self-duality for complete tensor rings*, Algebra Berichte 56, Verlag Reinhard Fischer, München 1987.
7. K. Morita, *Duality for modules and its applications to the theory of rings with minimum condition*, Tokyo Kyoiku Daigaku, Ser. A6 (1958), 83–142.
8. B. J. Müller, *On Morita duality*, Canad. J. Math. **21** (1969), 1338–1347.

9. —, *The structure of quasi-Frobenius rings*, *Canad. J. Math.* **26** (1974), 1141–1151.
10. A. H. Schofield, *Representations of rings over skew fields*, London Math. Soc. Lecture Note Series 92, Cambridge Univ. Press 1985.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MAINE, ORONO, MAINE 04469

Current address: Department of Mathematics, Fujian Normal University, Fuzhou, Fujian 350007, People's Republic of China