

REMARK ON WALTER'S INEQUALITY FOR SCHUR MULTIPLIERS

MAREK BOŹEJKO

(Communicated by Paul S. Muhly)

ABSTRACT. We extend and give another proof of Walter's inequality: For a linear operator $T \in L(l^2(X))$

$$\|T\|_{V_2(X)}^2 \leq \| |T|_l \|_{V_2(X)} \| |T|_r \|_{V_2(X)},$$

where $V_2(X)$ is the Banach algebra of Schur multipliers on $L(l^2(X))$ and $|T|_l = (T T)^{*1/2}$, $|T|_r = |T^*|_l$.

1. INTRODUCTION

Let X be a set with counting measure. The algebra of all bounded operators on $l^2(X)$, denoted $L(l^2(X))$, can be identified with the functions $k: X \times X \rightarrow C$, called also kernels, in the following way: For $T \in L(l^2(X))$ and $x, y \in X$, $k(x, y) = \langle T \delta_x, \delta_y \rangle$.

After C. Herz [H] and G. Bennett [Be], we say that a kernel $a(x, y)$ is a *Schur multiplier* of $L(l^2(X))$ if for every $k \in L(l^2(X))$, $a \circ k \in L(l^2(X))$ and $\|a\|_{V_2(X)} = \sup\{\|a \cdot k\|: \|k\| = 1\}$ is finite. Here the multiplication $a \circ k$ is the pointwise product of kernels called also the *Schur product* i.e. $(a \circ k)(x, y) = a(x, y)k(x, y)$ and the norm $\| \cdot \|$ is the operator norm on $l^2(X)$.

It was shown by M. Walter [W] that if X is a finite set, then for a kernel $a \in L(l^2(X))$ we have the following inequality:

$$(**) \quad \|a\|_{V_2(X)}^2 < \| |a|_l \|_{V_2(X)} \| |a|_r \|_{V_2(X)},$$

where $|a|_l$ is the left modulus of an operator a defined as follows: $|a|_l = (aa^*)^{1/2}$ and the right modulus $|a|_r = |a^*|_l$. In this note we give another and rather natural proof of Walter's result and we even prove that the inequality (**) holds for all bounded operators for an arbitrary set X .

2. WALTER'S INEQUALITY

Let us start with the following Lemma, which is probably well known, but for completeness we include a proof.

Received by the editors December 4, 1987 and, in revised form, September 20, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 43A65; Secondary 20S05, 22D12, 43A22.

Key words and phrases. Schur multipliers, Schur product.

Lemma. (i) If $b(x, y) = \sum_z A(x, z)B(z, y)$, where

$$B: l^1(X) \longrightarrow l^2(X) \quad \text{and} \quad A: l^2(X) \longrightarrow l^\infty(X),$$

then

$$\|b\|_{V_2(X)} \leq \|B\|_{1,2} \|A\|_{2,\infty}.$$

(ii) If the kernel $a(x, y)$ is positive definite and bounded on $X \times X$, then

$$\|a\|_{V_2(X)} = \sup_{x \in X} a(x, x).$$

(iii) If $a \in L(l^2(X))$, then

$$\|a^* a\|_{V_2(X)} = \|a\|_{1,2}^2, \quad \|aa^*\|_{V_2(X)} = \|a\|_{2,\infty}^2.$$

Proof. (i) From the definition of V_2 -norm we have

$$\|b\|_{V_2(X)} = \sup_{\|k\|=1} \left\{ \left| \sum_{x,y} b(x, y)k(x, y)u(x)v(y) \right| : |u|_2 = |v|_2 = 1, u \text{ and } v \text{ have finite supports} \right\}.$$

Since functions u and v have finite supports, therefore we can interchange the summation and by the Cauchy-Schwarz inequality we get

$$\begin{aligned} \left| \sum_{x,y} b(x, y)u(x)v(y) \right| &= \left| \sum_z \sum_{x,y} k(x, y)A(x, z)u(x)B(z, y)v(y) \right| \\ &\leq \|k\| \sum_z \left(\sum_x |A(x, z)u(x)|^2 \right)^{1/2} \left(\sum_y |B(z, y)v(y)|^2 \right)^{1/2} \\ &\leq \|k\| \left(\sum_{z,x} |A(x, z)u(x)|^2 \right)^{1/2} \left(\sum_{z,y} |B(z, y)v(y)|^2 \right)^{1/2} \\ &\leq \|k\| \|A\|_{2,\infty} \|B\|_{1,2} |u|_2 |v|_2. \end{aligned}$$

The last inequality follows from the well-known fact (see for example [Be]) that

$$\|A\|_{2,\infty} = \sup_x \left(\sum_z |A(x, z)|^2 \right)^{1/2} \quad \text{and} \quad \|B\|_{1,2} = \sup_y \left(\sum_z |B(z, y)|^2 \right)^{1/2}.$$

Proof of (ii). Since $\|a\|_{V_2(X)} \geq \sup_{x,y} |a(x, y)|$ always and the kernel a is positive and bounded, by the Cauchy-Schwarz inequality we have therefore

$$|a(x, y)|^2 \leq a(x, x)a(y, y),$$

which implies that

$$\sup_{x,y} |a(x, y)| = \sup_x |a(x, x)|.$$

On the other hand since

$$a(x, y) = \langle \delta_x, \delta_y \rangle_a, \quad x, y \in X,$$

where the scalar product $\langle \cdot, \cdot \rangle_a$ is induced by the kernel a .

Now we can apply part (i) of our Lemma and we get

$$\|a\|_{V_2(X)} \leq \sup_x \langle \delta_x, \delta_x \rangle_a = \sup_x a(x, x).$$

Hence $\|a\|_{V_2(X)} = \sup_x a(x, x)$.

Proof of (iii). Since a^*a is a positive definite operator, by (ii) we obtain

$$\|a^*a\|_{V_2(X)} = \sup_x \sum_z a^*(x, z)a(z, x) = \|a\|_{1,2}^2.$$

By the duality argument we have $\|aa^*\|_{V_2(X)} = \|a\|_{2,\infty}^2$.

We are now ready to prove the extension of M. Walter's inequality.

Theorem. *If a is a bonded linear operator on $l^2(X)$, then*

- (i) $\|a\|_{V_2(X)}^2 \leq \| |a|_r \|_{V_2(X)} \| |a|_l \|_{V_2(X)}$.
- (ii) For $0 \leq t \leq 2$ we have

$$\|a\|_{V_2(X)}^2 \leq \| |a|_r^t \|_{V_2(X)} \| |a|_l^{(2-t)} \|_{V_2(X)}.$$

Proof. First we give a slightly different interpretation of the Lemma. We showed that if a matrix a is a Schur multiplier of the form $a = AB$, where $B: l_1(X) \rightarrow l_2(X)$ and $A: l_2(X) \rightarrow l_\infty(X)$, then

$$\|a\|_{V_2(X)} \leq \|A\|_{2,\infty} \|B\|_{1,2}.$$

We shall prove (ii) of our Theorem using the polar decomposition of the operator a and finding a quite good factorization. It is well known (see [GK] that $a = u|a|_r = |a|_l u$, where u is a suitable partial isometry operator on $l_2(X)$, and $|a|_r = (a^*a)^{1/2}$ and $|a|_l = (aa^*)^{1/2}$.

Hence one can show that in our case for $t > 0$ we have $u|a|_r^t = |a|_l^t u$. Therefore

$$a = u|a|_r = |a|_l^{(1-t/2)} u |a|_r^{t/2}.$$

Since $\|u\|_{2,2} \leq 1$, by the Lemma we get

$$\begin{aligned} \|a\|_{V_2(X)}^2 &\leq \|u|a|_r^{t/2}\|_{1,2}^2 \| |a|_l^{(1-t/2)} \|_{2,\infty}^2 \\ &\leq \|u\|_{2,2}^2 \| |a|_r^{t/2} \|_{1,2}^2 \| |a|_l^{(2-t)} \|_{V_2(X)} \\ &\leq \| |a|_r^t \|_{V_2(X)} \| |a|_l^{(2-t)} \|_{V_2(X)}. \end{aligned}$$

REFERENCES

[Be] G. Bennett, *Schur multipliers*, Duke Math. J. **44** (1977), 603–639.
 [Bo] M. Bozejko, *Positive definite bounded matrices and a characterization of amenable groups*, Proc. Amer. Math. Soc. **95** (1985), 357–360.

- [GK] I. C. Goh' berg and M. G. Krein, *Introduction to the theory of linear nonselfadjoint operators*, Amer. Math. Soc., Providence, R. I., 1969.
- [H] C. Herz, Une generalization de le notion de transformé le Fourier-Stieltjes, *Ann. Inst. Fourier (Grenoble)*, 24 (1974), 145–157.
- [W] M. E. Walter, *On the Norm of a Schur product*, *Linear Algebra Appl.* 79 (1986), 209–213.

INSTYTUT MATEMATYCZNY UNIWERSYTETU WROCLAWSKIEGO, WROCLAW, PLAC GRUNWALDZKI
2/4, POLAND