

POSITIVE SOLUTIONS OF DIFFERENCE EQUATIONS

CH. G. PHILOS AND Y. G. SFICAS

(Communicated by Kenneth R. Meyer)

ABSTRACT. Consider the difference equation

$$(E) \quad (-1)^{m+1} \Delta^m A_n + \sum_{k=0}^{\infty} p_k A_{n-l_k} = 0,$$

where m is a positive integer, $(p_k)_{k \geq 0}$ is a sequence of positive real numbers and $(l_k)_{k \geq 0}$ is a sequence of integers with $0 \leq l_0 < l_1 < l_2 < \dots$. The characteristic equation of (E) is

$$(*) \quad -(1 - \lambda)^m + \sum_{k=0}^{\infty} p_k \lambda^{-l_k} = 0.$$

We prove the following theorem.

Theorem. (i) For m even, (E) has a positive solution $(A_n)_{n \in \mathbb{Z}}$ with $\limsup_{n \rightarrow \infty} A_n < \infty$ if and only if (*) has a root in $(0, 1)$.

(ii) For m odd, (E) has a positive solution $(A_n)_{n \in \mathbb{Z}}$ if and only if (*) has a root in $(0, 1)$.

1. INTRODUCTION

Recently, there has been a lot of activity concerning the oscillatory behavior of the solutions of difference equations. See, for example, [1,2,3 and 4] and the references cited therein. Our aim in this paper is to obtain necessary and sufficient conditions for the existence of positive solutions of certain difference equations.

Let $Z = \{\dots, -1, 0, 1, \dots\}$. The forward difference operator Δ is defined as usual, i.e.

$$\Delta S_n = S_{n+1} - S_n, \quad n \in Z$$

for any sequence $(S_n)_{n \in Z}$ of real numbers. Moreover, if $(A_n)_{n \in Z}$ is a sequence, we define

$$\Delta^0 A_n = A_n, \quad \text{and} \quad \Delta^i A_n = \Delta(\Delta^{i-1} A_n) \quad (i = 1, 2, \dots)$$

for every $n \in Z$.

Received by the editors February 7, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 39A10.

Key words and phrases. Difference equation, solution, positive solution.

Consider the difference equation

$$(E) \quad (-1)^{m+1} \Delta^m A_n + \sum_{k=0}^{\infty} p_k A_{n-l_k} = 0,$$

where m is a positive integer, $(p_k)_{k \geq 0}$ is a sequence of positive real numbers and $(l_k)_{k \geq 0}$ is a sequence of integers with $0 \leq l_0 < l_1 < l_2 < \dots$.

By a solution of (E) we mean a sequence $(A_n)_{n \in \mathbb{Z}}$ which satisfies (E) for all $n \in \mathbb{Z}$. A solution $(A_n)_{n \in \mathbb{Z}}$ of (E) is called *positive* if $A_n > 0$ for every $n \in \mathbb{Z}$. Moreover, a positive solution $(A_n)_{n \in \mathbb{Z}}$ of (E) is said to be *bounded at ∞* if $\limsup_{n \rightarrow \infty} A_n < \infty$.

The characteristic equation of (E) is

$$(*) \quad -(1 - \lambda)^m + \sum_{k=0}^{\infty} p_k \lambda^{-l_k} = 0.$$

In this paper we prove the following result.

Theorem. (i) For m even, (E) has a positive solution which is bounded at ∞ if and only if (*) has a root in $(0, 1)$.

(ii) For m odd, (E) has a positive solution if and only if (*) has a root in $(0, 1)$.

To prove our theorem we need two lemmas. These lemmas are established in §2. The proof of the theorem will be given in §3.

It is easy to verify that

$$\sup_{\lambda \in (0, 1)} [(1 - \lambda)^m \lambda^l] = \frac{m^m l^l}{(m + l)^{m+l}} \quad \text{for } l \in \{0, 1, \dots\}.$$

(Here, we use the convention that $0^0 = 1$.) Hence, for every $\lambda \in (0, 1)$

$$\begin{aligned} -(1 - \lambda)^m + \sum_{k=0}^{\infty} p_k \lambda^{-l_k} &= (1 - \lambda)^m \left[-1 + \sum_{k=0}^{\infty} p_k \frac{1}{(1 - \lambda)^m \lambda^{l_k}} \right] \\ &\geq (1 - \lambda)^m \left[-1 + \sum_{k=0}^{\infty} p_k \frac{(m + l_k)^{m+l_k}}{m^m l_k^{l_k}} \right] \end{aligned}$$

and so the assumption

$$(C) \quad \sum_{k=0}^{\infty} p_k \frac{(m + l_k)^{m+l_k}}{m^m l_k^{l_k}} > 1$$

implies that (*) has no roots in $(0, 1)$. Therefore, our theorem leads to the following corollary.

Corollary. Suppose that (C) holds. Then:

(i) For m even, there is no positive solution of (E) which is bounded at ∞ .

(ii) For m odd, there is no positive solution of (E).

The oscillatory behavior of solutions of the difference equation

$$(-1)^{m+1} \Delta^m A_n + \sum_{k=0}^N p_k A_{n-l_k} = 0,$$

where N is a non-negative integer, p_k ($k = 0, 1, \dots, N$) are positive constants and l_k ($k = 0, 1, \dots, N$) are integers with $0 \leq l_0 < l_1 < \dots < l_N$, can be studied by a detailed analysis of the representation of the solutions in terms of the roots of the characteristic equation. The above equation is the discrete version of the delay differential equation

$$(-1)^{m+1} x^{(m)}(t) + \sum_{k=0}^N p_k x(t - \tau_k) = 0,$$

where $N \geq 0$ is an integer, the coefficients p_0, p_1, \dots, p_N are positive numbers and the delays are constants such that $0 \leq \tau_0 < \tau_1 < \dots < \tau_N$. The oscillation of solutions of this differential equation is treated in [5]. However, this is the first paper dealing with difference equations of the form (E). For such difference equations no representation of solutions in terms of the roots of (*) is known.

2. LEMMAS

The following two lemmas will be useful in §3.

Lemma 1. *Let $(A_n)_{n \in \mathbb{Z}}$ be a positive solution of (E) which is bounded at ∞ . Then*

$$(-1)^j \Delta^j A_n > 0 \quad \text{for all } n \in \mathbb{Z} \quad (j = 0, 1, \dots, m - 1, m).$$

Proof. From (E) we obtain for $n \in \mathbb{Z}$

$$(-1)^m \Delta^m A_n = \sum_{k=0}^{\infty} p_k A_{n-l_k}$$

and consequently

$$(1) \quad (-1)^m \Delta^m A_n > 0 \quad \text{for all } n \in \mathbb{Z}.$$

For $m = 1$ the proof is complete. So, we assume that $m > 1$. We now claim that

$$(2) \quad (-1)^{m-1} \Delta^{m-1} A_n > 0 \quad \text{for every } n \in \mathbb{Z}.$$

Otherwise, there exists an integer n_1 with

$$(-1)^{m-1} (\Delta^{m-1} A_n)_{n=n_1} \leq 0.$$

From (1) it follows that the sequence $((-1)^{m-1} \Delta^{m-1} A_n)_{n \in \mathbb{Z}}$ is strictly decreasing. Hence, if we choose an integer $n_2 > n_1$, then we derive for $n \geq n_2$

$$(-1)^{m-1} \Delta^{m-1} A_n \leq (-1)^{m-1} (\Delta^{m-1} A_n)_{n=n_2} < (-1)^{m-1} (\Delta^{m-1} A_n)_{n=n_1} \leq 0.$$

Therefore

$$(-1)^{m-1} \Delta^{m-1} A_n \leq -\gamma \quad \text{for every } n \geq n_2,$$

where $\gamma = -(-1)^{m-1}(\Delta^{m-1}A_n)_{n=n_2} > 0$. So, we obtain for $n > n_2$

$$\begin{aligned} &(-1)^{m-1}\Delta^{m-2}A_n - (-1)^{m-1}(\Delta^{m-2}A_n)_{n=n_2} \\ &= [(-1)^{m-1}\Delta^{m-2}A_n - (-1)^{m-1}\Delta^{m-2}A_{n-1}] \\ &\quad + [(-1)^{m-1}\Delta^{m-2}A_{n-1} - (-1)^{m-1}\Delta^{m-2}A_{n-2}] \\ &\quad + \\ &\quad \vdots \\ &\quad + [(-1)^{m-1}(\Delta^{m-2}A_n)_{n=n_2+1} - (-1)^{m-1}(\Delta^{m-2}A_n)_{n=n_2}] \\ &= (-1)^{m-1}\Delta^{m-1}A_{n-1} + (-1)^{m-1}\Delta^{m-1}A_{n-2} + \dots \\ &\quad \dots + (-1)^{m-1}(\Delta^{m-1}A_n)_{n=n_2} \\ &\leq -\gamma(n - n_2), \end{aligned}$$

which gives

$$(3) \quad \lim_{n \rightarrow \infty} (-1)^{m-1}\Delta^{m-2}A_n = -\infty.$$

Since $(A_n)_{n \in \mathbb{Z}}$ is bounded at ∞ , (3) is a contradiction if $m = 2$. So, we suppose that $m > 2$ and we consider a positive constant γ_1 . Then (3) implies the existence of an integer n_3 such that

$$(-1)^{m-1}\Delta^{m-2}A_n \leq -\gamma_1 \quad \text{for every } n \geq n_3.$$

Thus, by applying the method used previously, we can obtain

$$\lim_{n \rightarrow \infty} (-1)^{m-1}\Delta^{m-3}A_n = -\infty.$$

Next, repeating the above procedure if $m > 3$, we finally find

$$\lim_{n \rightarrow \infty} (-1)^{m-1}A_n = -\infty,$$

which contradicts the fact that $(A_n)_{n \in \mathbb{Z}}$ is positive and that this sequence is bounded at ∞ . We have thus proved that (2) is true. If $m = 2$, the proof of the lemma is complete. If $m > 2$, then, repeating the above arguments, we can show that

$$(-1)^{m-2}\Delta^{m-2}A_n > 0 \quad \text{for all } n \in \mathbb{Z}.$$

By using this technique, we can complete the proof of the lemma.

Lemma 2. *Let m be odd. Then every positive solution of (E) is bounded at ∞ .*

Proof. Assume, for the sake of contradiction, that (E) has a positive solution $(A_n)_{n \in \mathbb{Z}}$ which is not bounded at ∞ . Since m is odd, from (E) it follows that

$$(4) \quad \Delta^m A_n < 0 \quad \text{for all } n \in \mathbb{Z}.$$

Thus, we always have $m > 1$. Furthermore, (4) implies that, if $j \in \{1, \dots, m - 1\}$, then $\Delta^j A_n$ is either positive for all large n or negative for all large n .

In particular, since $(A_n)_{n \in \mathbb{Z}}$ is not bounded at ∞ , there exists an integer n_0 such that

$$(5) \quad \Delta A_n > 0 \quad \text{for every } n \geq n_0.$$

Furthermore, we have

$$(6) \quad \Delta^{m-1} A_n > 0 \quad \text{for all } n \in \mathbb{Z}.$$

Indeed, in the opposite case we can apply the method used in the proof of Lemma 1 to obtain $\lim_{n \rightarrow \infty} (-1)^{m-1} A_n = -\infty$. So, since m is odd,

$$\lim_{n \rightarrow \infty} A_n = -\infty.$$

This contradicts the positiveness of $(A_n)_{n \in \mathbb{Z}}$ and hence (6) is true. Now, we observe that, by (5), the sequence $(A_n)_{n \geq n_0}$ is strictly increasing. By using this fact, (6) and (E), for $n > N \equiv n_0 + l_0$ we obtain

$$\begin{aligned} -(\Delta^{m-1} A_n)_{n=N} &< \Delta^{m-1} A_n - (\Delta^{m-1} A_n)_{n=N} \\ &= (\Delta^{m-1} A_n - \Delta^{m-1} A_{n-1}) + (\Delta^{m-1} A_{n-1} - \Delta^{m-1} A_{n-2}) + \dots \\ &\quad \dots + [(\Delta^{m-1} A_n)_{n=N+1} - (\Delta^{m-1} A_n)_{n=N}] \\ &= \Delta^m A_{n-1} + \Delta^m A_{n-2} + \dots + (\Delta^m A_n)_{n=N} \\ &= -\sum_{k=0}^{\infty} p_k A_{n-1-l_k} - \sum_{k=0}^{\infty} p_k A_{n-2-l_k} - \dots - \sum_{k=0}^{\infty} p_k A_{N-l_k} \\ &< -p_0 A_{n-1-l_0} - p_0 A_{n-2-l_0} - \dots - p_0 A_{N-l_0} \\ &\leq -p_0 A_{n_0} (n - N) \end{aligned}$$

and so we arrive at the contradiction

$$n < N + \frac{1}{p_0 A_{n_0}} (\Delta^{m-1} A_n)_{n=N} \quad \text{for every } n > N.$$

3. PROOF OF THE THEOREM

In view of Lemma 2, for m odd there is no positive solution of (E) which is not bounded at ∞ . Hence, it is enough to prove part (ii) of our theorem only for positive solutions which are bounded at ∞ . Therefore, the proof of the theorem has been reduced to proving, for arbitrary m , the following result: Equation (E) has a positive solution which is bounded at ∞ if and only if (*) has a root in $(0, 1)$.

Assume first that (*) has a root $\lambda \in (0, 1)$. Then we set $A_n = \lambda^n$, $n \in \mathbb{Z}$ and we obtain

$$\begin{aligned} (-1)^{m+1} \Delta^m A_n + \sum_{k=0}^{\infty} p_k A_{n-l_k} &= (-1)^{m+1} (\lambda - 1)^m \lambda^n + \sum_{k=0}^{\infty} p_k \lambda^{n-l_k} \\ &= [-(1 - \lambda)^m + \sum_{k=0}^{\infty} p_k \lambda^{-l_k}] \lambda^n = 0 \end{aligned}$$

for all $n \in Z$. Thus, $(A_n)_{n \in Z}$ is a positive solution of (E) which obviously is bounded at ∞ .

Suppose, conversely, that there is a positive solution $(A_n)_{n \in Z}$ of (E) which is bounded at ∞ . Also assume, for the sake of contradiction, that the characteristic equation (*) has no roots in $(0, 1)$. From Lemma 1 it follows that $\Delta A_n < 0$ for all $n \in Z$ and consequently $(A_n)_{n \in Z}$ is a strictly decreasing sequence. So, from (E) we obtain for every $n \in Z$

$$0 = (-1)^{m+1} \Delta^m A_n + \sum_{k=0}^{\infty} p_k A_{n-l_k} > (-1)^{m+1} \Delta^m A_n + \left(\sum_{k=0}^{\infty} p_k \right) A_n$$

and therefore

$$(7) \quad 0 < \sum_{k=0}^{\infty} p_k < \infty.$$

Set

$$F(\lambda) = -(1 - \lambda)^m + \sum_{k=0}^{\infty} p_k \lambda^{-l_k} \quad \text{for } \lambda \in (0, 1].$$

Then

$$F(1) = \sum_{k=0}^{\infty} p_k \in (0, \infty).$$

Moreover, we have $F(\lambda) > -(1 - \lambda)^m + p_1 \lambda^{-l_1}$ for every $\lambda \in (0, 1)$, and so

$$F(0+0) = \infty.$$

Hence, as $F(\lambda) = 0$ has no roots in $(0, 1)$, there exists a positive number μ such that

$$(8) \quad -(1 - \lambda)^m + \sum_{k=0}^{\infty} p_k \lambda^{-l_k} \geq \mu \quad \text{for all } \lambda \in (0, 1).$$

Next, by taking into account (7), we put

$$\lambda_0 = 1 - \left(\sum_{k=0}^{\infty} p_k \right)^{1/m}, \quad \text{and } \lambda_r = 1 - [(1 - \lambda_{r-1})^m + \mu]^{1/m} \quad (r = 1, 2, \dots).$$

Furthermore, we define

$$A_n^{[0]} = A_n \quad \text{for } n \in Z$$

and

$$A_n^{[r]} = \sum_{j=0}^{m-1} (1 - \lambda_{r-1})^{m-1-j} (-1)^j \Delta^j A_n^{[r-1]} \quad \text{for } n \in Z \quad (r = 1, 2, \dots).$$

Then, for any $r \in \{0, 1, \dots\}$, $(A_n^{[r]})_{n \in Z}$ is a positive solution of (E) which is bounded at ∞ . Indeed, consider a positive solution $(\tilde{A}_n)_{n \in Z}$ of (E) which is bounded at ∞ . In view of Lemma 1, we have

$$(-1)^j \Delta^j \tilde{A}_n > 0 \quad \text{for all } n \in Z \quad (j = 0, 1, \dots, m - 1, m).$$

Thus, if $j \in \{0, 1, \dots, m - 1\}$, then $((-1)^j \Delta^j \tilde{A}_n)_{n \in Z}$ is a positive sequence which is strictly decreasing (and, therefore, bounded at ∞). Moreover, since (E) is linear and the coefficients p_k ($k = 0, 1, \dots$) and the indices l_k ($k = 0, 1, \dots$) are independent of n , it follows that, for each $j \in \{0, 1, \dots, m - 1\}$, the sequence $((-1)^j \Delta^j \tilde{A}_n)_{n \in Z}$ is a solution of (E). Hence, each one of the sequences $((-1)^j \Delta^j \tilde{A}_n)_{n \in Z}$ ($j = 0, 1, \dots, m - 1$) is a positive solution of (E) which is bounded at ∞ . Therefore, because of the linearity of (E), it follows that, if c_0, c_1, \dots, c_{m-1} are positive constants, then the sequence $(\sum_{j=0}^{m-1} c_j (-1)^j \Delta^j \tilde{A}_n)_{n \in Z}$ is a positive solution of (E) which is bounded at ∞ . Now, we can easily see that $1 - \lambda_r > 0$ ($r = 0, 1, \dots$). So, by the above particular result and by mathematical induction, we can show that: If $r \in \{0, 1, \dots\}$, then $(A_n^{[r]})_{n \in Z}$ is a positive solution of (E) which is bounded at ∞ .

We have

$$(9) \quad (-1)^{m+1} \Delta^m A_n^{[r]} + (1 - \lambda_r)^m A_n^{[r]} = A_{n+1}^{[r+1]} - \lambda_r A_n^{[r+1]} \quad \text{for } n \in Z \quad (r = 0, 1, \dots).$$

In fact, for any $r \in \{0, 1, \dots\}$ and every $n \in Z$, we obtain

$$\begin{aligned} & A_{n+1}^{[r+1]} - \lambda_r A_n^{[r+1]} \\ &= \sum_{j=0}^{m-1} (1 - \lambda_r)^{m-1-j} (-1)^j \Delta^j A_{n+1}^{[r]} - \lambda_r \sum_{j=0}^{m-1} (1 - \lambda_r)^{m-1-j} (-1)^j \Delta^j A_n^{[r]} \\ &= \sum_{j=0}^{m-1} (1 - \lambda_r)^{m-1-j} (-1)^j (\Delta^j A_{n+1}^{[r]} - \Delta^j A_n^{[r]}) \\ &\quad + (1 - \lambda_r) \sum_{j=0}^{m-1} (1 - \lambda_r)^{m-1-j} (-1)^j \Delta^j A_n^{[r]} \\ &= \sum_{j=0}^{m-1} (1 - \lambda_r)^{m-1-j} (-1)^j \Delta^{j+1} A_n^{[r]} + \sum_{j=0}^{m-1} (1 - \lambda_r)^{m-j} (-1)^j \Delta^j A_n^{[r]} \\ &= - \sum_{j=1}^m (1 - \lambda_r)^{m-j} (-1)^j \Delta^j A_n^{[r]} + \sum_{j=0}^{m-1} (1 - \lambda_r)^{m-j} (-1)^j \Delta^j A_n^{[r]} \\ &= (-1)^{m+1} \Delta^m A_n^{[r]} + (1 - \lambda_r)^m A_n^{[r]}. \end{aligned}$$

Now, we will prove that

$$(10) \quad A_{n+1}^{[r+1]} - \lambda_r A_n^{[r+1]} < 0 \quad \text{for all } n \in Z \quad (r = 0, 1, \dots).$$

Indeed, in view of Lemma 1, the sequence $(A_n)_{n \in \mathbb{Z}}$ is strictly decreasing. Therefore, we obtain for every $n \in \mathbb{Z}$

$$\begin{aligned} (-1)^{m+1} \Delta^m A_n^{[0]} + (1 - \lambda_0)^m A_n^{[0]} &= (-1)^{m+1} \Delta^m A_n + \left(\sum_{k=0}^{\infty} p_k \right) A_n \\ &< (-1)^{m+1} \Delta^m A_n + \sum_{k=0}^{\infty} p_k A_{n-l_k} = 0 \end{aligned}$$

and hence, by (9), we conclude that (10) is true for $r = 0$. Next, assuming that (10) holds for some $r \in \{0, 1, \dots\}$ we should prove that it is also true for $r + 1$. By the inductive assumption,

$$A_{n+1}^{[r+1]} - \lambda_r A_n^{[r+1]} < 0 \quad \text{for every } n \in \mathbb{Z}.$$

This implies in particular that $\lambda_r > 0$. On the other hand, $\lambda_r < 1$. So, we must have $0 < \lambda_r < 1$. Furthermore, we have

$$A_n^{[r+1]} > \lambda_r^{-1} A_{n+1}^{[r+1]} \quad \text{for } n \in \mathbb{Z}.$$

By applying this inequality, we can verify that

$$A_{n-l_0}^{[r+1]} \geq \lambda_r^{-l_0} A_n^{[r+1]} \quad \text{for all } n \in \mathbb{Z}$$

and

$$A_{n-l_k}^{[r+1]} > \lambda_r^{-l_k} A_n^{[r+1]} \quad \text{for all } n \in \mathbb{Z} \quad (k = 1, 2, \dots).$$

Hence, from (E) we obtain for $n \in \mathbb{Z}$

$$\begin{aligned} 0 &= (-1)^{m+1} \Delta^m A_n^{[r+1]} + \sum_{k=0}^{\infty} p_k A_{n-l_k}^{[r+1]} \\ &> (-1)^{m+1} \Delta^m A_n^{[r+1]} + \left(\sum_{k=0}^{\infty} p_k \lambda_r^{-l_k} \right) A_n^{[r+1]}. \end{aligned}$$

But (8) ensures that

$$\sum_{k=0}^{\infty} p_k \lambda_r^{-l_k} \geq (1 - \lambda_r)^m + \mu = (1 - \lambda_{r+1})^m.$$

So, in view of (9), we have for every $n \in \mathbb{Z}$

$$\begin{aligned} 0 &> (-1)^{m+1} \Delta^m A_n^{[r+1]} + (1 - \lambda_{r+1})^m A_n^{[r+1]} \\ &= A_{n+1}^{[r+2]} - \lambda_{r+1} A_n^{[r+2]}. \end{aligned}$$

That is, (10) is also satisfied for $r + 1$.

Finally, since $A_n^{[r+1]} > 0$ for all $n \in \mathbb{Z}$ ($r = 0, 1, \dots$), from (10) it follows that

$$\lambda_r > 0 \quad (r = 0, 1, \dots).$$

On the other hand, it is easy to verify that the sequence $(\lambda_r)_{r=0,1,\dots}$ is strictly decreasing. So, $L \equiv \lim_{r \rightarrow \infty} \lambda_r$ exists and $0 \leq L < \lambda_0 < 1$. Since

$$\lambda_r = 1 - [(1 - \lambda_{r-1})^m + \mu]^{1/m} \quad (r = 1, 2, \dots),$$

we obtain

$$L = 1 - [(1 - L)^m + \mu]^{1/m},$$

which gives $\mu = 0$, a contradiction. The proof of our theorem is complete.

ACKNOWLEDGMENT

The authors wish to thank Professor G. Ladas for helpful conversations on the subject of this paper.

REFERENCES

1. L. H. Erbe and B. G. Zhang, *Oscillation of discrete analogues of delay equations*, *Differential and Integral Equations* **2** (1989), 300–309.
2. I. Györi and G. Ladas, *Linearized oscillations for equations with piecewise constant arguments*, *Differential and Integral Equations* **2** (1989), 123–131.
3. G. Ladas, *Oscillations of equations with piecewise constant mixed arguments*, *Proceedings of the International Conference on Theory and Applications of Differential Equations*, March 21–25, 1988, Ohio University.
4. G. Ladas, Ch. G. Philos and Y. G. Sficas, *Necessary and sufficient conditions for the oscillation of difference equations* *Libertas Math.* **9** (1989).
5. G. Ladas, Y. G. Sficas and I. P. Stavroulakis, *Necessary and sufficient conditions for oscillations of higher order delay differential equations*, *Trans. Amer. Math. Soc.* **285** (1984), 81–90.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOANNINA, P. O. BOX. 1186, 451 10 IOANNINA, GREECE