

DIRICHLET-FINITE ANALYTIC AND HARMONIC FUNCTIONS ARE BMO

J. L. SCHIFF

(Communicated by Paul S. Muhly)

ABSTRACT. Based on a result of F. John, an elementary proof is given of the fact that Dirichlet-finite analytic and Dirichlet-finite harmonic functions are of bounded mean oscillation in the unit disk.

1. In [5] Metzger proved the rather surprising result that the space of Dirichlet-finite analytic functions on a hyperbolic Riemann surface belong to the space BMO. Subsequently, in [4], Kusunoki and Taniguchi found that the same result holds for Dirichlet-finite harmonic functions on Riemann surfaces of finite type. In this note we give an elementary proof, based on a result of John [2], that in the unit disk Dirichlet-finite analytic and harmonic functions are BMO.

2. Denote the unit disk by $U : |z| < 1$, and

$$AD(U) = \left\{ f \in A(U) : D_U(f) = \int_U \int |f'(z)|^2 dx dy < \infty \right\},$$
$$HD(U) = \left\{ u \in H(U) : D_U(u) = \int_U \int |\text{grad } u|^2 dx dy < \infty \right\},$$

as the spaces of Dirichlet-finite analytic, and Dirichlet-finite harmonic, functions, respectively.

Let $BMOA(U)$ be the space of analytic functions on U which belong to the Hardy class $H^2(U)$ and satisfy

$$\sup_{\zeta \in U} \int_U \int |f'(z)|^2 \log \left| \frac{1 - \bar{\zeta}z}{z - \zeta} \right| dx dy < \infty,$$

and define $BMOH(U)$ analogously for harmonic functions, replacing $|f'|^2$ by $|\text{grad } u|^2$.

A sufficient condition for a real-valued differentiable function u to belong to $BMO(U)$ in the sense of John-Nirenberg [3] has been given by John [2], namely: $\sup_{z \in U} (1 - |z|) |\text{grad } u(z)| < \infty$.

Received by the editors July 19, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 30D99; Second 30D55.

© 1990 American Mathematical Society
0002-9939/90 \$1.00 + \$.25 per page

We can now readily establish the following:

Theorem. $AD(U) \subseteq BMOA(U)$.

Proof. Firstly, $AD(U) \subseteq H^2(U)$ (cf. Heins [1]). Take $f \in AD(U)$, $f = u + iv$. By the areal mean value property applied to $\operatorname{Re}[(f')^2]$, $\operatorname{Im}[(f')^2]$,

$$(f'(z))^2 = \frac{1}{\pi \rho^2} \int_0^\rho \int_0^{2\pi} [f'(z + re^{i\theta})]^2 r dr d\theta,$$

where $\rho = 1 - |z|$. Then

$$|\operatorname{grad} u(z)|^2 = |f'(z)|^2 \leq \frac{1}{\pi(1 - |z|)^2} \int_U \int |f'(z)|^2 dx dy = \frac{1}{\pi(1 - |z|)^2} D_U(f).$$

Therefore,

$$\sup_{z \in U} (1 - |z|) |\operatorname{grad} u(z)| \leq \sqrt{\frac{1}{\pi} D_U(f)} < \infty.$$

Since $u \in h^2(U)$, the John result implies $u \in BMOH(U)$, and hence $f \in BMOA(U)$.

For any function $u \in HD(U)$, $u \in \operatorname{Re} f$ for some $f \in AD(U)$, and the preceding proof yields:

Corollary. $HD(U) \subseteq BMOH(U)$.

REFERENCES

1. M. Heins, *Hardy classes on Riemann surfaces*, Lecture Notes in Math., 98, Springer-Verlag, 1969.
2. F. John, *Functions whose gradients are bounded by the reciprocal distance from the boundary of their domain*, Russian Math. Surveys **29** (1974), 170–175.
3. F. John and L. Nirenberg, *On functions of bounded mean oscillation*, Comm. Pure Appl. Math. **14** (1961), 415–426.
4. Y. Kusunoki and M. Taniguchi, *Remarks on functions of bounded mean oscillation on Riemann surfaces*, Kodai Math. J. **6** (1983), 434–442.
5. T. Metzger, *On BMOA for Riemann surfaces*, Canad. Math. J. **33** (1981), 1255–1260.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AUCKLAND, AUCKLAND, NEW ZEALAND

Current address: Department of Mathematics, University of York, Heslington, York YO1 5DD England