

A REMARK ON A PAPER BY C. FEFFERMAN

FILIPPO CHIARENZA AND MICHELE FRASCA

(Communicated by John B. Conway)

ABSTRACT. We give a simplified proof of an imbedding theorem by C. Fefferman [3].

The purpose of this paper is to provide a simplified proof of a deep result by C. Fefferman (see [3,1]) concerning the imbedding

$$(1.1) \quad \int_{\mathbf{R}^n} |u(x)|^p V(x) dx \leq c \int_{\mathbf{R}^n} |\nabla u(x)|^p dx, \quad \forall u \in C_0^\infty(\mathbf{R}^n).$$

In fact (1.1) was proved in [3], for $p = 2$, assuming

$$V \in L^{r, n-2r}(\mathbf{R}^n) \quad 1 < r \leq n/2.$$

Here $L^{r, n-2r}(\mathbf{R}^n) \equiv L^{r, n-2r}$ is the classical Morrey space of the $L_{\text{loc}}^r(\mathbf{R}^n)$ functions such that

$$\sup_{\substack{x \in \mathbf{R}^n \\ \rho > 0}} \frac{1}{\rho^{n-2r}} \int_{B(x, \rho)} |V(y)|^r dy \equiv \|V\|_{r, n-2r}^r < +\infty$$

where we set $B(x, \rho) = \{y \in \mathbf{R}^n : |x - y| \leq \rho\}$.

Our proof rests on the following nice feature of the space $L^{r, n-2r}$: given $V \in L^{r, n-2r}$ there exists an A_1 weight in the same class majorizing V . Such a property is not shared by $L^{1, n-2}$ which is well known to be necessary but not sufficient for (1.1) to hold.

Our result is the following:

Theorem. Let $1 < p < n$, $1 < r \leq n/p$, $V \in L^{r, n-pr}$. Then

$$(1.2) \quad \int_{\mathbf{R}^n} |u(x)|^p V(x) dx \leq c \|V\|_{r, n-pr} \int_{\mathbf{R}^n} |\nabla u(x)|^p dx, \quad \forall u \in C_0^\infty(\mathbf{R}^n).$$

Here c depends on n and p only.

Proof. To prove (1.2) we assume for a moment that $V \in A_1$ (i.e. $MV(x) \leq cV(x)$ a.e. in \mathbf{R}^n , where MV is the usual Hardy–Littlewood maximal function). This assumption will be removed later (see Lemma 1).

Received by the editors March 10, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 46E35; Secondary 42B25.

This work was partially financially supported by a national project of the Italian Ministero della Pubblica Istruzione.

For any fixed $u \in C_0^\infty(\mathbf{R}^n)$ let B be a ball such that $u \in C_0^\infty(B)$. Consider the solution z of the Dirichlet problem: $-\Delta z = V$ in B , $z = 0$ on ∂B . Then

$$(1.3) \quad \int_{\mathbf{R}^n} |u(x)|^p V(x) dx \leq p \int_{\mathbf{R}^n} |u(x)|^{p-1} |\nabla u(x)| |\nabla z(x)| dx.$$

Now we observe that using an idea of Hedberg [6] it is easy to prove that the assumption $V \in L^{r, n-pr}$ implies the punctual estimate

$$(1.4) \quad |\nabla z(x)| \leq c(n, r, p) [MV(x)]^{(p-1)/p} \|V\|_{r, n-pr}^{1/p}.$$

For the sake of completeness we will prove (1.4) in Lemma 2 below (see also the proof of Theorem 2 in [2]). Substituting in (1.3) we have

$$\begin{aligned} \int_B |u(x)|^p V(x) dx &\leq c \|V\|_{r, n-pr}^{1/p} \int_B |u(x)|^{p-1} |\nabla u(x)| [MV(x)]^{(p-1)/p} dx \\ &\leq c \|V\|_{r, n-pr}^{1/p} \left(\int_B |u(x)|^p MV(x) dx \right)^{1-1/p} \left(\int_B |\nabla u(x)|^p dx \right)^{1/p} \\ &\leq c \|V\|_{r, n-pr}^{1/p} \left(\int_B |u(x)|^p V(x) dx \right)^{1-1/p} \left(\int_B |\nabla u(x)|^p dx \right)^{1/p}, \end{aligned}$$

where in the last inequality we used the A_1 assumption on V . The conclusion now follows.

The theorem will be completely proved after we prove the following lemmas.

Lemma 1. *Let $V \in L^{r, n-pr}$, $1 < p < n$, $1 < r \leq n/p$. Let $r_1: 1 < r_1 < r$. Then $(MV^{r_1})^{1/r_1} \in A_1 \cap L^{r, n-pr}$.*

Proof. $(MV^{r_1})^{1/r_1} \in A_1$ clearly (see [5], p. 158). Using the inequality proved in [4] (Lemma 1, p. 111) we have

$$(1.5) \quad \int_{\mathbf{R}^n} (MV^{r_1})^{r/r_1} \chi(x) dx \leq c \int_{\mathbf{R}^n} [|V(x)|^{r_1}]^{r/r_1} M\chi(x) dx,$$

for any χ nonnegative function. Now take as $\chi(x)$ the characteristic function of a ball $B_\rho \equiv B_\rho(x_0)$. We have from (1.5)

$$\begin{aligned} \int_{B_\rho} (MV^{r_1})^{r/r_1} dx &\leq c \left\{ \int_{B_{2\rho}} |V(x)|^r M\chi dx + \sum_{k=1}^{+\infty} \int_{B_{2^{k+1}\rho} \setminus B_{2^k\rho}} |V(x)|^r M\chi dx \right\} \\ &\leq c \left\{ \int_{B_{2\rho}} |V(x)|^r M\chi dx + \sum_{k=1}^{+\infty} \int_{B_{2^{k+1}\rho} \setminus B_{2^k\rho}} |V(x)|^r \frac{\rho^n}{(|x-x_0|-\rho)^n} dx \right\} \end{aligned}$$

Finally

$$\begin{aligned} \int_{B_\rho} [MV^{r_1}]^{r/r_1} dx &\leq c \|V\|_{r, n-pr}^r \left\{ (2\rho)^{n-pr} + \sum_{k=1}^{+\infty} \frac{1}{(2^{k-1})^n} (2^{k+1}\rho)^{n-pr} \right\} \\ &\leq c \|V\|_{r, n-pr}^r \rho^{n-pr}. \end{aligned}$$

Lemma 2. Let $V \in L^{r, n-pr}$, $1 < p < n$, $1 < r \leq n/p$. Then

$$|I_1(V)(x)| = \left| \int_{\mathbb{R}^n} \frac{V(y)}{|x-y|^{n-1}} dy \right| \leq c(n, r, p)[MV(x)]^{(p-1)/p} \|V\|_{r, n-pr}^{1/p}.$$

Proof. Let $\rho > 0$. Then

$$I_1(V)(x) = \int_{|x-y| \leq \rho} \frac{V(y)}{|x-y|^{n-1}} dy + \int_{|x-y| > \rho} \frac{V(y)}{|x-y|^{n-1}} dy \equiv I' + I''.$$

By Hedberg [6] we have

$$|I'| \leq c_n \rho MV.$$

For I'' , setting $\sigma = n - (r/2)(p + 1)$, we have

$$\begin{aligned} |I''| &\leq \left(\int_{|x-y| > \rho} \frac{|V(y)|^r}{|x-y|^\sigma} dy \right)^{1/r} \left(\int_{|x-y| > \rho} |x-y|^{(\sigma/r+1-n)r/(r-1)} dy \right)^{1-1/r} \\ &\equiv J' \cdot J''. \end{aligned}$$

For J' using the assumption on V and the trick by Hedberg we have

$$J' \leq c \rho^{(1-p)/2-1} \|V\|_{r, n-pr}.$$

Finally calculating J'' we get

$$|I_1 V| \leq c' \rho MV + c'' \rho^{1-p} \|V\|_{r, n-pr}.$$

For $\rho = \left(\frac{MV}{\|V\|_{r, n-pr}} \right)^{-1/p}$ we have

$$|I_1 V| \leq c(MV)^{1-1/p} \|V\|_{r, n-pr}^{1/p}.$$

REFERENCES

1. S. Y. A. Chang, J. M. Wilson, and T. H. Wolf, *Some weighted norm inequalities concerning the Schrodinger operators*, Comment. Math. Helvetici **60** (1985), 217-246.
2. F. Chiarenza and M. Frasca, *Morrey spaces and Hardy-Littlewood maximal function*, Rend. Mat. (to appear).
3. C. Fefferman, *The uncertainty principle*, Bull. Amer. Math. Soc. **9** (1983), 129-206.
4. C. Fefferman and E. M. Stein, *Some maximal inequalities*, Amer. J. Math. **93** (1971), 107-115.
5. J. Garcia-Cuerva and J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, North-Holland Mat. Stud. **116** 1985.
6. L. I. Hedberg, *On certain convolution inequalities*, Proc. Amer. Math. Soc. **36** (1972), 505-510.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI CATANIA, V. LE A. DORIA 6, 95125 CATANIA

(Current address of F. Chiarenza): DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI MESSINA
 CONTRADA PAPARDO, 98010 SANT'AGATA, MESSINA