

LOWER BOUNDS FOR THE EXTRINSIC TOTAL CURVATURES  
OF A SPACE-LIKE CODIMENSION 2 SURFACE  
IN MINKOWSKI SPACE

MAREK KOSSOWSKI

(Communicated by Jonathan M. Rosenberg)

**ABSTRACT.** There are three invariant curvature functions defined on any smooth space-like 2-surfaces in four-dimensional Minkowski space. (If the surface lies in a Euclidean hyperplane then the functions agree with  $H^2$ ,  $K^2$ , and  $(H^2 - K^2)$ . For each of these functions we show that there exists a space-like immersion of any oriented compact (or noncompact complete) surface with associated total curvature arbitrarily small.

A two dimensional oriented surface  $M$  in four dimensional Minkowski space which inherits a positive definite metric is called space-like. Such a surface carries three functions  $\mathbf{H}$ ,  $\mathbf{K}$ , and  $\mathbf{U}$  that are determined by the first and second fundamental forms. (If the surface lies in a Euclidean hyperplane then these functions reduce to  $H^2$ ,  $K^2$ , and  $(H^2 - K^2)$  respectively.) It is natural to ask for the infimum of the respective total curvatures  $\int_M |\mathbf{H}| dA$ ,  $\int_M |\mathbf{K}|^{1/2} dA$  and  $\int_M \mathbf{U}^{1/2} dA$  over the set of embedded compact space-like surfaces. In this paper we show that there exists an embedded smooth compact orientable surface  $M_g$  of genus  $g \geq 0$  with  $\int_{M_g} |\mathbf{H}| dA$  arbitrarily small (Theorem 5). In the process we relate  $\mathbf{H}$  to the split mean curvatures in the manner of reference [2]. We then show that there exist compact (and complete noncompact) embedded orientable surfaces of arbitrary genus with either  $\mathbf{K}$  or  $\mathbf{U}$  identically zero, (Theorems 6 and 7). The author would like to thank P. Ehrlich, R. Howard, T. K. Milnor and R. Penrose for inspiration.

Throughout this paper  $\mathbf{M}^4$  will denote Minkowski space, the real four dimensional vector space equipped with bilinear form  $\langle \cdot, \cdot \rangle$  of type  $(3, 1)$ , (i.e., the normal form has 3 plus signs and 1 minus sign). We will assume that  $\mathbf{M}^4$  is oriented and time oriented (i.e., a 4-volume form  $dV$ , and future time-like vector field  $\text{Fut}$ , have been chosen.) We will write  $LC = \{v \in \mathbf{M}^4 | \langle v, v \rangle = 0\}$  to denote the light cone in Minkowski space. A space-like surface in  $\mathbf{M}^4$  is

---

Received by the editors February 24, 1989 and, in revised form, September 11, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 53C50; Secondary 53B30, 53C42.

Research partially supported by NSF grant DMS 88-03585.

©1990 American Mathematical Society  
0002-9939/90 \$1.00 + \$.25 per page

a smooth immersion of a smooth compact oriented 2-manifold  $k: M \rightarrow \mathbf{M}^4$ , such that the induced metric  $k^*\langle \cdot, \cdot \rangle$  is positive definite. It follows that the normal bundle  $M^\perp \rightarrow M$  is a trivial rank two bundle with fibres carrying a type  $(1, 1)$  metric. Now choose an orthogonal splitting of Minkowski space  $\mathbf{M}^4 = \mathbf{E}^3 + \mathbf{E}^-$  with projections  $t: \mathbf{E}^3 + \mathbf{E}^- \rightarrow \mathbf{E}^-$  and  $\pi_E: \mathbf{E}^3 + \mathbf{E}^- \rightarrow \mathbf{E}^3$  and let  $U_s^i, i = F, P$  denote the two unique sections of  $M^\perp$  which satisfy the following:

- (a)  $\langle U_s^i, U_s^i \rangle = 0, \quad i = F, P,$
- (E1) (b)  $t_*(U_s^i(x))$  is of unit length for all  $x \in M, \quad i = F, P,$
- (c)  $\langle U_s^F, \text{FUT} \rangle < 0,$
- (d)  $dV(-, -, U_s^P, U_s^F) \equiv dA(-, -),$

(Here  $dA$  is the induced area form on  $M$  and equivalence is in the sense of orientations. The subscript  $s$  refers to the choice of splitting.)

We also have the splitting dependent function  $\frac{1}{\sqrt{2}}\langle U_s^F, U_s^P \rangle: M \rightarrow R$ , which we denote by  $\text{TILT}_s$ , and which satisfies:

- (a)  $0 < \text{TILT}_s \leq 1$
- (E2) (b)  $\text{TILT}_s$  is constant iff  $\text{TILT}_s = 1$  and  $M$  lies in a fiber of  $t: \mathbf{E}^3 + \mathbf{E}^- \rightarrow \mathbf{E}^-$ .

See Lemma 2 in [1].

Now for any choice of splitting  $\mathbf{M}^4 = \mathbf{E}^3 + \mathbf{E}^-$  we get two split second fundamental forms  $\Pi_s^i: TM \times TM \rightarrow R, i = F, P$  defined by  $\Pi_s^i(X, Y) = \langle \nabla_X U_s^i, Y \rangle$  where  $X, Y$  are vector fields on  $M$  and  $\nabla$  is the ambient connection. We define the *split curvatures* by  $K_s^i = \frac{\text{DET} \Pi_s^i}{\text{DET} I}, i = F, P$  and *split mean curvatures* by  $H_s^i = \frac{1}{2} \frac{\text{TR} \Pi_s^i}{\text{DET} I}, i = F, P$  where  $I$  is the first fundamental form on  $M$ .

For any normal vector  $u_x \in M_x$ , we have an endomorphism of  $T_x M$  defined by  $v_x \rightarrow$  (tangential projection of  $\nabla_v u$  at  $x$ ). There are unique global sections  $R^F, R^P$  of  $M$  that satisfy  $\langle R^F, R^P \rangle = \sqrt{2}$  and a,c, and d of (E1) above. For such a pair we have the *quadric mean curvature*  $\mathbf{H} = \frac{1}{4} \text{TR}(\nabla R^F) \cdot \text{TR}(\nabla R^P)$  and the *quartic curvature*  $\mathbf{K} = \text{DET}(\nabla R^F) \cdot \text{DET}(\nabla R^P)$ . Observe that if we fix a splitting  $\mathbf{M}^4 = \mathbf{E}^3 + \mathbf{E}^-$ , then

$$\mathbf{H} = \frac{H_s^F H_s^P}{\text{TILT}_s}, \text{ and}$$

$$\mathbf{K} = \frac{K_s^F K_s^P}{(\text{TILT}_s)^2}$$

(E3)

From these splitting dependent descriptions of  $\mathbf{H}$  and  $\mathbf{K}$  it is apparent that

$$\mathbf{U} = \frac{(K_s^F - (H_s^F)^2)(K_s^P - (H_s^P)^2)}{(\text{TILT}_s)^2} \geq 0$$

(E4)

is also an invariant, which we will call the *umbilic curvature*. All three functions  $\mathbf{H}, \mathbf{K}, \mathbf{U}: M \rightarrow \mathbf{R}$  are congruence invariants for  $(M, k)$ .

**Proposition 1.** *Given  $(M, k)$ , then for all splittings  $\mathbf{M}^4 = \mathbf{E}^3 + \mathbf{E}^-$ ,*

$$\int_M |\mathbf{H}| dA \leq \left[ \sup \frac{1}{\text{TILT}_s} \right] \int_M (H_s^F)^2 + (H_s^P)^2 dA.$$

*Further if equality holds for some splitting then  $(M, k)$  is an immersion into a space-like hyperplane  $\mathbf{E}^3 \subset \mathbf{M}^4$ . (The supremum is taken over all splittings  $\mathbf{M}^4 = \mathbf{E}^3 + \mathbf{E}^-$ ).*

*Proof.* Choose a splitting. Then (E3) implies that

$$2|\mathbf{H}| \leq \frac{1}{\text{TILT}_s} \left[ (H_s^F)^2 + (H_s^P)^2 \right],$$

from which follows the inequality

$$2 \int_M |\mathbf{H}| dA \leq \int_M \frac{(H_s^F)^2 + (H_s^P)^2}{\text{TILT}_s} dA \leq \left[ \sup \frac{1}{\text{TILT}_s} \right] \int_M (H_s^F)^2 + (H_s^P)^2 dA.$$

Since equality implies  $(\text{TILT}_s)$  is constant (E2b) implies our claim.

**Theorem 2.** *Given  $(M, k)$  then for all splittings  $\mathbf{M}^4 = \mathbf{E}^3 + \mathbf{E}^-$ ,*

$$8\pi \leq \left[ \sup \frac{1}{\text{TILT}_s} \right] \int_M (H_s^F)^2 + (H_s^P)^2 dA.$$

*Further if equality holds for some splitting then  $M = S^2$  and  $k$  embeds  $S^2$  as a round sphere in a space-like hyperplane  $\mathbf{E}^3 \subset \mathbf{M}^4$ .*

*Proof.* Recall that for any splitting we have:

$$K_s^i \leq (H_s^i)^2 \quad i = F, P,$$

with equality at  $x \in M$  if and only if  $x$  is an umbilic point for  $\Pi_s^i, i = F, P$ . Now with (E7) of [1] we have that

$$\begin{aligned} 8\pi &\leq \int_{\{K_s^F > 0\}} K_s^F dA + \int_{\{K_s^P > 0\}} K_s^P dA \\ &\leq \int_{\{K_s^F > 0\}} (H_s^F)^2 dA + \int_{\{K_s^P > 0\}} (H_s^P)^2 dA \\ (E5) \quad &\leq \int_M (H_s^F)^2 + (H_s^P)^2 dA \\ &\leq \left[ \sup \frac{1}{\text{TILT}_s} \right] \int_M (H_s^F)^2 + (H_s^P)^2 dA \end{aligned}$$

Finally if we have equality, then  $\text{TILT}_s = 1$ , hence  $(M, k)$  is an immersion into an  $\mathbf{E}^3$  factor of the splitting with  $\mathbf{E}^3$ -mean curvature given by  $H_s^F = H_s^P$  and Gaussian curvature given by  $K_s^F = K_s^P$ . Thus  $(M, k)$  is a totally umbilic surface in a  $\mathbf{E}^3 \subset \mathbf{M}^4$ .

Next we examine how the split mean curvatures behave individually.

**Theorem 3.**

(a) Given  $(M, k)$  with  $M = S^2$ , then for all splittings  $\mathbf{M}^4 = \mathbf{E}^3 + \mathbf{E}^-$

$$4\pi \leq \int_{S^2} (H_s^i)^2 dA \quad i = F, P$$

Further equality for  $i = F$  or  $P$  implies that  $k$  embeds  $S^2$  in a light cone  $LC \subset \mathbf{M}^4$ . Equality in both  $i = F$  and  $P$  implies that  $(M, k)$  is congruent to a round  $S^2$  in a space-like hypersurface  $\mathbf{E}^3 \subset \mathbf{M}^4$ .

(b) Given  $\varepsilon > 0$  and an integer  $g > 0$ , then there exists an orientable compact space-like embedded surface of genus  $g$  and a splitting  $\mathbf{M}^5 = \mathbf{E}^3 + \mathbf{E}^-$  such that for  $i = F$  or  $P$

$$0 \leq \int_{M_g} (H_s^i)^2 dA \leq \varepsilon.$$

(Note that because of (E5) above this holds for only one of the split mean curvatures.)

*Proof.* (a) Referring to Theorem 1 of [1] we have that

$$4\pi \leq \int_{\{K_s^i > 0\}} K_s^i dA, \quad i = F, P$$

where the integration is over the subset of  $M$  where  $K_s^i > 0$ . As a consequence we have that for  $i = F, P$ ,

$$\begin{aligned} 4\pi &\leq \int_{\{K_s^i > 0\}} K_s^i dA \\ &\leq \int_{\{K_s^i > 0\}} (H_s^i)^2 dA \leq \int_{S^2} (H_s^i)^2 dA. \end{aligned}$$

Now suppose there is equality for  $i = F$  or  $P$ . From (E6) of [2] it follows that as conformal structures on  $M$ ,  $\Pi_s^i$  for  $(M, k)$  agrees with  $\bar{\Pi}$  for the light-like hypersurface  $LL^i(M, k)$ . Thus if we choose any (affine) space-like hyperplane  $\mathbf{E}^3 \subset \mathbf{M}^4$ , at any immersed point in  $\{\mathbf{E}^3 \cap LL^i(M, k)\}$  this intersection is locally an umbilic surface in  $\mathbf{E}^3$ , and hence is a piece of a round sphere or of a plane. Since we are assuming  $M$  is compact, we will have our result if we show that there exists an  $\mathbf{E}^3 \subset \mathbf{M}^4$  with intersection  $\{\mathbf{E}^3 \cap LL^i(M, k)\}$  consisting of only immersive points (i.e.,  $LL^i(M, k)$  is an icon). To prove this choose a hyperplane  $\tilde{\mathbf{E}}^3 \subset \mathbf{M}^4$  so that the intersection  $\{\tilde{\mathbf{E}}^3 \cap LL^i(M, k)\}$  contains an immersive point. Recall that  $LL^i(M, k) \xrightarrow{\pi^i} M$  is fibered by null lines and each fiber intersects the chosen  $\tilde{\mathbf{E}}^3$  in a unique point. (See [1] for details.) Since  $k: M \rightarrow \mathbf{M}^4$  has image, which is a section of this fibering of  $LL^i(M, k)$ ,

we now have smooth maps

$$S^2 = M \xrightarrow{k} \mathbb{L}^i(M, k) \subset \mathbb{M}^4$$

$$\downarrow p$$

$$\tilde{\mathbb{E}}^3 \subset \mathbb{M}^4$$

where  $p \circ k(x)$  is intersection point of the  $\pi^i$ -fibered through  $k(x)$  and the chosen  $\tilde{\mathbb{E}}^3$ . We know the immersive points of  $p \circ k$  are open in  $S^2 = M$ , we need only show that they are closed. Let  $x_n \rightarrow x$  be a convergent sequence of points in  $S^2$  with  $p \circ k$  immersive at  $x_n$ . Now consider the  $\mathbb{E}^3$  parallel to the above chosen  $\tilde{\mathbb{E}}^3$  that contains  $k(x)$ . We know that  $\{\mathbb{E}^3 \cap \mathbb{L}^i(M, k)\} = V$  is a piece of a sphere or a plane. Hence  $(\pi^i)^{-1}(\pi^i(V))$  is a subset of a light cone or a null 3-plane. But for any affine  $\mathbb{E}^3 \subset \mathbb{M}^4$ , the intersection of  $\mathbb{E}^3$  with a light cone or a null 3-plane consists entirely of immersive points or entirely of nonimmersive points. Thus the intersection of  $(\pi^i)^{-1}(\pi^i(V)) \subset \mathbb{L}^i(M, k)$  with the chosen  $\tilde{\mathbb{E}}^3$  contains  $p \circ k(x_n)$  (for  $n$  sufficiently large). These are immersive points for  $p \circ k$  and hence immersive points for the intersection with  $\tilde{\mathbb{E}}^3$ . Since  $p \circ k(x)$  lies in this intersection it is also an immersive point. We conclude  $\mathbb{L}^i(M, k)$  is congruent to a light cone. For the last claim in Theorem 3 (a) we now know that both  $\mathbb{L}^i(M, k)$  are congruent to light cones. Hence  $(M, k)$  is space-like, compact, and lies in their intersection. It must be a round  $S^2 \subset \mathbb{E}^3 \subset \mathbb{M}^4$ .

(b) The construction of  $(M_g, k)$  is but a slight modification of the construction presented in Appendix 2 of [1]. Consider a fixed splitting  $\mathbb{M}^4 = \mathbb{E}^3 + \mathbb{E}^-$  with projection  $t: \mathbb{E}^3 + \mathbb{E}^- \rightarrow \mathbb{E}^-$  and the torus of revolution  $(T^2, k)$  obtained by rotating an “embedded Figure 8” about a  $\mathbb{M}^2$  in  $\mathbb{M}^4$  so tha a linear segment in the figure 8 generates a cylinder in  $\mathbb{E}^3$ ; and the  $S^2$  valued Gauss maps are not surjective. For this torus and splitting  $H_s^i = \frac{1}{2} \frac{\text{TR}\langle \nabla U_s^i, - \rangle}{\text{DETI}}$ ,  $i = F, P$ . Now because the Gauss maps are not surjective, there exist isometries  $b_\theta: \mathbb{M}^4 \rightarrow \mathbb{M}^4$  such that for all positive integers  $n > 0$ , there exist  $\theta_n$  so that  $n < \text{INF}|t_*(b_{\theta_n} \circ U_s^F)|$  where the infimum is over points of the torus. Since  $b_{\theta_n}$  is an isometry we have that  $(T^2, b_{\theta_n} \circ k)$  has  $U_{s\theta_n}^F = \lambda(b_n \circ U_s^F)$  where  $\lambda > \frac{1}{n}$  is a function on  $T^2$  and  $U_{s\theta_n}^F$  is the vector field satisfying (E1) above relative to the fixed splitting. We have that,

$$H_{s\theta_n}^F = \frac{\text{TR}\langle \nabla U_{s\theta_n}^F, - \rangle}{\text{DETI}} < \frac{1}{n} \frac{\text{TR}\langle \nabla U_s^F, - \rangle}{\text{DETI}} < \frac{H_s^F}{n}$$

and hence a torus with  $\int_{T^2} (H_s^F)^2 dA$  arbitrarily small. Note that the key step of constructing  $b_{\theta_n}$  requires only that the Gauss map be nonsurjective. Thus to construct higher genus examples take two copies of  $(T^2, k)$  and shrink one

copy so that the two copies do not intersect, and the cylinders in  $E^3$  mentioned at the outset are concentric. We can connect these two cylinders in the  $E^3$  with catenoidal necks in such a way that removing the  $E^3$ -minimal parts of the necks results in two punctured tori with  $M^4$ -Gauss maps nonsurjective. If we denote this connected sum by  $(M_g, k)$  then for any splitting  $M^4 = E^3 + E^-$ , both the  $H_s^i$ ,  $i = F, P$  will vanish on the  $E^3$ -minimal part of the necks. Thus we may repeat the above argument on  $(M_g - \{H_s^F = 0\}, b_{\theta_n} \circ k)$  and we have surfaces which satisfy our claim for  $i = F$ . The same argument adapts to construct an example for  $i = P$ .

By a *space-like graph* in  $M^4$  we mean a smooth space-like hypersurface which, relative to some (hence any) splitting of  $M^4 = E^3 + E^-$  is globally the graph of a function  $f: E^3 \rightarrow E^-$ .

**Theorem 4.** *If  $(M, k)$  is an embedding into a space-like graph, then for all splittings of  $M^4 = E^3 + E^-$*

$$4\pi \leq \int_M (H_s^i)^2 dA, \quad i = F, P.$$

Further equality for  $i = F$  or  $P$  implies that  $M = S^2$  so that Theorem 3, Part (a) applies.

*Proof.* Recall from the proof of Theorem 9 in [1] that for  $(M, k)$  a surface embedded in a graph we must have

$$4\pi \leq \int_{\{K_s^i > 0\}} K_s^i dA, \quad i = F, P.$$

(This is a restatement of the first line in the proof of Theorem 9.) Now for the case of equality we must have by Theorem 1 of [1] that  $M = S^2$ .

**Theorem 5.** *Given  $\varepsilon > 0$  and an integer  $g \geq 0$ , there exists an orientable compact embedded space-like surface of genus  $g$  with,*

$$0 < \int_{M_g} |\mathbf{H}| dA < \varepsilon.$$

*Proof.* We first construct an example with  $g = 0$ . Take two round spheres  $S_1^2$  and  $S_2^2$  in  $E^3 \subset M^4$  and smoothly connect them with a ‘‘Hopf neck’’  $HN \subset E^3$  of small total squared  $E^3$ -mean curvature to yield an embedded sphere,  $S^2 \# S^2$ . This can be done as in Figure 1 so that the transition region between round sphere and Hopf neck consist of a pair of planar parallel annuli in  $E^3$ . (For details concerning the Hopf neck see Lemma 6b in [3].) Let  $P_1$  and  $P_2$  be the parallel planes in which these annuli lie. Now extend the  $S^2 \# S^2 \subset E^3$  to a Light-like hypersurface  $LL(S^2 \# S^2)$  in  $M^4$  so that the vertex of round spherical parts lie in the ‘‘future’’ of the  $E^3$  fixed at the outset. Similarly extend the round spheres  $S_1^2$  and  $S_2^2$  to light cones  $LC_1$  and  $LC_2$  so that their vertices are in the future of the  $E^3$ . Next extend the 2-planes  $P_1$  and  $P_2$  to null 3-planes  $N_1^3$




FIGURE 1

and  $N_2^3$  so that  $N_\alpha^3$  intersects the future of  $LC_\alpha$  in a bounded set. Let  $\bar{N}_\alpha^{3F}$ ,  $\alpha = 1, 2$  denote the closure of the subsets of  $N_\alpha^3$  that lie in the future of  $E^3$  and let  $D_\alpha = LL(S^2 \# S^2) \cap \bar{N}_\alpha^{3F}$ ,  $\alpha = 1, 2$ . These  $D_\alpha$  are embedded space-like 2-disks. Thus by construction  $\{D_1 \cup HN \cup D_2\}$  is an embedded space-like two-sphere that intersects the fixed  $E^3$  in the Hopf neck along with its pair of planar annuli. Now since each  $D_\alpha$  lies in a null 3-plane  $H$  vanishes on both  $D_\alpha$ . Thus the support of  $H$  lies entirely in the Hopf neck where it agrees with the  $E^3$ -mean curvature squared. We have our sphere. To construct higher genus examples we need only connect the parallel annuli with additional Hopf necks with small total squared  $E^3$ -mean curvature and we are finished.

We now construct space-like surfaces with  $K$  and  $U$  identically zero. We remark that the existence of  $K$ -flat surfaces is relevant to the conjecture discussed in Appendix 3 of [1] in that these surfaces indicate that there need not exist a point  $x \in M$  where both split curvatures are positive  $K_s^i(x) > 0 \quad i = F, P$ .

**Theorem 6.** *Given  $g \geq 0$ , there exist orientable embedded compact (or noncompact complete) space-like surfaces of genus  $g$  with  $K = 0$ .*

*Proof.* Consider any two  $E^3$ -flat surfaces  $\bar{k}^i: FLT^i \rightarrow E^3 \subset M^4$ ,  $i = F, P$  that intersect transversely on an immersed circle  $\tilde{S}^1 \rightarrow FLT^F \cap FLT^P$ . (The reason for indexing these surfaces by  $F$  and  $P$  will be apparent below.) Next choose  $E^3$ -normals for each surface  $n^i: FLT^i \rightarrow S^2 \subset E^3$ ,  $i = F, P$ . As in the proof of Proposition 6 of [2] we use these  $E^3$ -normals to construct two light-like hypersurfaces  $j^i: LL^i(FLT^i) \rightarrow M^4$ ,  $i = F, P$ . Now two null 3-planes in  $M^4$  intersect in an  $E^2 \subset M^4$  or coincide. Since  $FLT^i$  intersect transversely in  $E^3$  we have that  $LL^F(FLT^F)$  intersects  $LL^P(FLT^P)$  transversely in  $M^4$ . Thus near

the intersecting circle we have that  $LL^F(FLT^F) \cap LL^P(FLT^P)$  is an immersed space-like cylinder  $\tilde{S}^1 \times (-\delta, \delta)$ . (It will be embedded if the  $FLT^i \subset E^3$ ,  $i = F, P$  are embedded.) It follows that this small cylinder is transverse to the fibers of both  $\pi^i: LL^i(FLT^i) \rightarrow FLT^i$ . We may view the original  $FLT^i$   $i = F, P$  as “zero sections” of  $\pi^i$  and observe that the intersection cylinder is transverse to both of these zero sections in  $LL^i(FLT^i)$ . Thus we may extend the  $\tilde{S}^1 \times (-\delta, \delta)$  so that one boundary edge  $\tilde{S}^1 \times \{-\delta\}$  extends so as to agree with the zero section of  $\pi^F$  outside of a compact set and the other boundary edge  $\tilde{S}^1 \times \{\delta\}$  extends so as to agree with the zero section of  $\pi^P$  outside a compact set. We may construct this extension so that it is everywhere transverse to (at least) one of the fibrations  $\pi^i$ ,  $i = F, P$ , thence this *extended intersection*  $(FLT^F \# FLT^P, k)$  is a space-like surface in  $M^4$ .

In Figure 3 we illustrate our construction and include the image of the extended intersection under the projection  $\pi_E: M^4 \rightarrow E^3$ . Now (E10) of [2] tells us that any section of either  $LL^i(FLT^i)$   $i = F, P$  must have  $K$  identically zero. Using the construction on flat surfaces as represented by Figures 2 and 3 we can build spheres or tori and take connected sums so that the resulting surfaces have  $K$  identically zero. We have our surfaces.

**Theorem 7.** *Given  $g \geq 0$  there exists an orientable embedded compact (or non-compact complete) space-like surface of genus  $g$  with  $U = 0$ .*

*Proof.* The proof parallels that of Theorem 6 with the modification that flat surfaces in  $E^3$  are replaced by round spheres  $S_i^2$ ,  $i = F, P$  in  $E^3$ . The key point is again (E10) of [2], which in this context tells us that any section of either  $LL^i(S_i^2)$   $i = F, P$  must have  $U$  identically zero.

It should be apparent to the reader that these *subumbilic* immersions are highly nonrigid. So that  $F(M, k) = \int_M U^{1/2} dA$  defines a geometric functional




FIGURE 2






FIGURE 3

on  $\text{Immer}^\infty(M, M^4)$  with highly degenerate second variation at the minima constructed above.

## BIBLIOGRAPHY

1. J. K. Beem and P. E. Ehrlich, *Global Lorentzian, Geometry*, Dekker, 1981.
2. M. Kossowski, *The  $S^2$  valued Gauss maps and split total curvature of a space-like codimension 2 surface in Minkowski space*, J. London Math. Soc. **40**.
3. —, *The intrinsic conformal structure and Gauss map of a Light-like hypersurface in Minkowski space*, Trans. Amer. Math. Soc. **316** (1989).
4. W. Kuhnel and U. Pinkall, *On total mean curvatures*, Quart. J. Math. Oxford **37** (1986).
5. T. K. Milnor, *Harmonic maps and classical surface theory in Minkowski 3-space*, Trans. Amer. Math. Soc. **280** (1983).
6. B. O'Neil, *Semi-Riemannian geometry with applications to relativity*, Academic Press, New York, 1983.
7. R. Penrose, *Battille Recontres*, Benjamin, NY, 1968.
8. T. J. Wilmore, *Total curvature in Riemannian geometry*, Wiley, 1982.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SOUTH CAROLINA 29208