

## NONEXISTENCE OF ALMOST COMPLEX STRUCTURES ON GRASSMANN MANIFOLDS

PARAMESWARAN SANKARAN

(Communicated by Frederick R. Cohen)

**ABSTRACT.** In this paper we prove that, for  $3 \leq k \leq n-3$ , none of the oriented Grassmann manifolds,  $\tilde{G}_{n,k}$ —except for  $\tilde{G}_{6,3}$ , and a few as yet undecided cases—admits a weakly almost complex structure. The result for  $k = 1, 2, n-1, n-2$  are well known and classical. The proofs make use of basic concepts in  $K$ -theory, the property that  $\tilde{G}_{n,k}$  is  $(n-k)$ -universal, known facts about  $K(\mathbb{H}P^4)$ , and characteristic classes.

### 1. INTRODUCTION

For  $1 \leq k < n$ , let  $\tilde{G}_{n,k}$  denote the oriented Grassmann manifold of oriented  $k$ -dimensional vector subspaces of  $\mathbb{R}^n$ .  $\tilde{G}_{n,k}$  is a smooth manifold of dimension  $k(n-k)$ . Note that  $\tilde{G}_{n,1} \cong S^{n-1}$ , the  $(n-1)$ -sphere, and that  $\tilde{G}_{n,k} \cong \tilde{G}_{n,n-k}$  under the diffeomorphism that sends an oriented  $k$ -plane  $V$  to  $V^\perp$  together with that orientation on  $V^\perp$  which induces the standard orientation on  $V \oplus V^\perp = \mathbb{R}^n$ .

Recall that a smooth manifold  $M$  is said to be (weakly) almost complex if its tangent bundle  $\tau M$  is (stably) isomorphic to a complex vector bundle over  $M$ . For example,  $\tilde{G}_{n,1} \cong S^{n-1}$  is weakly almost complex for all  $n$ , but is almost complex only when  $n = 3$  or  $7$ . (See [14, p. 217] and [5, 15.1].) It is a classical result that  $\tilde{G}_{n,2} \cong SO(n)/(SO(2) \times SO(n-2))$  is a Hermitian symmetric space, and is therefore almost complex for all  $n$ .

In this paper, we investigate which of the remaining  $\tilde{G}_{n,k}$ 's are weakly almost complex. Since  $\tilde{G}_{n,k} \cong \tilde{G}_{n,n-k}$ , we assume, without loss of generality, that  $2k \leq n$ . Our main result is

**1.1. Theorem.** *Let  $3 \leq k \leq n/2$ . Then*

- (i)  $\tilde{G}_{n,k}$  is not weakly almost complex if  $n$  is odd or if  $(n-k) \geq 16$ .
- (ii)  $\tilde{G}_{6,3}$  is weakly almost complex.  $\tilde{G}_{6,3} \times \tilde{G}_{6,3}$  is almost complex.

---

Received by the editors January 8, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 57R15.

This work was done when the author was at the Institute of Mathematical Sciences, C.I.T. Campus, Madras.

Attempts by the author to settle the remaining cases for which  $n$  is even and  $3 \leq k \leq (n - k) \leq 15$ , left unanswered in the above theorem, have failed.

Let  $\gamma_{n,k}$  denote the canonical  $k$ -plane bundle over  $\tilde{G}_{n,k}$ , and let  $\beta_{n,k}$  be its orthogonal complement, whose fiber over a  $V \in \tilde{G}_{n,k}$  is the vector space  $V^\perp \subset \mathbb{R}^n$ . One has the bundle equivalence

$$(1.2) \quad \gamma_{n,k} \oplus \beta_{n,k} \approx n\varepsilon,$$

where  $\varepsilon$  denotes a trivial line bundle.

It is well known that the tangent bundle  $\tau\tilde{G}_{n,k}$  of  $\tilde{G}_{n,k}$  has the following description (see [9]):

$$(1.3) \quad \tau\tilde{G}_{n,k} \approx \gamma_{n,k} \otimes_{\mathbb{R}} \beta_{n,k}.$$

Using (1.3) or [6], we obtain

$$(1.4) \quad \tau\tilde{G}_{n,k} \oplus (\gamma_{n,k} \otimes_{\mathbb{R}} \gamma_{n,k}) \approx n\gamma_{n,k}.$$

For a topological space  $X$ , let  $r : K(X) \rightarrow KO(X)$  denote the homomorphism (of Abelian groups) gotten by restriction of scalars to  $\mathbb{R}$ , and let  $c : KO(X) \rightarrow K(X)$  denote the complexification,  $c[\xi] = [\xi \otimes_{\mathbb{R}} \mathbb{C}]$ , which is a ring homomorphism.

One has the following identities:

$$(1.5) \quad r \circ c(x) = 2x \quad \forall x \in KO(X)$$

$$(1.6) \quad c \circ r(y) = y + y' \quad \forall y \in K(X),$$

where  $y'$  stands for complex conjugation of  $y$ .

Note that a smooth manifold  $M$  is weakly almost complex if and only if  $[\tau M] + \delta$  is in the image of  $r$ , where  $\delta = 0$  or  $[\varepsilon]$  according to whether  $\dim M$  is even or odd.

## 2. K-THEORY OF QUATERNIONIC PROJECTIVE SPACES

Let  $\mathbb{H}P^n$  denote the quaternionic projective  $n$ -space. Let  $\eta_n$  denote the canonical right  $\mathbb{H}$ -vector bundle over  $\mathbb{H}P^n$ , and let  $\xi_n$  denote the complex vector bundle obtained from  $\eta_n$  by restriction of scalars to  $\mathbb{C}$ . Let  $v = -c_2(\xi_n) \in H^4(\mathbb{H}P^n; \mathbb{Z}) \subset H^4(\mathbb{H}P^n; \mathbb{Q})$ . Then  $v$  generates the ring  $H^*(\mathbb{H}P^n; \mathbb{Q})$ . The Chern character  $\text{ch}(\xi_n)$  of  $\xi_n$  is

$$\text{ch}(\xi_n) = \exp(y_1) + \exp(y_2),$$

where  $(1 + y_1)(1 + y_2) = 1 - v = c(\xi_n)$ , the total Chern class of  $\xi_n$ . Hence  $y_1 + y_2 = 0$ , and  $y_1 y_2 = -v$ . Therefore  $y_1 = -y_2$ , and  $y_1^2 = +v$ . Thus

$$(2.1) \quad \text{ch}(\xi_n) = \exp(\sqrt{v}) + \exp(-\sqrt{v}).$$

2.2. **Proposition.** *The Chern character  $ch : K(\mathbb{H}P^n) \rightarrow H^*(\mathbb{H}P^n; \mathbb{Q})$  is a monomorphism. The image of the Chern character is freely generated over the integers by  $\{1, w, w^2, \dots, w^n\}$ , where*

$$w = 2 \cosh(\sqrt{v}) - 2 = v + \frac{2v^2}{4!} + \dots + \frac{2v^n}{(2n!)}.$$

The proof can be found in [11]. (See also [2, 3.1].)

Note that  $w^{n+1} = 0$ .

Let  $\eta_n^* = \text{Hom}_{\mathbb{H}}(\eta_n, \mathbb{H})$  be the dual of  $\eta_n$ , which is a left  $\mathbb{H}$ -vector bundle of rank 1. Consider the bundle  $\omega = \eta_n \otimes_{\mathbb{H}} \eta_n^*$  over  $\mathbb{H}P^n$ . Then  $\omega$  is a real vector bundle whose rank is 4. The map  $\langle q \rangle \mapsto q \otimes f_q$ , where  $f_q : q\mathbb{H} = \langle q \rangle \rightarrow \mathbb{H}$  is the  $\mathbb{H}$ -linear map defined by  $f_q(q) = 1$ , is a well-defined, continuous, nowhere vanishing section of the bundle  $\omega$ . Hence  $\omega$  splits as  $\omega \approx \varepsilon \oplus \zeta$ , where  $\zeta$  is a 3-plane bundle.  $\zeta$  is necessarily orientable, since  $\mathbb{H}P^n$  is simply connected.

2.3. **Lemma.** *Let  $n \geq 4$ . Then  $[\zeta \otimes_{\mathbb{R}} \zeta] + [\varepsilon] \in KO(\mathbb{H}P^n)$  is not in the image of  $r : K(\mathbb{H}P^n) \rightarrow KO(\mathbb{H}P^n)$ .*

*Proof.* Note that the rank of  $\zeta \otimes_{\mathbb{R}} \zeta$  is 9. Also,  $[\omega \otimes_{\mathbb{R}} \omega] = [\zeta \otimes_{\mathbb{R}} \zeta] + [\varepsilon] + 2[\zeta]$ , and  $2[\zeta] \in \text{Im}(r)$  by (1.5). Thus it suffices to show that  $[\omega]^2 \notin \text{Im}(r)$ .

Let, if possible,  $[\omega]^2 = r(y)$ , for some  $y \in K(\mathbb{H}P^n)$ . Then, by (1.6),

$$c([\omega]^2) = c \circ r(y) = y + y'.$$

That is,

$$(2.4) \quad c([\omega]^2) = y + y'.$$

Now  $c[\omega] = c[\eta_n \otimes_{\mathbb{H}} \eta_n^*] = [\xi_n \otimes_{\mathbb{C}} \xi_n^*]$ , where  $\xi_n^*$  is the complex bundle obtained from  $\eta_n^*$  by restricting the scalars to  $\mathbb{C}$ . The last equality follows from the fact that there exists a functorial isomorphism between  $(V \otimes_{\mathbb{H}} W) \otimes_{\mathbb{R}} \mathbb{C}$  and  $V \otimes_{\mathbb{C}} W$ , where  $V$  is a right  $\mathbb{H}$ -vector space,  $W$  is a left  $\mathbb{H}$ -vector space and they are regarded as  $\mathbb{C}$ -vector spaces by restriction of scalars. (See [1, 3.7–3.9].)

Taking Chern characters on both sides of (2.4), we get

$$\text{ch}(y) + \text{ch}(y') = \text{ch}(c[\omega]^2) = (\text{ch}(c[\omega]))^2 = \text{ch}([\xi_n])^2 \text{ch}([\xi_n^*])^2.$$

Since  $H^j(\mathbb{H}P^n; \mathbb{Q}) = 0$  unless  $j \equiv 0 \pmod{4}$ , the total Chern class  $c(y)$  equals  $c(y')$ , and  $c(\xi_n) = c(\xi_n^*)$ . Therefore we get

$$\begin{aligned} \text{ch}(y) &= 1/2(\text{ch}([\xi_n])^4) = 1/2(2 \cosh(\sqrt{v}))^4 \\ &= 1/2w^4 + \text{terms involving lower powers of } w. \end{aligned}$$

This is a contradiction because, by Proposition 2.2,  $\text{ch}(y)$  must be an integral linear combination of powers of  $w$  and  $w^4 \neq 0$  for  $n \geq 4$ . This proves the lemma.

3. PROOF OF THEOREM 1.1

Let  $3 \leq k \leq n/2$ .

The manifold  $\tilde{G}_{6,3}$  is parallelizable [13]. Therefore, it is weakly almost complex. (Note that  $\dim \tilde{G}_{6,3}$  is odd.) It also follows that  $\tilde{G}_{6,3} \times \tilde{G}_{6,3}$  is almost complex.

Let  $n$  be odd. As  $3 \leq k \leq n/2$ , we have  $n \geq 7$ . Let  $\pi : \tilde{G}_{n,k} \rightarrow G_{n,k}$  be the double covering map onto the Grassmann manifold  $G_{n,k}$  of  $k$ -planes in  $\mathbb{R}^n$ . It can be seen from [3, Theorem 1.1] that the Stiefel-Whitney class  $w_3(\tilde{G}_{n,k}) = \pi^*(w_3(G_{n,k})) = w_3(\gamma_{n,k})$ , for  $n$  odd and  $k \geq 3$ . Using the Gysin sequence of the double covering map  $\pi$ , and the knowledge of  $H^*(G_{n,k}; \mathbb{Z}_2)$ , it can be shown that  $w_3(\gamma_{n,k}) \neq 0$  for  $n \geq 7$ . Since the odd-dimensional Stiefel-Whitney classes of any weakly almost complex manifold must vanish [12], it follows that  $\tilde{G}_{n,k}$  is not weakly almost complex.

Now let  $n$  be even,  $k \geq 3$ , and  $n - k \geq 16 = \dim \mathbb{H}P^4$ . Since  $\tilde{G}_{n,k}$  is  $(n - k)$ -universal for orientable  $k$ -plane bundles, there exists a map  $f : \mathbb{H}P^4 \rightarrow \tilde{G}_{n,k}$  such that  $f^*(\gamma_{n,k}) = \zeta \oplus m\varepsilon$ , where  $m = k - 3$ , and  $\zeta$  is the orientable 3-plane bundle of §2. One has

$$f^*(\gamma_{n,k} \otimes_{\mathbb{R}} \gamma_{n,k}) \approx (\zeta \otimes_{\mathbb{R}} \zeta) \oplus 2m\zeta \oplus m^2\varepsilon.$$

Using (1.4), (1.5), Lemma 2.3, and the fact that  $n$  is even, we see that  $\tilde{G}_{n,k}$  is not weakly almost complex. This completes the proof of Theorem 1.1.

We now turn to the Grassmann manifolds. Denote the unique (up to bundle equivalence) nontrivial line bundle over  $G_{n,k}$  by  $\psi$ . Note that the tangent bundle of the real projective space  $\mathbb{R}P^{n-1} \cong G_{n,1}$  is stably isomorphic to the bundle  $n\psi$ . Hence  $G_{n,1}$  is weakly almost complex for  $n$  even. For  $n$  odd  $G_{n,1}$  is not orientable and hence it is not weakly almost complex.

3.1. **Lemma.** *Let  $2 \leq k \leq n/2$ .*

- (i) (Borel-Hirzebruch [4, p. 526]) *For  $n \geq 5$ ,  $G_{n,2}$  is not almost complex.*
- (ii)  *$G_{n,k}$  is not weakly almost complex if  $n$  is odd, or if  $k \geq 3$ , and  $n - k \geq 16$ .*
- (iii)  *$G_{4,2}$  and  $G_{6,3}$  are weakly almost complex.  $G_{4,2}$  is not almost complex.*

*Proof of (ii).* For  $n$  odd,  $G_{n,k}$  is not orientable, and is therefore not weakly almost complex. The remaining cases now follow from Theorem 1.1, the naturality of the homomorphism  $r$ , and the observation that  $\tau\tilde{G}_{n,k} \approx \pi^*(\tau G_{n,k})$ , where  $\pi : \tilde{G}_{n,k} \rightarrow G_{n,k}$  is the double covering map.

*Proof of (iii).* Denote by  $\gamma$  and  $\beta$  the canonical  $k$ - and  $(n - k)$ -plane bundles over  $G_{n,k}$ . According to Lam [9], a stable normal bundle to  $G_{n,k}$  is  $\lambda^2(\gamma) \oplus \lambda^2(\beta)$ . When  $n = 4$  and  $k = 2$ ,  $\lambda^2(\gamma) \approx \lambda^2(\beta) \approx \psi$ . Hence the stable normal bundle in this case is in the image of  $r$ . It follows that  $G_{4,2}$  is weakly almost complex.

On  $G_{6,3}$ ,  $\lambda^2(\gamma) \approx \gamma \otimes_{\mathbb{R}} \psi$ , and  $\lambda^2(\beta) \approx \beta \otimes_{\mathbb{R}} \psi$  as can be shown using [7, Proposition 10.3, Chapter 12] the Hodge duality  $\lambda^{n-1}(\nu) \approx \nu$  for orientable  $n$ -plane bundles, and  $\theta \otimes_{\mathbb{R}} \theta \approx \varepsilon$  for any line bundle  $\theta$ . Thus, on  $G_{6,3}$ ,

$$\lambda^2(\gamma) \oplus \lambda^2(\beta) \approx (\gamma \otimes_{\mathbb{R}} \psi) \oplus (\beta \otimes_{\mathbb{R}} \psi) \approx (\gamma \oplus \beta) \otimes_{\mathbb{R}} \psi \approx 6\varepsilon \otimes_{\mathbb{R}} \psi \approx 6\psi,$$

which is in the image of  $r$ . It follows that  $G_{6,3}$  is weakly almost complex.

To see that  $G_{4,2}$  is not almost complex, notice, first, that  $H^i(G_{4,2}; \mathbb{Q}) \cong \mathbb{Q}$  for  $i = 0, 4$ , and is zero for  $i \neq 0, 4$ . Since  $2\psi$  is a stable normal bundle for  $G_{4,2}$ , we have the following formula for the rational Pontrjagin class  $p_1(G_{4,2}) \in H^4(G_{4,2}; \mathbb{Q})$ :

$$(3.2) \quad p_1(G_{4,2}) = -p_1(2\psi) = c_2(2\psi \otimes_{\mathbb{R}} \mathbb{C}) = (c_1(\psi \otimes_{\mathbb{R}} \mathbb{C}))^2 = 0$$

because  $H^2(G_{4,2}; \mathbb{Q}) = 0$ . On the other hand, if  $\tau = \tau_{G_{4,2}}$  were a complex vector bundle, then [12],

$$-p_1(G_{4,2}) = 2c_2(\tau) - (c_1(\tau))^2 = 2c_2(\tau).$$

Since the top Chern class of  $\tau$  is its Euler class, we must have  $-\frac{1}{2}\langle p_1(G_{4,2}), [G_{4,2}] \rangle = \langle c_2(\tau), [G_{4,2}] \rangle = \chi(G_{4,2}) = 2$ , the Euler characteristic of  $G_{4,2}$ . Hence  $p_1(G_{4,2}) \neq 0$ , contradicting (3.2). This shows that  $G_{4,2}$  is not almost complex.

As a corollary to the above theorem and Theorem 1.1., we obtain the following:

**3.3. Theorem.**

- (i) *A product of any finite number of oriented Grassmann manifolds,  $\tilde{G}_{n,k}$ 's, (resp. the Grassmann manifolds  $G_{m,p}$ 's) is not weakly almost complex if, for one of the factors,  $n \geq 7$  is odd,  $3 \leq k \leq n - 3$ ; or  $k \geq 3$ ,  $(n - k) \geq 16$ ; (resp.  $m$  is odd; or  $p \geq 3$ ,  $(m - p) \geq 16$ ).*

- (ii) *The oriented flag manifold*

$$\tilde{G}(n_1, n_2, \dots, n_s) = SO(n)/(SO(n_1) \times SO(n_2) \times \dots \times SO(n_s)),$$

where  $n = n_1 + \dots + n_s$  is not weakly almost complex if, for some  $i \neq j$ , any one of the following holds: (a)  $n_i + n_j$  is odd,  $n_i, n_j \geq 3$ ; (b)  $n_i \geq 3$ ,  $n_j \geq 16$ .

- (iii) (Korbaš) *Let  $s \geq 3$ . The flag manifold*

$$G(n_1, n_2, \dots, n_s) = O(n)/(O(n_1) \times \dots \times O(n_s))$$

is weakly almost complex if and only if  $n_1 = \dots = n_s = 1$ .  $M = G(1, \dots, 1)$  is almost complex if  $\dim M = \binom{n}{2}$  is even.

*Proof.* (i), (ii), and parts of (iii) follow from the observation that in each of the cases the manifold in question can be regarded as the total space of a differentiable bundle with fiber, an oriented Grassmann manifold, or a Grassmann manifold which is not weakly almost complex by Theorem 1.1 or 3.1 (cf. [13]).

Korbaš proves (iii) by a Stiefel-Whitney class argument. The positive results follow from the fact that  $G(1, \dots, 1)$  is parallelizable. See [15] for details.

### 3.4. Remarks.

- (i) M. Markl [10] has observed using his ‘ $J$ -genus’ that  $(\tilde{G}_{7,3})^n$  is not almost complex. He shows, also, that none of the quaternionic flag manifolds other than  $\mathbb{H}G(1, \dots, 1)$  admits a weakly almost complex structure, using the corresponding negative results of Hsiang and Szczarba [6] for quaternionic Grassmannians.
- (ii) J. Korbaš [8] has shown that  $\tilde{G}(n_1, \dots, n_s)$ ,  $s \geq 2$ ,  $n_1 \equiv \dots \equiv n_{s-1} \equiv 0 \pmod{4}$ ,  $n_2 \equiv 1 \pmod{2}$ , is not almost complex. For  $n_s \geq 3$ , this is weaker than the result of Theorem 3.3(ii), whereas our theorem does not cover completely the case  $n_s = 1$ .

### REFERENCES

1. J. F. Adams, *Lectures on Lie groups*, Benjamin, New York, 1969.
2. M. F. Atiyah and J. A. Todd, *On the complex Stiefel manifolds*, Proc. Cambridge Math. Soc. **56** (1960), 343–353.
3. V. Bartik and J. Korbaš, *Stiefel-Whitney characteristic classes and parallelizability of Grassmann manifolds*, Suppl. Rend. Circ. Mat. Palermo (2) **6** (1984), 19–29.
4. A. Borel and F. Hirzebruch, *Characteristic classes and homogeneous spaces-I*, Amer. J. Math. **80** (1958), 458–535.
5. A. Borel and J.-P. Serre, *Groupes de Lie et puissances reduites de Steenrod*, Amer. J. Math. **75** (1953), 409–448.
6. W. C. Hsiang and R. H. Szczarba, *On the tangent bundle of a Grassmann manifold*, Amer. J. Math. **86** (1964), 698–704.
7. D. Husemoller, *Fibre bundle*, Graduate Texts in Mathematics, no. 20, Springer-Verlag, New York, 1975.
8. J. Korbaš, *On the nonexistence of almost complex structures on some oriented flag manifolds*, Diff. Geom. and Appl., communications (Brno, 1986), 175–179, Univ. J. E. Purkyně, Brno, 1987.
9. K. Y. Lam, *A formula for the tangent bundle of flag manifolds and related manifolds*, Trans. Amer. Math. Soc. **213** (1975), 305–314.
10. M. Markl, *Note on the existence of almost complex structures on compact manifolds*, Ann. Glob. Anal. Geom. **4** (1986), 263–269.
11. W. S. Massey, *Non-existence of almost complex structures on quaternionic projective spaces*, Pacific J. Math. **12** (1962), 1379–1384.
12. J. W. Milnor and J. D. Stasheff, *Characteristic classes*, Ann. Math. Studies, vol. 76, Princeton Univ. Press, Princeton, NJ, 1974.
13. P. Sankaran and P. Zvengrowski, *Stable parallelizability of partially oriented flag manifolds*, Pacific J. Math. **128** (1987), 349–359.
14. N. Steenrod, *Topology of fibre bundles*, Princeton Univ. Press, Princeton, NJ, 1951.
15. J. Korbaš, *Note on Stiefel-Whitney classes of flag manifolds*, Suppl. Rend. Circ. Mat. Palermo (2) **16** (1987), 109–111.