

THE DUNFORD-PETTIS PROPERTY IN THE PREDUAL OF A VON NEUMANN ALGEBRA

L. J. BUNCE

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. The von-Neumann algebras whose predual has the Dunford-Pettis property are characterised as being Type I finite. This answers the question asked by Chu and Iochum in *The Dunford Pettis property in C^* -algebras*, *Studia Math.* **97** (1990), 59–64.

The purpose of this paper is to characterise those von Neumann algebras M for which the predual M_* has the Dunford-Pettis property thereby settling the question raised in [1].

A Banach space X is said to have the Dunford-Pettis property if for each Banach space Y each weakly compact linear operator from X to Y sends weakly convergent sequences to norm convergent sequences. Classically, all L_1 spaces (Dunford and Pettis) and all $C(X)$ spaces (Gröthendieck) have the Dunford-Pettis property (see [2] for much more). A thorough study of the Dunford-Pettis property in C^* -algebras and von Neumann algebras was undertaken in [1] and [3], to which we refer the reader for any unmentioned details. But one question remained unanswered. It was proved in [1] and [3] that if M is a von Neumann algebra then

- (a) if M_* has the Dunford-Pettis property then M is finite;
- (b) if M is Type I finite then M_* has the Dunford-Pettis property.

An obstacle to a characterisation was that it was not known whether it was possible for the predual of a Type II_1 von Neumann algebra to have the Dunford-Pettis property. We show that is it not possible. Thus we establish the following.

Theorem. *The following are equivalent for a von Neumann algebra M .*

- (i) M_* has the Dunford-Pettis property.
- (ii) M is Type I finite.

Proof. In order to establish the validity of the converse of (b) let M be a von Neumann algebra for which M_* has the Dunford-Pettis property. In view of (a), and since the predual of every summand of M clearly inherits the property, it can be supposed that M is of Type II_1 . At this point we appeal to the Jordan operator theory of spin factors contained in [4–6]. Since M contains a

Received by the editors October 31, 1990 and, in revised form, February 9, 1991.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 46L05, 46L10, 46B20.

© 1992 American Mathematical Society
0002-9939/92 \$1.00 + \$.25 per page

Type II_1 subfactor it certainly contains a countably infinite spin system $\{s_n\}$, consisting of nontrivial symmetries s_n (in M_{sa}) satisfying $s_n s_m + s_m s_n = 0$ whenever $n \neq m$. The real Banach subspace V of M_{sa} generated by $\{s_n\}$ is a JW-subalgebra of M_{sa} (called a spin factor). In addition, V is an infinite dimensional real Hilbert space in an equivalent norm. Evidently, then, there exists a sequence (x_n) in V such that $\|x_n\| = 1$ for all n and $x_n \rightarrow 0$ in the $\sigma(V, V^*)$ topology. Clearly $x_n \rightarrow 0$ in the $\sigma(M, M^*)$ topology. But as M_* has the Dunford-Pettis property this means that $x_n^2 \rightarrow 0$ in the $\sigma(M, M_*)$ topology, by [3, Lemma 1] or [1, Corollary 5]. Hence $x_n \rightarrow 0$ in the strong operator topology. But as proved in [5, Theorem 7.1] the latter coincides on V with the norm topology. So $x_n \rightarrow 0$ in norm, a contradiction which completes the proof.

REFERENCES

1. C.-H. Chu and B. Iochum, *The Dunford Pettis property in C^* -algebras*, *Studia Math.* **97** (1990), 59–64.
2. J. Diestel, *A survey of results related to the Dunford-Pettis property*, *Contemp. Math.*, vol. 2, Amer. Math. Soc., Providence, RI, 1980, pp. 15–60.
3. M. Hamana, *On linear topological properties of some C^* -algebras*, *Tôhoku Math. J.* **29** (1977), 157–163.
4. H. Hanche-Olsen and E. Stormer, *Jordan operator algebras*, Pitman, London, 1984.
5. E. Stormer, *Jordan algebras of type I*, *Acta Math.* **115** (1966), 165–184.
6. D. Topping, *An isomorphism invariant for spin factors*, *J. Math. Mech.* **15** (1966), 1055–1063.

DEPARTMENT OF MATHEMATICS, READING UNIVERSITY, P.O. BOX 220, READING RG6 2AX, ENGLAND