

A CHARACTERIZATION OF THE SECOND DUAL OF $C_0(S, A)$

STEPHEN T. L. CHOY AND JAMES C. S. WONG

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Let S be a locally compact Hausdorff space, and let A be a Banach space. The space of the continuous functions from S to A vanishing at infinity is denoted by $C_0(S, A)$. Let $MW(S, A^*)$ be the space of the representing measures of all the bounded linear functionals on $C_0(S, A)$. For $\mu \in MW(S, A^*)$ let

$$L_\infty(|\mu|, A^{**}, A^*) = \{f: S \rightarrow A^{**} \mid f(\cdot)x^* \in L_\infty(|\mu|) \forall x^* \in A^*\}.$$

The second dual of $C_0(S, A)$ is characterized in the general case by means of certain elements in the product linear space $\prod\{L_\infty(|\mu|, A^{**}, A^*) : \mu \in MW(S, A^*)\}$.

1. INTRODUCTION

Let A be a Banach space, S be a locally compact Hausdorff space, and $\mathcal{B}(S)$ be the σ -algebra of all the Borel sets of S . The space of the continuous functions from S to A vanishing at infinity endowed with the uniform norm is denoted by $C_0(S, A)$. The second dual of $C_0(S, A)$ is considered for the case where S is compact and the dual A^* has the Radon-Nikodym property in [3], for the case where S is locally compact and A is a Banach algebra with a positive cone satisfying certain conditions in [4]. Recently a characterization of $C_0^{**}(S, A)$ by means of the "generalized functions" was given in [5] by the authors in the case where A^* and A^{**} have the Radon-Nikodym property. The purpose of this paper is to give a characterization of $C_0^{**}(S, A)$ in general by means of the elements in $gl(S, A^{**})$ (see Definition 3.1).

Let $T: C_0(S, A) \rightarrow \mathbb{C}$ be a bounded linear functional, and let $M(S)$ be the space of all the bounded regular Borel measures on S . Then, since \mathbb{C} is reflexive, T is weakly compact. There is a unique representing measure $m: \mathcal{B}(S) \rightarrow A^*$ such that $m(\cdot)x \in M(S)$ for all $x \in A$, $T(f) = \int f dm$ for all $f \in C_0(S, A)$ and $\|T\| = \hat{m}(S)$, the semivariation of m (see, e.g., [2, 4]). Note that the total variation $|m|$ and \hat{m} are the same in this case [8, p. 54]. Let $MW(S, A^*)$ be the set of the representing measures m of all such functionals T . Since $|m|(S) = \hat{m}(S) = \|T\|$ is finite, we see that $|m| \in M(S)$ [2, Theorem

Received by the editors December 2, 1991 and, in revised form, May 4, 1992.

1991 *Mathematics Subject Classification.* Primary 46G10, 46E40; Secondary 46J10.

Key words and phrases. Second dual, vector-valued function space, Bochner integrable functions. The second author is deceased.

© 1993 American Mathematical Society
0002-9939/93 \$1.00 + \$.25 per page

2.8] and that

$$C_0^*(S, A) = MW(S, A^*).$$

For $\mu \in MW(S, A^*)$ let $L_1(|\mu|, A^*)$ be the space of all the equivalence classes of A^* -valued Bochner integrable functions defined on S . Measures defined by densities in $L_1(|\mu|, A^*)$ are studied in §2. It is shown in Theorem 2.2 that the linear span M of all such measures in $MW(S, A^*)$ is contained between $M(S) \otimes A^*$ and its closure $[M(S) \otimes A^*]^-$ with respect to the semivariation norm. The second dual $C_0^{**}(S, A)$ is then characterized in §3.

2. MEASURES DEFINED BY DENSITIES IN $L_1(|\mu|, A^*)$

Let $\varphi \in L_1(|\mu|, A^*)$, and define $\nu = \varphi|\mu| \in MW(S, A^*)$ by the Riesz Representation Theorem

$$\int g \, d\nu = \int \langle g, \varphi \rangle \, d|\mu| \quad (g \in C_0(S, A)).$$

It follows that, for $E \in \mathcal{B}(S)$,

$$\nu(E)x = \int \langle \chi_E x, \varphi \rangle \, d|\mu| \quad (x \in A)$$

and

$$(*) \quad \nu(E) = \int_E \varphi \, d|\mu|.$$

Also $|\nu| = |\varphi|\mu| = |\varphi| |\mu|$ by Dinculeanu [8, Theorem 6, p. 186]. Hence

$$(**) \quad \|\nu\| = |\nu|(S) = (|\varphi| |\mu|)(S) = \int |\varphi| \, d|\mu| = \|\varphi\|_{1, |\mu|}.$$

We see from (*) that $\nu \ll |\mu|$ (see [7, Definition 3, p. 11]). Therefore $\varphi \rightarrow \nu$ is a linear isometry from $L_1(|\mu|, A^*)$ into $\{\nu \in MW(S, A^*) : \nu \ll |\mu|\}$.

Definition 2.1. Denote by M the linear span in $MW(S, A^*)$ of all the measures defined by densities in $L_1(|\mu|, A^*)$ for all $\mu \in MW(S, A^*)$. That is,

$$M = \text{span} \left\langle \bigcup \{L_1(|\mu|, A^*)|\mu| : \mu \in MW(S, A^*)\} \right\rangle.$$

We shall show in Theorem 2.2 that this subspace M is large enough for us to apply the Hahn-Banach Extension Theorem to characterize $C_0^{**}(S, A)$. Let us recall first a result from Cambern and Grein [3, Corollary 2] that $M(S) \otimes A^*$ can be embedded in $MW(S, A^*)$ in such a way that $m \otimes x^*$ corresponds to $m(\cdot)x^*$ for all $m \in M(S)$ and $x^* \in A^*$.

Theorem 2.2. Let $[M(S) \otimes A^*]^-$ be the closure of $M(S) \otimes A^*$ with respect to the semivariation norm of $MW(S, A^*)$. Then

$$[M(S) \otimes A^*]^- \supseteq M \supseteq M(S) \otimes A^*.$$

Proof. We show first that $M \supseteq M(S) \otimes A^*$. Let $m \in M(S)$ and $x^* \in A^*$. Then, by the Radon-Nikodym Theorem for scalar measures, $m = \phi|m|$ for

some $\phi \in L_1(|m|)$. For any $\sum_{i=1}^N f_i \otimes x_i \in C_0(S) \otimes A$,

$$\begin{aligned} (m \otimes x^*) \left(\sum_1^N f_i \otimes x_i \right) &= \sum_1^N m(f_i)x^*(x_i) = \sum_1^N \phi|m|(f_i) \cdot x^*(x_i) \\ &= \sum_1^N \int \langle f_i x_i, \phi x^* \rangle d|m| = \sum_1^N (\phi x^*)|m|(f_i x_i) \\ &= (\phi x^*)|m| \left(\sum_1^N f_i \otimes x_i \right). \end{aligned}$$

Since $C_0(S, A) = C_0(S) \hat{\otimes} A$ with the least cross norm, we see that $m \otimes x^* = (\phi x^*)|m| \in M$, so $M \supseteq M(S) \otimes A^*$.

Next we shall prove that $(L_1(|\mu|) \otimes A^*)|\mu| \subset M(S) \otimes A^*$. Recall that $(L_1(|\mu|) \hat{\otimes} A^*) = L_1(|\mu|, A^*)$, where the closure on the left is with respect to the greatest cross norm (see, e.g., Diestel and Uhl [7, Example 10, p. 228]). For $\sum_{i=1}^N f_i \otimes x_i \in C_0(S) \otimes A$, $\sum_{j=1}^M (\phi_j \otimes x_j^*) \in L_1(|\mu|) \otimes A^*$,

$$\begin{aligned} \sum_{j=1}^M (\phi_j \otimes x_j^*)|\mu| \left(\sum_{i=1}^N f_i \otimes x_i \right) &= \int \left\langle \sum_{i=1}^N f_i \otimes x_i, \sum_{j=1}^M \phi_j \otimes x_j^* \right\rangle d|\mu| \\ &= \sum_{j=1}^M \int \left\langle \sum_{i=1}^N f_i \otimes x_i, \phi_j \otimes x_j^* \right\rangle d|\mu| = \sum_{j=1}^M \sum_{i=1}^N \int \langle f_i, \phi_j \rangle x_j^*(x_i) d|\mu| \\ &= \sum_{j=1}^M \sum_{i=1}^N \int \langle f_i, \phi_j \rangle d|\mu| \cdot x_j^*(x_i) = \sum_{j=1}^M \sum_{i=1}^N (\phi_j |\mu| \otimes x_j^*)(f_i \otimes x_i) \\ &= \sum_{j=1}^M (\phi_j |\mu| \otimes x_j^*) \left(\sum_{i=1}^N f_i \otimes x_i \right). \end{aligned}$$

By taking the limit process we see that

$$\sum_{j=1}^M (\phi_j \otimes x_j^*)|\mu| = \sum_{j=1}^M (\phi_j |\mu| \otimes x_j^*).$$

Thus

$$(L_1(|\mu|) \otimes A^*)|\mu| \subset M(S) \otimes A^* \subseteq MW(S, A^*).$$

Finally let $\varphi \in L_1(|\mu|, A^*)$, and let

$$\varphi_j = \sum_{i=1}^{N_j} \phi_i \otimes x_i^* \in L_1(|\mu|) \otimes A^*$$

be such that $\varphi_j \rightarrow \varphi$ in $L_1(|\mu|, A^*)$. For $f \in C_0(S, A)$, $\|f\| \leq 1$,

$$\begin{aligned} \left| \int f d\varphi_j |\mu| - \int f d\varphi |\mu| \right| &= \left| \int \langle f, \varphi_j - \varphi \rangle d|\mu| \right| \\ &= \left| \int f d(\varphi_j - \varphi) |\mu| \right| \leq \|f\| \cdot \|\varphi_j - \varphi\|_{1, |\mu|} \\ &\leq \|\varphi_j - \varphi\|_{1, |\mu|} \rightarrow 0. \end{aligned}$$

That is, $\varphi_j|\mu| \rightarrow \varphi|\mu|$ in $MW(S, A^*)$ with semivariation (= total variation) norm. Hence $M \subseteq [M(S) \otimes A^*]^-$. This completes the proof of the theorem.

Remark 2.3. It is worthwhile to note that if S is compact then $[M(S) \otimes A^*]^- = MW(S, A^*)$ if S is dispersed or A^* has the Radon-Nikodym property (see [3, Theorem 1]).

3. CHARACTERIZATION OF THE SECOND DUAL

For $\mu \in MW(S, A^*)$ let $L_\infty(|\mu|)$ be the Banach space of all the equivalence classes of bounded Borel measurable (real-valued) functions on S with essential supremum norm (see, e.g., [11, p. 85]). Denote

$$L_\infty(|\mu|, A^{**}, A^*) = \{f: S \rightarrow A^{**} | f(\cdot)x^* \in L_\infty(|\mu|) \forall x^* \in A^*\}.$$

For details of the definition of these spaces see [10, Definition 3, p. 75 and §5, p. 78]. Functions in $L_\infty(|\mu|, A^{**}, A^*)$ are identified by the equivalence relation $f \equiv g(w)$, which means by definition

$$f(\cdot)x^* = g(\cdot)x^* \quad (x^* \in A^*)$$

$|\mu|$ -a.e. (the null set can depend on x^*) [10, p. 76].

Note that functions in $L_\infty(|\mu|, A^{**}, A^*)$ need not be $|\mu|$ -measurable (hence not necessarily Bochner integrable) unless for example when A^{**} is separable [10, Corollary to Theorem 10, p. 73], in which case, $L_\infty(|\mu|, A^{**}, A^*) = L_\infty(|\mu|, A^{**})$, where $L_\infty(|\mu|, A^{**})$ stands for the space of the equivalence classes of A^{**} -valued Bochner integrable functions defined on S that are $|\mu|$ -essentially bounded.

The space $L_\infty(|\mu|, A^{**}, A^*)$ is a Banach space with norm

$$N_\infty(g) = \inf\{\alpha > 0 : \{s : |g(s)| > \alpha\} \text{ is } |\mu|\text{-null}\}$$

[10, p. 74]. Note that, by the uniform boundedness principle, we have

$$N_\infty(g) \leq \|g\| = \sup\{\|g(s)\| : s \in S\} \leq \infty$$

[10, p. 75].

It is also shown in [10, Chapter VII, §4, Theorem 7, and Corollary] that

$$L_1(|\mu|, A^*)^* \cong L_\infty(|\mu|, A^{**}, A^*)$$

is an isometric isomorphism under the correspondence

$$(g, \varphi) = \int \langle g(s), \varphi(s) \rangle d|\mu|(s)$$

for $\varphi \in L_1(|\mu|, A^*)$, $g \in L_\infty(|\mu|, A^{**}, A^*)$.

Definition 3.1. Consider the product linear space $\prod\{L_\infty(|\mu|, A^{**}, A^*) : \mu \in MW(S, A^*)\}$. Let $f = (f_\mu)_{\mu \in MW(S, A^*)}$ be an element in this space satisfying the following two conditions:

- (1) $\|f\|_\infty = \sup\{N_\infty(f_\mu) : \mu \in MW(S, A^*)\} < \infty$ and
- (2) if $\mu, \nu \in M(S, A^*)$ are such that $|\mu| \ll |\nu|$, then $f_\mu \equiv f_\nu(w)$ with respect to $|\mu|$.

The set of all such elements f on S is denoted by $\text{gl}(S, A^{**})$. With the norm $\|f\|_\infty$, $\text{gl}(S, A^{**})$ is a Banach space.

Lemma 3.2. *Let $\mu, \nu \in MW(S, A^*)$. If $|\mu| \ll |\nu|$ with $|\mu| = \omega|\nu|$ for some $\omega \in L_1(|\nu|)$ and $\varphi \in L_1(|\mu|, A^*)$, $\psi = \omega\varphi$, then $\varphi|\mu| = \psi|\nu|$.*

Proof. Since $|\mu| \ll |\nu|$, by the Radon-Nikodym Theorem, there is $\omega \in L_1(|\nu|)$ such that $|\mu| = \omega|\nu|$ and

$$\int g d|\mu| = \int g\omega d|\nu| \quad (g \in L_1(|\mu|)).$$

Fix $\varphi \in L_1(|\mu|, A^*)$ and let $\psi = \omega\varphi$. Then $\psi \in L_1(|\nu|, A^*)$. Let $f \in C_0(S, A)$

$$\begin{aligned} \int f d\varphi|\mu| &= \int \langle f, \varphi \rangle d|\mu| = \int \langle f, \varphi \rangle \omega d|\nu| \\ &= \int \langle f, \omega\varphi \rangle d|\nu| \quad (\omega \text{ is scalar-valued}) \\ &= \int \langle f, \psi \rangle d|\nu| = \int f d\psi|\nu|. \end{aligned}$$

Hence $\varphi|\mu| = \psi|\nu|$.

Theorem 3.3. *For each $F \in C_0^{**}(S, A)$, there is a (unique) $f \in \text{gl}(S, A^{**})$ such that*

$$F(\varphi|\mu) = \int \langle f_\mu, \varphi \rangle d|\mu| \quad (\varphi \in L_1(|\mu|, A^*))$$

for all $\mu \in MW(S, A^*)$.

Proof. For each $F \in C_0^{**}(S, A)$, consider the bounded linear functional

$$\varphi \rightarrow F(\varphi|\mu) \quad (\varphi \in L_1(|\mu|, A^*)).$$

From the result of Tulcea and Tulcea [10, Chapter VII, §4] mentioned earlier, there is $f_\mu \in L_\infty(|\mu|, A^{**}, A^*) = L_1(|\mu|, A^*)^*$ such that

$$F(\varphi|\mu) = \int \langle f_\mu, \varphi \rangle d|\mu| \quad (\varphi \in L_1(|\mu|, A^*)).$$

Hence we have an $f = (f_\mu)_{\mu \in MW(S, A^*)} \in \text{gl}(S, A^{**})$ if we can show properties (1) and (2) of Definition 3.1. Now

(1) For each $\mu \in MW(S, A^*)$, since $L_1(|\mu|, A^*)^* \cong L_\infty(|\mu|, A^{**}, A^*)$ is an isometric isomorphism and $\|\varphi|\mu|\| = \|\varphi\|_{1, |\mu|}$ (see (**)) in §2),

$$\begin{aligned} N_\infty(f_\mu) &= \sup \left\{ \left| \int \langle f_\mu, \varphi \rangle d|\mu| \right| : \|\varphi\|_{1, |\mu|} \leq 1 \right\} \\ &= \sup \left\{ \int |F(\varphi|\mu)| : \|\varphi\|_{1, |\mu|} \leq 1 \right\} \\ &\leq \sup \{ |F(\nu)| : \|\nu\| \leq 1 \} \\ &\leq \sup \{ \|F\| \|\nu\| : \|\nu\| \leq 1 \} \leq \|F\| < \infty. \end{aligned}$$

(2) Suppose $|\mu| \ll |\nu|$ and $\varphi \in L_1(|\mu|, A^*)$. By the Riesz Representation Theorem, there is $\omega \in L_1(|\nu|)$ such that $|\mu| = \omega|\nu|$. Let $\psi = \omega\varphi$. Then, from

Lemma 3.2. $\varphi|\mu| = \psi|\nu|$. Hence, for all $\varphi \in L_1(|\mu|, A^*)$,

$$\begin{aligned} \int \langle f_\mu, \varphi \rangle d|\mu| &= F(\varphi|\mu|) = F(\psi|\nu|) \\ &= \int \langle f_\nu, \psi \rangle d|\nu| = \int \langle f_\nu, \omega\varphi \rangle d|\nu| \\ &= \int \langle f_\nu, \varphi \rangle \omega d|\nu| = \int \langle f_\nu, \varphi \rangle d|\mu|. \end{aligned}$$

Hence $f_\mu \equiv f_\nu(w)$. Thus $f = (f_\mu) \in \text{gl}(S, A^{**})$.

The uniqueness is due to the fact that functions in $L_\infty(|\mu|, A^{**}, A^*)$ are identified by the equivalence relation $f \equiv g(w)$. This completes the proof of the theorem.

For $f_\mu \in L_\infty(|\mu|, A^{**}, A^*)$, define F_μ on $L_1(|\mu|, A^*)|\mu| \subseteq MW(S, A^*)$ by

$$F_\mu(\varphi|\mu|) = \int \langle f_\mu, \varphi \rangle d|\mu| \quad (\varphi \in L_1(|\mu|, A^*)).$$

Lemma 3.4. Let $\mu, \nu \in MW(S, A^*)$ with $|\mu| \ll |\nu|$. Then F_μ is linear and bounded and $F_\mu = F_\nu$ on $L_1(|\mu|, A^*)|\mu|$.

Proof. Clearly F_μ is linear and bounded because

$$\begin{aligned} \|F_\mu\| &= \sup\{|F_\mu(\varphi|\mu|)| : \|\varphi|\mu|\| \leq 1\} \\ &\leq \sup\left\{\left|\int \langle f_\mu, \varphi \rangle d|\mu|\right| : \|\varphi\|_{1,|\mu|} \leq 1\right\} \\ &\leq \sup\{N_\infty(f_\mu) \cdot \|\varphi\|_{1,|\mu|} : \|\varphi\|_{1,|\mu|} \leq 1\} \\ &\leq N_\infty(f_\mu) < \infty. \end{aligned}$$

Since $|\mu| \ll |\nu|$, there is some $\omega \in L_1(|\nu|)$ such that $|\mu| = \omega|\nu|$. Let $\psi = \omega\varphi$. Then

$$\begin{aligned} F_\nu(\psi|\nu|) &= \int \langle f_\nu, \psi \rangle d|\nu| = \int \langle f_\nu, \varphi \rangle \omega d|\nu| \\ &= \int \langle f_\nu, \varphi \rangle d|\mu| = \int \langle f_\mu, \varphi \rangle d|\mu| = F_\mu(\varphi|\mu|), \end{aligned}$$

since $f_\mu \equiv f_\nu(w)$ and $L_\infty(|\mu|, A^{**}, A^*) = L_1(|\mu|, A^*)^*$ is an isometry. Now, since $\psi|\nu| = \varphi|\mu|$ by Lemma 3.2,

$$F_\mu(\varphi|\mu|) = F_\nu(\psi|\nu|) = F_\nu(\varphi|\mu|).$$

Consequently, $F_\mu = F_\nu$ on $L_1(|\mu|, A^*)|\mu|$.

Theorem 3.5. For all $\mu \in MW(S, A)$ and $f = (f_\mu)_\mu \in \text{gl}(S, A^{**})$, there is an $F \in C_0^{**}(S, A)$, such that

$$F(\varphi|\mu|) = \int \langle f_\mu, \varphi \rangle d|\mu| \quad (\varphi \in L_1(|\mu|, A^*)).$$

Proof. We shall define a bounded linear functional F on the linear span M defined in Definition 2.1 such that

$$F(\varphi|\mu|) = \int \langle f_\mu, \varphi \rangle d|\mu|$$

for $\varphi \in L_1(|\mu|, A^*)$ and $\mu \in MW(S, A^*)$. In general, an element ϕ in M is of the form

$$\phi = \sum_{i=1}^m \varphi_i |\mu_i|,$$

where $\varphi_i \in L_1(|\mu_i|, A^*)$ and $\mu_i \in MW(S, A^*)$. Define

$$F(\phi) = \sum_{i=1}^m F_{\mu_i}(\varphi_i |\mu_i|).$$

We shall show that F is independent of the special representation of ϕ . Suppose also $\phi = \sum_{j=1}^n \psi_j |\nu_j|$. Pick $x^* \in A^*$ with $\|x^*\| = 1$, and put

$$\tau = \left(\sum_{i=1}^m |\mu_i| + \sum_{j=1}^n |\nu_j| \right) x^*.$$

Then clearly $\tau \in MW(S, A^*)$ and $|\mu_i| \ll |\tau|$, $|\nu_j| \ll |\tau|$ for $1 \leq i \leq m$, $1 \leq j \leq n$. Thus $|\mu_i| = \omega_i |\tau|$, $|\nu_j| = \eta_j |\tau|$ for some $\omega_i, \eta_j \in L_1(|\tau|)$. Thus

$$\begin{aligned} \phi &= \sum_{i=1}^m \varphi_i |\mu_i| = \sum_{i=1}^m \varphi_i \omega_i |\tau| \\ &= \left(\sum_{i=1}^m \varphi_i \omega_i \right) |\tau| \in L_1(|\tau|, A^*) |\tau| \end{aligned}$$

by Lemma 3.2. Let $\sum_{i=1}^m \varphi_i \omega_i = g$. Similarly, we have

$$\phi = \sum_{j=1}^n \psi_j |\nu_j| = h |\tau|$$

for some $h \in L_1(|\tau|, A^*)$. Since $\phi = g|\tau| = h|\tau|$, we see from (*) in §2 that $g = h|\tau|$ a.e. Since they are elements in $L_1(|\tau|, A^*)$, $g = h$. Now

$$\begin{aligned} \sum_{i=1}^m F_{\mu_i}(\varphi_i |\mu_i|) &= \sum_{i=1}^m \int \langle f_{\mu_i}, \varphi_i \rangle d|\mu_i| \\ &= \sum_{i=1}^m \int \langle f_{\tau}, \varphi_i \rangle d|\mu_i| \quad (|\mu_i| \ll |\tau|) \\ &= \sum_{i=1}^m \int \langle f_{\tau}, \varphi_i \rangle \omega_i d|\tau| = \sum_{i=1}^m \int \langle f_{\tau}, \varphi_i \omega_i \rangle d|\tau| \\ &= \int \left\langle f_{\tau}, \sum_{i=1}^m \int \varphi_i \omega_i \right\rangle d|\tau| = \int \langle f_{\tau}, g \rangle d|\tau| = F_{\tau}(g|\tau). \end{aligned}$$

With similar argument we also have

$$\sum_{j=1}^n F_{\nu_j}(\psi_j |\nu_j|) = F_{\tau}(h|\tau|).$$

Since $g = h$, we see

$$\sum_{i=1}^m F_{\mu_i}(\varphi_i|\mu_i|) = \sum_{j=1}^m F_{\nu_j}(\psi_j|\nu_j|).$$

Thus F is well defined on M .

Additivity of F on M follows from definition. Also, for $\lambda \in \mathbb{C}$,

$$\begin{aligned} F\left(\lambda \sum_{i=1}^n \varphi_i|\mu_i|\right) &= F\left(\sum_{i=1}^n \lambda \varphi_i|\mu_i|\right) \\ &= \sum_{i=1}^n F_{\mu_i}(\lambda \varphi_i|\mu_i|) = \lambda \sum_{i=1}^n F_{\mu_i}(\varphi_i|\mu_i|), \end{aligned}$$

so F is linear.

By definition

$$F(\varphi|\mu|) = \int \langle f_\mu, \varphi \rangle d|\mu| \quad (\varphi \in L_1(|\mu|, A^*)).$$

We shall show that F is bounded on M . Clearly

$$|F_\mu(\varphi|\mu|)| \leq N_\infty(f_\mu)\|\varphi\|_{1,|\mu|} = N_\infty(f_\mu)\|\varphi|\mu|\|.$$

So, for $f = (f_\mu) \in \text{gl}(S, A^{**})$,

$$\|F_\mu\| \leq N_\infty(f_\mu) \leq \|f\|_\infty < \infty.$$

That is, F_μ is bounded on $L_1(|\mu|, A^*)|\mu|$. Next suppose $\phi = \sum_{i=1}^m \varphi_i|\mu_i| \in M$. Let $\gamma = (\sum_{i=1}^m |\mu_i|)x^*$ for some $x^* \in A^*$, $\|x^*\| = 1$. Then $\gamma \in MW(S, A^*)$ and $|\mu_i| \ll |\gamma|$ for each i . Let $|\mu_i| = \omega_i|\gamma|$ for some $\omega_i \in L_1(|\gamma|)$, and let $\omega = \sum_{i=1}^m \varphi_i\omega_i$. Then $\phi = \omega|\gamma|$ and

$$|F(\phi)| = |F_\tau(\phi)| \leq \|F_\tau\| \|\phi\| \leq \|f\|_\infty \|\phi\|.$$

That is, F is bounded on M with $\|F\| \leq \|f\|_\infty$. It follows that F has a unique extension (also denoted by F) to the closed linear span of M with $\|F\| \leq \|f\|_\infty$ and then to $MW(S, A^*)^* = C_0^{**}(S, A)$ by the Hahn-Banach Extension Theorem (the extension need not be unique). This completes the proof of the theorem.

ACKNOWLEDGMENT

The author is greatly indebted to the referee for his helpful comments on the presentation of this paper.

REFERENCES

1. J. Batt and E.J. Berg, *Linear bounded transformations on the space of continuous functions*, J. Funct. Anal. **4** (1969), 215–239.
2. J. K. Brooks and P. W. Lewis, *Linear operators and vector measures*, Trans. Amer. Math. Soc. **192** (1974), 139–162.
3. M. Cambern and P. Grein, *The bidual of $C(X, E)$* , Proc. Amer. Math. Soc. **85** (1982), 53–58.
4. S. T. L. Choy, *Positive operators and algebras of dominated measures*, Rev. Roumaine Math. Pures Appl., vol. 34, Ed. Acad. R. S. România, Bucharest, 1989, pp. 213–219.

5. S. T. L. Choy and J. C. S. Wong, *The second dual of $C_0(S, A)$* , J. Austral. Math. Soc. (to appear).
6. J. Diestel, *Sequences and series in Banach spaces*, Graduate Texts in Math., vol. 92, Springer-Verlag, New York, 1984.
7. J. Diestel and J. J. Uhl, *Vector measures*, Math. Surveys Monographs, vol. 15, Amer. Math. Soc., Providence, RI, 1977.
8. N. Dinculeanu, *Vector measures*, Pergamon Press, New York, 1967.
9. J. Duncan and S. A. R. Hosseini, *The second dual of a Banach algebra*, Proc. Roy. Soc. Edinburgh Sect. A **84** (1979), 309–325.
10. A. I. Tulcea and C. I. Tulcea, *Topics in the theory of lifting*, Springer-Verlag, Heidelberg and New York, 1969.
11. J. C. Wong, *Abstract harmonic analysis of generalized functions on locally compact semi-groups with applications to invariant means*, J. Austral. Math. Soc. Ser. A **23** (1977), 84–94.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, 10 KENT RIDGE CRES-
CENT, SINGAPORE 0511, REPUBLIC OF SINGAPORE

E-mail address: matct1@nusvm.bitnet

DEPARTMENT OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF CALGARY, CALGARY,
ALBERTA, CANADA T2N 1N4