

A FORBIDDEN SET FOR EMBEDDED EIGENVALUES

RAFAEL RENE DEL RIO CASTILLO

(Communicated by Andrew Bruckner)

ABSTRACT. We study the problem of embedding eigenvalues to the spectrum of a Sturm-Liouville operator in the half axis when this spectrum is a perfect set. We prove the existence of an uncountable dense subset of the spectrum for which, by modifying the condition at the left or by locally perturbing the potential, it is not possible to add any eigenvalues.

1. INTRODUCTION

In this paper we consider Sturm-Liouville operators generated by the differential expression $lu = -u'' + q(x)u$ in the half line $[0, \infty)$.

It is known that, through local perturbations of the potential or by considering arbitrary conditions at the left, it is possible to add eigenvalues anywhere in the resolvent set (see [4, Theorem 2.5.3]) or to produce an infinite number of embedded eigenvalues (see [2, Remark 5]).

Nevertheless, we prove in this paper that there exists a specific subset of the spectrum for which, assuming the spectrum of the unperturbed operator is a perfect set, it is not possible to generate embedded eigenvalues by means of the above perturbations. We also show that this “forbidden” set, which depends only on the behaviour of the potential at infinity, is a dense and uncountable set. Moreover, every point of the spectrum is a condensation point of this set.

This paper is organized as follows. In §2 we define the unperturbed operator L and the perturbed operator \tilde{L} . We prove that eigenvalues can appear only in points where the symmetric derivative of the spectral function of the unperturbed operator is zero. This is proved by using a theorem of Aronszajn [1]. In §3, using tools of elementary real analysis, we show that the set of points where a given series diverges is “big” in some sense. This result is crucial for proving our main theorem in the presence of only pure point spectrum. Section 4 is devoted to the proof of the main result where a theorem of Kundu [5] is applied.

Received by the editors July 31, 1992; presented to the Midwest and Southeastern Atlantic Second Joint Regional Conference on Differential Equations, November 1992, Lexington, Kentucky.
1991 *Mathematics Subject Classification*. Primary 34L99, 34B24, 47E05.

2. PRELIMINARIES

Consider the selfadjoint operator L generated by the differential expression

$$lu = -u'' + q(x)u, \quad 0 \leq x < \infty,$$

where q is a real-valued, locally integrable function, as $Lu = lu$ with domain $D(L) = \{u \in L_2(0, \infty) | u, u' \text{ are locally absolutely continuous, } lu \in L_2(0, \infty), \text{ and } u(0) \cos \alpha + u'(0) \sin \alpha = 0\}$, $\alpha \in [0, \pi)$.

The limit point case occurs at ∞ , and 0 is a regular point. We shall denote the spectral function of L by ρ . Sometimes to emphasize the dependence on α we shall write L_α and ρ_α .

The perturbed operator \tilde{L} will be any selfadjoint realization of the differential expression

$$\tilde{l}u = -u'' + \{q(x) + v(x)\}u, \quad a \leq x < \infty,$$

where $-\infty \leq a$ and $v(x)$ is a locally integrable function with compact support. If the limit circle case (l.c.c.) occurs at a , then a boundary condition will be needed.

The operator \tilde{L} is defined as $\tilde{L}u = \tilde{l}u$ with domain

$$D(\tilde{L}) = \{u \in L_2(a, \infty) | u, u' \text{ are locally absolutely continuous,}$$

$$\tilde{l}u \in L_2(a, \infty), \text{ and } [v, u]_a = 0 \text{ if we have l.c.c. at } a\}.$$

Here v is a nontrivial solution of $(\tilde{l} - \lambda)u = 0$ ($\lambda \in \mathbb{R}$) and

$$[v, u]_a = \lim_{x \rightarrow a} (\bar{v}(x)u'(x) - \bar{v}'(x)u(x)).$$

See, for example, Theorem 5.8 of [6].

The following observation will be useful in the sequel.

Remark 1. If λ is eigenvalue of \tilde{L} then, for some $\alpha \in [0, \pi)$, λ is an eigenvalue of L_α .

Let us define

$$D\rho(\lambda) = \lim_{\varepsilon \rightarrow 0} \frac{\rho(\lambda + \varepsilon) - \rho(\lambda - \varepsilon)}{2\varepsilon}.$$

Lemma 1. $\int_{-\infty}^{\infty} d\rho(\lambda)/(\lambda_0 - \lambda)^2 < \infty \Rightarrow D\rho(\lambda_0) = 0$.

Proof. Assume that $\int_{-\infty}^{\infty} d\rho(\lambda)/(\lambda_0 - \lambda)^2 < \infty$, and, for all $\varepsilon > 0$, define the interval $I_\varepsilon = [\lambda_0 - \varepsilon, \lambda_0 + \varepsilon]$. We shall denote the length of I_ε by $|I_\varepsilon|$. Then we have

$$\int_{-\infty}^{\infty} \frac{d\rho(\lambda)}{(\lambda_0 - \lambda)^2} \geq \int_{I_\varepsilon} \frac{d\rho(\lambda)}{(\lambda_0 - \lambda)^2} \geq \frac{1}{|I_\varepsilon|^2} \int_{I_\varepsilon} d\rho(\lambda) = \frac{1}{|I_\varepsilon|} \frac{\rho(I_\varepsilon)}{|I_\varepsilon|}$$

and, therefore,

$$|I_\varepsilon| \int_{-\infty}^{\infty} \frac{d\rho(\lambda)}{(\lambda_0 - \lambda)^2} \geq \frac{\rho(\lambda_0 + \varepsilon) - \rho(\lambda_0 - \varepsilon)}{2\varepsilon}.$$

Since

$$\lim_{\varepsilon \rightarrow 0} |I_\varepsilon| \int_{-\infty}^{\infty} \frac{d\rho(\lambda)}{(\lambda_0 - \lambda)^2} = 0$$

and the right-hand side of the above inequality is nonnegative, it follows that $D\rho(\lambda_0)$ exists and equals zero. Q.E.D.

Lemma 2. *If λ_0 is an eigenvalue of \tilde{L} then $D\rho(\lambda_0) = 0$ or λ_0 is an eigenvalue of L .*

Proof. Assume that λ_0 is an eigenvalue of \tilde{L} and $D\rho(\lambda_0) > 0$. Then from Lemma 1 it follows that $\int_{-\infty}^{\infty} d\rho(\lambda)/(\lambda_0 - \lambda)^2$ is not finite. Using Theorem 4 of [1] we can conclude that λ_0 is not an eigenvalue of L_β for $\beta \neq \alpha$. If λ_0 is not an eigenvalue of L_α then, from Remark 1, it follows that λ_0 is not an eigenvalue of \tilde{L} . Q.E.D.

3. A REAL VARIABLE RESULT

Let \mathbb{N} denote the set of positive integers, \mathbb{R}^+ the positive real numbers, and \bar{I} the closure of I . The difference of two sets A, B will be denoted by $A - B$. Recall that a set is said to be *dense in itself* if it is contained in the set of its limit points and that a point p in a metric space X is said to be a *condensation point* of a set $A \subset X$ if every neighborhood of p contains uncountable many points of A .

Let $F: S \rightarrow \mathbb{R}^+$ be an arbitrary function, where $S \subset [a, b]$ is countable and dense in itself. Define

$$A = \left\{ \xi \in [a, b] - S \mid \sum_{s \in S} \frac{F(s)}{|s - \xi|^2} = \infty \right\}.$$

Lemma 3. *The set A defined above is uncountable, and every point of S is a condensation point of A .*

Proof. Let us consider the family of functions $\mathcal{F} := \{\delta \mid \delta: \mathbb{N} \rightarrow \{0, 1\}\}$. Choose $\delta \in \mathcal{F}$, and, for each $k \in \mathbb{N}$, denote by δ_k the restriction of δ to the finite set $\{1, \dots, k\}$. We shall define for each δ_k an interval I_{δ_k} .

In what follows I_{sF} will denote the open interval $(s - F^{1/2}(s), s + F^{1/2}(s))$, and the index i will take the values $0, 1$.

Let $I \subset [a, b]$ be an arbitrary open interval such that $S \cap I \neq \emptyset$, and choose $s_0 \in S \cap I$. Define $I_{s_0} := I \cap I_{s_0F}$. Select and fix two different points a_i such that $a_i \neq s_0$ and $a_i \in I_{s_0} \cap S$. We can do this because S is dense in itself.

Let I_{a_i} be two open intervals which satisfy $a_i \in I_{a_i}$, $I_{a_i} \subset I_{s_0}$, $\bigcap_{i=0}^1 \bar{I}_{a_i} = \emptyset$. Define $I_{\delta_1} := I_{a_{\delta(1)}} \cap I_{a_{\delta(1)}F}$. This completes the construction for the case $k = 1$. Now define $s_1 = a_{\delta(1)}$.

Let us assume that we have defined an interval $I_{\delta_{k-1}}$ and a point $s_{k-1} \in I_{\delta_{k-1}} \cap S$. Choose two points a_i such that $a_i \in I_{\delta_{k-1}} \cap S$ and $a_i \neq s_{k-1}$, and fix two open intervals I_{a_i} satisfying $a_i \in I_{a_i}$, $I_{a_i} \subset I_{\delta_{k-1}}$, $\bigcap_{i=0}^1 \bar{I}_{a_i} = \emptyset$. Define $I_{\delta_k} := I_{a_{\delta(k)}} \cap I_{a_{\delta(k)}F}$ and $s_k = a_{\delta(k)}$.

We can now define I_{δ_k} for each $k \in \mathbb{N}$. This definition can be done in many ways, but once we have chosen the points a_i and the intervals I_{a_i} , the definition is unique. In order to define $I_{\delta'_k}$ for any other function δ' , we choose exactly the same points a_i and the same intervals I_{a_i} we have chosen to construct I_{δ_k} .

Now, for each $\delta \in \mathcal{F}$, define $B_I(\delta) = \bigcap_{k=1}^{\infty} \bar{I}_{\delta_k}$ where \bar{I}_{δ_k} denotes the closure of the interval I_{δ_k} . Since $\bar{I}_{\delta_{k-1}} \supset \bar{I}_{\delta_k}$ for every $k \in \mathbb{N}$, it follows that $B_I(\delta) \neq \emptyset$.

Now define $B_I = \bigcup_{\delta \in \mathcal{F}} B_I(\delta)$ and $B = \bigcup B_I$ where the second union is taken over all open intervals contained in $[a, b]$ such that $I \cap S \neq \emptyset$.

We shall prove the following:

- (a) $B - S \subset A$.
- (b) $B - S$ is uncountable and every point of S is a condensation point of $B - S$.

Choose $\xi \in B - S$. Then $\xi \in B_I(\delta) = \bigcap_{k=1}^{\infty} \bar{I}_{\delta_k}$ for some δ and I . Hence, there is a sequence $s_k \in I_{\delta_k} \cap S$ such that, for all $k \in \mathbb{N}$, $|\xi - s_k| \leq F^{1/2}(s_k)$. Therefore,

$$1 \leq \frac{F(s_k)}{|\xi - s_k|^2} \quad \text{and} \quad \sum_{s \in S} \frac{F(s)}{|s - \xi|^2} \geq \sum_{k=1}^{\infty} \frac{F(s_k)}{|s_k - \xi|^2} = \infty,$$

implying that $\xi \in A$. This shows that (a) holds.

To prove (b), choose an open interval $I \subset [a, b]$ and construct B_I as before, i.e.,

$$B_I = \bigcup_{\delta \in \mathcal{F}} B_I(\delta).$$

Since $B_I(\delta) \neq \emptyset$, select $x_\delta \in B_I(\delta)$ and define the function

$$J: \mathcal{F} \rightarrow B_I \\ \delta \rightarrow x_\delta \in B_I(\delta).$$

We shall see that J is injective. If $\delta, \delta' \in \mathcal{F}$ and $\delta \neq \delta'$, then there is $k \in \mathbb{N}$ such that $\delta(k) \neq \delta'(k)$. From the construction above it follows that $\bar{I}_{a_{\delta(k)}} \cap \bar{I}_{a_{\delta'(k)}} = \emptyset$ and therefore $\bar{I}_{\delta_k} \cap \bar{I}_{\delta'_k} = \emptyset$. Hence, $B_I(\delta) \cap B_I(\delta') = \emptyset$ and so $x_\delta \neq x_{\delta'}$.

Since the set \mathcal{F} is uncountable, so are B_I and $B - S$. Since $B_I \subset I$, it follows that each point of S is a condensation point of A . Thus (b) holds, and the proof of the lemma is complete. Q.E.D.

4. THE MAIN RESULT

In [5] Kundu proved that if:

- (i) $\liminf_{x \rightarrow \xi - 0} f(x) \geq f(\xi) \geq \liminf_{x \rightarrow \xi + 0} f(x)$ for all $\xi \in [a, b]$,
- (ii) $\underline{D}f(x) \leq 0$ almost everywhere in (a, b) ,
- (iii) $\overline{D}f(x) < \infty$ except of a countable set in (a, b) ,

then f is decreasing in $[a, b]$.

Here we used the notation

$$\overline{D}f(x) = \limsup_{h \rightarrow 0} \frac{f(x+h) - f(x-h)}{2h}, \\ \underline{D}f(x) = \liminf_{h \rightarrow 0} \frac{f(x+h) - f(x-h)}{2h}.$$

The function f is said to be *decreasing* if $x < y \Rightarrow f(x) \geq f(y)$. Analogously, f is *increasing* if $x < y \Rightarrow f(x) \leq f(y)$.

We shall use this result to prove the following theorem. The spectrum of L will be denoted by $\sigma(L)$.

Theorem. *If for an interval J the set $C = J \cap \sigma(L)$ is a perfect set, then there exists an uncountable set $B \subset C$ such that every point of C is a condensation point of B and, moreover, \tilde{L} cannot have eigenvalues in B .*

Proof. The spectral function of L can be written as $\rho = \rho_c + \rho_d$ where ρ_c is an increasing continuous function and ρ_d is an increasing saltus function. Consider an arbitrary open interval such that $I \cap C \neq \emptyset$.

If ρ_c is not constant in I then ρ_c is not decreasing and, applying the theorem of Kundu mentioned above, it follows that there is an uncountable set $B \subset I$ such that, if $x \in B$, then it is not possible to have $D\rho_c(x) = 0$. Since $D\rho(x) = 0$ implies $D\rho_c(x) = 0$, it follows that the relation $D\rho(x) = 0$ is not possible. In view of Lemma 2, the two last conclusions of the theorem follow in this case. It remains to show that $B \subset C$, but this is a simple consequence of the fact that for x in the resolvent set of L we have $D\rho(x) = 0$.

Now assume that ρ_c is constant in I . In this case the spectrum is pure point in I and, for any $\xi \in I$, we have

$$\int_{-\infty}^{\infty} \frac{d\rho(\lambda)}{(\xi - \lambda)^2} \geq \int_I \frac{d\rho(\lambda)}{(\xi - \lambda)^2} = \int_I \frac{d\rho_d}{(\xi - \lambda)^2}.$$

Since ρ_d is a saltus function, the measure generated by this function is supported on a countable set $S \subset I$.

Let us denote the measure of a point $s \in S$ by $F(s)$. Then we have

$$\int_I \frac{d\rho_d(\lambda)}{(\xi - \lambda)^2} = \sum_{s \in S} \frac{F(s)}{(\xi - s)^2}.$$

Since $\bar{I} \cap C = \bar{S}$ and C is perfect, S is dense in itself and we can apply Lemma 3, showing that for every ξ in an uncountable set $A \subset I$ it happens that

$$\int_{-\infty}^{\infty} \frac{d\rho(\lambda)}{(\xi - \lambda)^2} = \infty.$$

An application of Theorem 4 of [1] and Remark 1 imply now the two last conclusions of theorem. From

$$\sum_{s \in S} F(s) = \int_I d\rho(s) < \infty \quad \text{and} \quad \sum_{s \in S} \frac{F(s)}{(\xi - s)^2} = \infty$$

for $\xi \in A$, it follows that $\inf_{s \in S} |\xi - s| = 0$ and therefore $A \subset C$. This can also be proven using Theorem 2.5.3 of [4]. Therefore, the remaining assertion is proven. Q.E.D.

Remark 2. The theorem holds for every perturbation \tilde{L} which has the property mentioned in Remark 1.

Remark 3. Using Theorem 1 of Donoghue [3] instead of Theorem 4 of Aronszajn [1], a similar result can be obtained for perturbations $H_\alpha = H_0 + \alpha P$ of a selfadjoint operator H_0 , where P is a selfadjoint projection on a fixed normalized element and $\alpha \in \mathbb{R}$.

Remark 4. If $\lambda \in B$ then $lu = \lambda u$ has no solution which lies in L_2 near ∞ .

ACKNOWLEDGMENT

I thank Professor Don Hinton for a stimulating conversation and the referee for useful comments.

REFERENCES

1. N. Aronszajn, *On a problem of Weyl in the theory of singular Sturm-Liouville equations*, Amer. J. Math. **79** (1957), 597–610.
2. R. R. del Rio Castillo, *Embedded eigenvalues of Sturm-Liouville operators*, Comm. Math. Phys. **142** (1991), 421–431.
3. W. F. Donoghue, *On the perturbation of spectra*, Comm. Pure Appl. Math. **18** (1965), 559–579.
4. M. S. P. Eastham and H. Kalf, *Schrödinger type operators with continuous spectra*, Pitman Research Notes in Math., vol. 65, Longman Sci. Tech., Harlow, 1982.
5. N. K. Kundu, *On some properties of symmetric derivatives*, Ann. Polon. Math. **30** (1974), 9–18.
6. J. Weidmann, *Spectral theory of ordinary differential operators*, Lecture Notes in Math., vol. 1258, Springer-Verlag, New York, 1987.

DEPARTAMENTO DE MÉTODOS MATEMÁTICOS Y NUMÉRICOS, IIMAS-UNAM, APARTADO POSTAL
20-726, MÉXICO, D. F., 01000, MÉXICO
E-mail address: delrio@redvax1.dgsca.unam.mx